



Fisheries New Zealand

Tini a Tangaroa

Fisheries Assessment Plenary

May 2018

Stock Assessment and Stock Status

Volume 1: Introductory Section to Groper



Fisheries Assessment Plenary May 2018
Stock Assessments and Stock Status

Volume 1: Introductory Sections to Groper



Fisheries New Zealand

Tini a Tangaroa

Fisheries Science and Information

Fisheries Assessment Plenary

May 2018

Stock Assessments and Stock Status
Volume 1: Introductory section to Groper

ISBN (print): 978-1-77665-573-1
ISBN (online): 978-1-77665-572-4

© Crown Copyright May 2018 – Ministry for Primary Industries

The written material contained in this document is protected by Crown copyright. This document is published by Fisheries New Zealand, a branded business unit within the Ministry for Primary Industries. All references to Fisheries New Zealand in this document should therefore, be taken to refer to the Ministry for Primary Industries.

The information in this publication is not governmental policy. While all reasonable measures have been made to ensure the information is accurate, the Ministry for Primary Industries does not accept any responsibility or liability for any error, inadequacy, deficiency, flaw in or omission from the information provided in this document or any interpretation or opinion that may be present, nor for the consequences of any actions taken or decisions made in reliance on this information. Any view or opinion expressed does not necessarily represent the view of the Ministry for Primary Industries.

Compiled and published by
Fisheries New Zealand
Fisheries Science and Information
Pastoral House, 25 The Terrace
PO Box 2526, Wellington 6140
New Zealand

Requests for further copies should be directed to:
Publications Logistics Officer
Ministry for Primary Industries
PO Box 2526
WELLINGTON 6140
Email: brand@mpi.govt.nz
Telephone: 0800 00 83 33
Facsimile: 04-894 0300

This publication is also available on the Ministry for Primary Industries websites at:
www.mpi.govt.nz/news-resources/publications.aspx

Or at
fs.fish.govt.nz under document library and stock assessment plenary.

Cover images: Scampi – William D. Gibson

Printed by: Graphic Press, Wellington

Preferred citation

Fisheries New Zealand (2018). Fisheries Assessment Plenary, May 2018: stock assessments and stock status. Compiled by the Fisheries Science and Information Group, Fisheries New Zealand, Wellington, New Zealand. 1 660p

PREFACE

Fisheries Assessment Plenary reports have represented a significant annual output of Fisheries New Zealand (FNZ) and its predecessors for the last 34 years. The Plenary is now more than 2000 pages long and is split into four volumes, three of which are produced in May and one in November of each year. The Plenary reports provide summaries of the available information and are in turn supported by 70–100 more detailed reports published on-line each year.

The May 2018 Plenary summarises fisheries, biological, environmental, stock assessment and stock status information for 84 of New Zealand's commercial fish species or species groups in a series of Science Working Group (SWG) or Plenary reports. Each species or species group is split into 1–10 stocks for management purposes. In addition, the mid-year Plenary that is produced each November for species that operate on different management cycles includes 17 SWG and Plenary summaries for highly migratory species, rock lobster, scallops and dredge oysters.

Over time, continual improvements have been made in data acquisition, stock assessment techniques, the development of reference points to guide fisheries management decisions, the provision of increasingly comprehensive and meaningful information from a range of sources, and peer review processes. SWG and Plenary meetings have continued the effort to populate the Status of the Stocks summary tables, which are used to provide comprehensive summary information about current stock status and the prognosis for these stocks, to evaluate fisheries performance relative to the 2008 Harvest Strategy Standard for New Zealand Fisheries and other management measures, and to rank the quality of stock assessment inputs and outputs based on the 2011 Research and Science Information Standard for New Zealand Fisheries.

Over the past few years, sections on environmental and ecosystem considerations have also been developed for some species by the SWGs that oversee aquatic environment and biodiversity. Sections on how ocean warming, ocean acidification and other ecosystem trends affect, for example, productivity and fish distributions will be incorporated as new information becomes available. FNZ recognises the need to increase our knowledge of the impacts of important environmental factors.

The Plenary reports take into account the most recent data and analyses available to SWGs and Fisheries Assessment Plenary meetings, and also incorporate relevant analyses undertaken in previous years. Due to time and resource constraints, recent data for some stocks may not yet have been fully analysed by the SWGs or the Plenary.

I would like to recognise and thank the large number of research providers and scientists from research organisations, academia, the seafood industry, marine amateur fisheries, environmental NGOs, Maori customary and FNZ; along with all other technical and non-technical participants in present and past SWG and Plenary meetings for their substantial contributions to this report. My sincere thanks to each and all who have contributed.

I would also like to pay particular tribute to the FNZ's past and present Science Officers who put tireless effort into checking and collating each Plenary report. The Science Officer for this report was William D. Gibson.

I am pleased to endorse this document as representing the best available scientific information relevant to fisheries and stock status, as at 31 May 2018.



Dr Pamela Mace
Principal Advisor Fisheries Science
Fisheries New Zealand



Fisheries New Zealand

Tini a Tangaroa

MAY 2018 PLENARY VOLUME CONTENTS

Volume 1

Alfonsino to Groper

Alfonsino (BYX)
Anchovy (ANC)
Arrow squid (SQU)
Barracouta (BAR)
Black cardinalfish (CDL)
Bladder kelp attached (KBB G)
Blue cod (BCO)
Blue mackerel (EMA)
Blue moki (MOK)
Blue warehou (WAR)
Bluenose (BNS)
Butterfish (BUT)
Cockles (COC)
 COC Introduction
 COC 1A
 COC 3
 COC 7A
Deepwater (King) clam (PZL)
Elephant fish (ELE)
Flatfish (FLA)
Freshwater eels (SFE, LFE)
Frostfish (FRO)
Garfish (GAR)
Gemfish (SKI)
Ghost shark
 Dark ghost shark (GSH)
 Pale ghost shark (GSP)
Giant spider crab (GSC)
Green-lipped mussel (GLM)
Grey mullet (GMU)
Groper (HPB)

Volume 2

Hake to Pilchard

Hake (HAK)
Hoki (HOK)
Horse mussel (HOR)
Jack mackerels (JMA)
John dory (JDO)
Kahawai (KAH)
Kina (SUR)
King crab (KIC)
Kingfish (KIN)
Knobbed whelk (KWH)
Leatherjacket (LEA)
Ling (LIN)
Lookdown dory (LDO)
Orange roughy (ORH)
 ORH Introduction
 ORH 1
 ORH 2A/2B/3A
 ORH 3B
 ORH 7A
 ORH 7B
 ORH ET
Oreos (OEO)
 OEO Introduction
 OEO 3
 OEO 4
 OEO 1 and 6
Paddle crabs (PAD)
Parore (PAR)
Paua (PAU)
 Paua Introduction
 PAU 2
 PAU 3
 PAU 4
 PAU 5A
 PAU 5B
 PAU 5D
 PAU 7
Pilchard (PIL)

Volume 3

Pipis to Yellow-eyed Mullet

Pipis (PPI)
 PPI 1
 PPI 1A
Porae (POR)
Prawn killer (PRK)
Queen scallops (QSC)
Redbait (RBT)
Red cod (RCO)
Red crab (CHC)
Red gurnard (GUR)
Red snapper (RSN)
Ribaldo (RIB)
Rig (SPO)
Rubyfish (RBY)
Scampi (SCI)
Sea cucumber (SCC)
Sea perch (SPE)
Silver warehou (SWA)
Skates
 Rough Skate (RSK)
 Smooth Skate (SSK)
Snapper (SNA)
Southern blue whiting (SBW)
Spiny dogfish (SPD)
Sprat (SPR)
Stargazer (STA)
Surf Clams
 Surf Clams Introduction
 Deepwater tuatua (PDO)
 Fine (Silky) dosinia (DSU)
 Frimled venus shell (BYA)
 Large trough shell (MMI)
 Ringed dosinia (DAN)
 Triangle shell (SAE)
 Trough shell (MDI)
Tarakihi (TAR)
Toothfish (TOT)
Trevally (TRE)
Trumpeter (TRU)
Tuatua (TUA)
White warehou (WWA)
Yellow-eyed mullet (YEM)



CONTENTS

Volume 1: Introductory section to Groper

	Page
Introduction.....	1
Glossary.....	3
Terms of Reference for Fisheries Assessment Working Groups.....	13
Fisheries Assessment Working Groups: Membership 2016–17.....	26
Guide to Biological Reference Points for Fisheries Assessment Meetings.....	29
Guidelines for Status of the Stocks Summary Tables.....	37
Alfonsino (BYX).....	45
Anchovy (ANC).....	55
Arrow squid (SQU).....	59
Barracouta (BAR).....	77
Black cardinalfish (CDL).....	97
Bladder kelp attached (KBB G)	115
Blue cod (BCO).....	129
Blue mackerel (EMA).....	171
Blue moki (MOK).....	183
Blue warehou (WAR).....	193
Bluenose (BNS).....	201
Butterfish (BUT).....	223
Cockles (COC)	
COC Introduction	231
COC 1A.....	237
COC 3.....	247
COC 7A.....	259
Deepwater (King) clam (PZL)	269
Elephant fish (ELE)	275
Flatfish (FLA)	299
Freshwater eels (SFE, LFE).....	331
Frostfish (FRO)	413
Garfish (GAR)	423
Gemfish (SKI)	429
Ghost shark	
Dark ghost shark (GSH).....	449
Pale ghost shark (GSP).....	465
Giant spider crab (GSC).....	475
Green-lipped mussel (GLM).....	479
Grey mullet (GMU).....	483
Groper (HPB)	489

Introduction

This report summarises the conclusions and recommendations from the meetings of the Fisheries Assessment Working Groups and the Fisheries Assessment Plenary held since last year's Plenary report was published. The meetings were convened to assess the fisheries managed within the Quota Management System, as well as other important fisheries in the New Zealand EEZ, and to discuss various matters that pertain to fisheries assessments.

In addition, summaries of environmental effects of fishing from research presented to the Aquatic Environment Working Group (AEWG) that have relevance to fishery management have been incorporated for selected species. Paragraph 11 (page 15) of the Terms of Reference for Fisheries Assessment Working Groups (FAWGs) includes "...information and advice on other management considerations (e.g., ...by-catch issues, effects of fishing on habitat...)", and states that "Sections of the Working Group reports related to bycatch and other environmental effects of fishing will be reviewed by the Aquatic Environment Working Group although the relevant FAWG is encouraged to identify to the AEWG Chair any major discrepancies between these sections and their understanding of the operation of relevant fisheries". In addition, the Terms of Reference for the AEWG (Paragraph 8, page 18) specifies the need "to review and revise existing environmental and ecosystem consideration sections of Fisheries Assessment Plenary report text based on new data or analyses, or other relevant information".

The report addresses, for each species, relevant aspects of the Fisheries Act 1996 and related considerations, as defined in the Terms of Reference for Fisheries Assessment Working Groups for 2018. In all cases, consideration has been based on and limited by the best available information. The purpose has been to provide objective, independent assessments of the current status of the fish stocks.

There are two types of catch limits used in this document – total allowable catch (TAC) and total allowable commercial catch (TACC). The current definition is that a TAC is a limit on the total removals from the stock, including those taken by the commercial, recreational and customary non-commercial sectors, illegal removals and all other mortality to a stock caused by fishing. A TACC is a limit on the catch taken by the commercial sector only. The definition of TAC was changed in the 1990 Fisheries Amendment Act when the term TACC was introduced. Before 1990, the term TAC applied only to commercial fishing. In the Landings and TAC tables in this report, the TAC figures equate to the TACC unless otherwise specified.

Only actual TACCs are provided. The actual TACCs are the values as of the last day of the fishing year; e.g., 30 September.

In considering customary non-commercial, and recreational interests, the focus has been on current interests and activities rather than historical activities. In most cases, there is little information available on the nature and extent of non-commercial interests, although estimates of recreational harvest are available in some instances. Information on illegal catches and other sources of mortality is provided where available.

Yield Benchmarks

The biological reference points, Maximum Constant Yield (*MCY*) and Current Annual Yield (*CAY*) first used in the 1988 assessment continue to be used in some stock assessments. This approach is described in the section of this report titled "Guide to Biological Reference Points for Fisheries Assessment Meetings".

Sources of Data

A major source of information for these assessments is the fisheries statistics system. It is important to maintain and develop this system to provide adequate and timely data for stock assessments.

Other Information

For some assessments, draft Fisheries Assessment Reports that more fully describe the data and the analyses have been prepared in time for the Working Group or Plenary process. Once finalised, these documents are placed on the Ministry's Fisheries website in a searchable database.

Environmental Effects of Fishing

The scientific information to assess the environmental effects of fishing and enable this outcome comes primarily from research commissioned by the Ministry and, for protected species only, the Department of Conservation (DOC). The work is reviewed by the Aquatic Environment Working Group (AEWG) (or a similar DOC technical working group) or by the Biodiversity Research Advisory Group (BRAG). The Ministry has developed an "Aquatic Environment and Biodiversity Annual Review", which summarises the current state of knowledge on the environmental interactions between fisheries and the aquatic environment. The Aquatic Environment and Biodiversity Annual Review assesses the various known and potential effects of fishing on an issue-by-issue basis (e.g., the total impact of all bottom trawl and dredge fisheries on benthic habitat), whereas relatively brief fishery-specific summaries have been progressively included in this report since 2005, starting with hoki. These fishery-specific sections are reviewed by AEWG rather than by the FAWGs responsible for the stock assessment sections in each Working Group report.

Status of Stocks Summary Tables

Since 2009, the key information relevant to providing more comprehensive and meaningful information for fisheries managers, stakeholders and other interested parties has been summarised at the end of each chapter in a table format using the Guidelines for Status of the Stocks Summary Tables on pages 39–44. Beginning in 2012, selected Status of Stocks tables have incorporated a new science information quality ranking system, as specified in the Research and Science Information Standard for New Zealand Fisheries (2011). Beginning in 2013, selected Status of Stocks tables have incorporated explicit statements regarding the status of fisheries relative to overfishing thresholds.

Glossary of Common Technical Terms

Abundance Index: A quantitative measure of fish density or abundance, usually as a relative time series. An abundance index can be specific to an area or to a segment of the **stock** (e.g., mature fish), or it can refer to abundance stock-wide; the index can reflect abundance in numbers or in weight (**biomass**).

AEWG: The Aquatic Environment (Science) Working Group.

Age frequency: The proportions of fish of different ages in the **stock**, or in the **catch** taken by either the commercial fishery or research fishing. This is often estimated based on a sample. Sometimes called an age composition.

Age-length key: The proportion of fish of each age in each length-group in a sample of fish.

Age-structured stock assessment: An assessment that uses a model to estimate how the numbers at age in the stock vary over time in order to determine the past and present **status** of a fish **stock**.

a₅₀: Either the age at which 50% of fish are mature ($=A_M$) or 50% are recruited to the fishery ($=A_R$).

AIC: The Akaike Information Criterion is a measure of the relative quality of a statistical model for a given set of data. As such, AIC provides a means for model selection; the preferred model is the one with the minimum AIC value.

A_M: *Age at maturity* is the age at which fish, of a given sex, are considered to be reproductively mature. See **a₅₀**.

AMP: *Adaptive Management Programme*. This involves increased **TACC's** (for a limited period, usually 5 years) in exchange for which the industry is required to provide data that will improve understanding of **stock status**. The industry is also required to collect additional information (biological data and detailed catch and effort) and perform the analyses (e.g. **CPUE** standardisation or age structure) necessary for monitoring the **stock**.

ANTWG: Antarctic (Science) Working Group.

A_R: *Age of recruitment* is the age when fish are considered to be **recruited** to the fishery. In **stock assessments**, this is usually the youngest age group considered in the analyses. See **a₅₀**.

a_{to95}: The number of ages between the age at which 50% of a stock is mature (or recruited) and the age at which 95% of the stock is mature (or recruited).

B₀: *Virgin biomass, unfished biomass*. This is the theoretical **carrying capacity** of the **recruited** or **vulnerable** or **spawning biomass** of a fish **stock**. In some cases, it refers to the average **biomass** of the **stock** in the years before fishing started. More generally, it is the average over recent years of the biomass that theoretically would have occurred if the stock had never been fished. **B₀** is often estimated from stock modelling and various percentages of it (e.g. 40% **B₀**) are used as **biological reference points (BRPs)** to assess the relative status of a **stock**.

B_{AV}: The average historical **recruited biomass**.

Bayesian stock assessment: an approach to stock assessment that provides estimates of uncertainty (**posterior distributions**) of the quantities of interest in the assessment. The method allows the initial uncertainty (that before the data are considered) to be described in the form of **priors**. If the data are informative, they will determine the posterior distributions; if they are uninformative, the posteriors will resemble the **priors**. The initial model runs are called **MPD** (mode of the posterior distribution) runs, and provide point estimates only, with no

uncertainty. Final runs (Markov Chain Monte Carlo runs or **MCMCs**), which are often very time consuming, provide both point estimates and estimates of uncertainty.

B_{BEG} : The estimated **stock biomass** at the beginning of the fishing year.

$B_{CURRENT}$: Current **biomass** in the year of the assessment (usually a **mid-year biomass**).

Benthic - the ecological region at the lowest level of a body of water, including the sediment surface and some sub-surface layers

Biological Reference Point (BRP): A benchmark against which the **biomass** or abundance of the **stock**, or the **fishing mortality rate** (or **exploitation rate**), or **catch** itself can be measured in order to determine **stock status**. These reference points can be **targets**, **thresholds** or **limits** depending on their intended use.

Biomass: Biomass refers to the size of the **stock** in units of weight. Often, biomass refers to only one part of the **stock** (e.g., **spawning biomass**, **vulnerable biomass** or **recruited biomass**, the latter two of which are essentially equivalent).

B_{MSY} : The average **stock biomass** that results from taking an average catch of **MSY** under various types of harvest strategies. Often expressed in terms of spawning **biomass**, but may also be expressed as **recruited** or **vulnerable biomass**.

Bootstrap: A statistical methodology used to quantify the uncertainty associated with estimates obtained from a **model**. The bootstrap is often based on **Monte Carlo** re-sampling of residuals from the initial **model** fit.

BRAG – Biodiversity Research Advisory Group

B_{REF} : A reference average biomass usually treated as a management target.

Bycatch: Refers to fish species, or size classes of those species, caught in association with key target species.

B_{YEAR} : Estimated or predicted **biomass** in the named year (usually a **mid-year biomass**).

Carrying capacity: The average **stock** size expected in the absence of **fishing**. Even without fishing the **stock** size varies through time in response to stochastic environmental conditions. See **B_o** : **virgin biomass**.

Catch (C): The total weight (or sometimes number) of fish caught by fishing operations.

CAY: **Current annual yield** is the one year **catch** calculated by applying a reference **fishing mortality**, F_{REF} , to an estimate of the fishable **biomass** at the beginning of the fishing year. Also see **MAY**.

CELR: Catch-Effort Landing Return.

CLR: Catch Landing Return.

Cohort: Those individuals of a **stock** born in the same spawning season. For annual spawners, a year's **recruitment** of new individuals to a **stock** is a single cohort or **year-class**.

Collapsed: Stocks that are below the **hard limit** are deemed to be **collapsed**.

Convergence: In reference to **MCMC** results from a **Bayesian stock assessment**, convergence means that the average and the variability of the parameter estimates are not changing as the **MCMC** chain gets longer.

CPUE: Catch per unit effort is the quantity of fish caught with one standard unit of fishing effort; e.g., the number of fish taken per 1000 hooks per day or the weight of fish taken per hour of trawling. CPUE is often assumed to be a relative **abundance index**.

Customary catch: Catch taken by tangata whenua to meet their customary needs.

CV: Coefficient of variation. A statistic commonly used to represent variability or uncertainty. For example, if a biomass estimate has a CV of 0.2 (or 20%), this means that the error in this estimate (the difference between the estimate and the true biomass) will typically be about 20% of the estimate.

Density-dependence: Fish populations are thought to self-regulate: as population biomass increases, growth may slow down, mortality may increase, recruitment may decrease or maturity may occur later. Growth is density-dependent if it slows down as biomass increases.

Depleted: Stocks that are below the **soft limit** are deemed to be **depleted**. Stocks can become **depleted** through **overfishing**, or environmental factors, or a combination of the two.

Discards – the portion of the catch thrown away at sea

DWWG: The Deepwater (Science) Working Group.

ECER: Eel Catch-Effort Return.

ECLR: Eel Catch Landing Return.

Ecosystem –a biological community of interacting organisms and their physical environment.

EEZ: An **Exclusive Economic Zone** is a maritime zone beyond the **Territorial Sea** over which the coastal state has sovereign rights over the exploration and use of marine resources. Usually, a state's EEZ extends to a distance of 200 nautical miles (370 km) out from its coast, except where resulting points would be closer to another country.

Equilibrium: A theoretical model state that arises when the **fishing mortality**, **exploitation pattern** and other fishery or **stock** characteristics (growth, natural mortality, **recruitment**) do not change from year to year.

Exploitable biomass: Refers to that portion of a **stock's biomass** that is available to the fishery. Also called **recruited biomass** or **vulnerable biomass**.

Exploitation pattern: The relative proportion of each age or size class of a **stock** that is vulnerable to fishing. See **selectivity ogive**.

Exploitation rate: The proportion of the **recruited** or **vulnerable biomass** that is caught during a certain period, usually a fishing year.

F: The **fishing intensity** or **fishing mortality rate** is that part of the total mortality rate applying to a fish **stock** that is caused by fishing. Usually expressed as an instantaneous rate.

$F_{0.1}$: The **fishing mortality rate** at which the increase in **equilibrium yield per recruit** in weight per unit of effort is 10% of the **yield per recruit** produced by the first unit of effort on the unexploited **stock** (i.e., the slope of the **yield per recruit** curve for the $F_{0.1}$ rate is only 1/10th of the slope of the **yield per recruit** curve at its origin).

$F_{40\%B_0}$: The **fishing mortality rate** associated with a biomass of 40% B_0 at **equilibrium** or on average.

$F_{40\%SPR}$: The **fishing mortality rate** associated with a spawning biomass per recruit (**SPR**) (or equivalently a spawning potential ratio) of 40% B_0 at equilibrium or on average.


FAWGs: Fisheries Assessment (Science) Working Groups.

Fishing intensity: A general term that encompasses the related concepts of **fishing mortality** and **exploitation rate**.

Fishing mortality: That part of the total mortality rate applying to a fish **stock** that is caused by fishing. Usually expressed as an instantaneous rate.

Fishing year: For most fish stocks, the fishing year runs from 1 October in one year to 30 September in the next. The second year is often used as shorthand for the split years. For example, 2015 is shorthand for 2014–15.

FMA: Fishery Management Area. The New Zealand **EEZ** is divided into 10 fisheries management units:



F_{MAX} : The **fishing mortality rate** that maximises **equilibrium yield per recruit**. F_{MAX} is the **fishing mortality** level that defines **growth overfishing**. In general, F_{MAX} is different from F_{MSY} (the **fishing mortality** that maximises **sustainable yield**), and is always greater than or equal to F_{MSY} , depending on the **stock-recruitment relationship**.

F_{MEY} : The fishing mortality corresponding to the maximum (**sustainable**) economic yield.

F_{MSY} : The **fishing mortality rate** that, if applied constantly, would result in an average catch corresponding to the **Maximum Sustainable Yield (MSY)** and an average biomass corresponding to B_{MSY} . Usually expressed as an instantaneous rate.

F_{REF} : The **fishing mortality** that is associated with an average biomass of B_{REF} .

FRML – Fisheries Related Mortality Limit.

Growth overfishing: Growth overfishing occurs when the **fishing mortality rate** is above F_{MAX} . This means that on average fish are caught before they have a chance to reach their maximum growth potential.

Hard Limit: A biomass limit below which fisheries should be considered for closure.

Harvest Strategy: For the purpose of the Harvest Strategy Standard, a harvest strategy simply specifies **target** and **limit reference points** and management actions associated with achieving the **targets** and avoiding the **limits**.

HMS: Highly Migratory Species.

HMSWG: Highly Migratory Species (Science) Working Group.

Hyperdepletion: The situation where an abundance index, such as **CPUE**, decreases faster than the true abundance.

Hyperstability: The situation where an abundance index, such as **CPUE**, decreases more slowly than the true abundance.

Incidental capture: Refers to non-fish and protected species which were not targeted, but were caught.

Index: Same as an **abundance index**.

LCER: Longline Catch-Effort Return.

Length frequency: The distribution of numbers at length from a sample of the **catch** taken by either the commercial fishery or research fishing. This is sometimes called a length composition.

Length-Structured Stock Assessment: An assessment that uses a model to estimate how the numbers at length in the stock vary over time in order to determine the past and present **status** of a fish **stock**.

Limit: a **biomass** or fishing mortality **reference point** that should be avoided with high probability. The Harvest Strategy Standard defines both **soft limits** and **hard limits**.

M: The (instantaneous) **natural mortality rate** is that part of the total mortality rate applying to a fish **stock** that is caused by predation and other natural events.

MAFWG: Marine Amateur Fisheries (Science) Working Group.

MALFIRM: Maximum Allowable Limit of Fishing Related Mortality.

Maturity: Refers to the ability of fish to reproduce.

Maturity ogive: A curve describing the proportion of fish of different ages or sizes that are mature.

MAY: Maximum average yield is the average **maximum sustainable yield** that can be produced over the long term under a constant fishing mortality strategy, with little risk of **stock** collapse. A constant fishing mortality strategy means catching a constant percentage of the biomass present at the beginning of each fishing year. **MAY** is the long-term average annual catch when the catch each year is the **CAY**. Also see **CAY**.

MCMC: Markov Chain Monte Carlo. See **Bayesian stock assessment**.

MCY: Maximum constant yield is the maximum sustainable yield that can be produced over the long term by taking the same catch year after year, with little risk of stock collapse.

MIDWG: Middle-depths (Science) Working Group.

Mid-year biomass: The biomass after half the year's catch has been taken.

MLS: Minimum Legal Size. Fish above the **MLS** can be retained while those below it must be returned to the sea.

Model: A set of equations that represents the population dynamics of a fish stock.

Monte Carlo Simulation: is an approach whereby the inputs that are used for a calculation are re-sampled many times assuming that the inputs follow known statistical distributions. The Monte Carlo method is used in many applications such as **Bayesian stock assessments**, parametric bootstraps and stochastic **projections**.

MPD: *Mode of the (joint) posterior distribution*. See **Bayesian stock assessment**.

MSY: Maximum sustainable yield is the largest long-term average catch or yield that can be taken from a **stock** under prevailing ecological and environmental conditions, and the current selectivity patterns exhibited by the fishery.

MSY-compatible reference points: *MSY-compatible reference points* include B_{MSY} , F_{MSY} and **MSY** itself, as well as analytical and conceptual **proxies** for each of these three quantities.

Natural mortality (rate): That part of the total mortality rate applying to a fish **stock** that is caused by predation and other natural events. Usually expressed as an instantaneous rate.

NCELR: Set Net Catch-Effort Landing Return.

NINS: Northern Inshore (Science) Working Group.

Objective function: An equation to be optimised (minimised or maximised) given certain constraints using non-linear programming techniques.

Otolith: One of the small bones or particles of calcareous substance in the internal ear of teleosts (bony fishes) that are used to determine their age.

Overexploitation: A situation where observed **exploitation** (or **fishing mortality**) rates are higher than **target levels**.

Overfishing: A situation where observed **fishing mortality** (or **exploitation**) rates are higher than **target** or **threshold** levels.

Partition: The way in which a fish stock or population is characterised, or split, in a stock assessment model; for example, by sex, age and maturity.

PCELR: Paua Catch Effort and Landing Return.

Population: A group of fish of one species that shares common ecological and genetic features. The **stocks** defined for the purposes of **stock assessment** and management do not necessarily coincide with self-contained populations.

Population dynamics: In general, refers to the biological and fishing processes that result in changes in fish **stock** abundance over time.

Posterior: a mathematical description of the uncertainty in some quantity (e.g., **biomass**) estimated in a **Bayesian stock assessment**. This is generally depicted as a frequency distribution (often plotted along with the **prior** distribution to show how much the two diverge).

Potential Biological Removal (PBR) - an estimate of the number of seabirds that may be killed without causing the population to decline below half the carrying capacity.

Pre-recruit: An individual that has not yet entered the fished component of the **stock** (because it is either too young or too small to be vulnerable to the fishery).

Prior: available information (often in the form of expert opinion) regarding the potential range of values of a parameter in a **Bayesian stock assessment**. Uninformative priors are used where there is no such information.

Production Model: A **stock model** that describes how the **stock biomass** changes from year to year (or, how **biomass** changes in **equilibrium** as a function of **fishing mortality**), but which does not keep track of the age or length frequency of the stock. The simplest production functions aggregate all of the biological characteristics of growth, **natural mortality** and reproduction into a simple, deterministic **model** using three or four parameters. Production models are primarily used in simple data situations, where total catch and effort data are available but age-structured information is either unavailable or deemed to be less reliable (although some versions of production models allow the use of age-structured data).

Productivity: Productivity is a function of the biology of a species and the environment in which it lives. It depends on growth rates, **natural mortality**, **age at maturity**, maximum average age and other relevant life history characteristics. Species with high **productivity** are able to sustain higher rates of **fishing mortality** than species with lower **productivity**. Generally, species with high productivity are more resilient and take less time to rebuild from a **depleted** state.

Projection: Predictions about trends in stock size and fishery dynamics in the future. Projections are made to address “what-if” questions of relevance to management. Short-term (1–5 years) projections are typically used in support of decision-making. Longer term projections become much more uncertain in terms of absolute quantities, because the results are strongly dependent on **recruitment**, which is very difficult to predict. For this reason, long-term projections are more useful for evaluating overall management strategies than for making short-term decisions.

Proxy: A surrogate for B_{MSY} , F_{MSY} or MSY that has been demonstrated to approximate one of these three metrics through theoretical or empirical studies.

q: Catchability is the proportion of fish that are caught by a defined unit of fishing effort. The constant relating an **abundance index** to the true biomass (the **abundance index** is approximately equal to the true biomass multiplied by the catchability).

Quota Management Areas (QMA): QMAs are geographic areas within which fish stocks are managed in the TS and EEZ.

Quota Management System (QMS): The QMS is the name given to the system by which the total commercial catch from all the main fish **stocks** found within New Zealand’s 200 nautical mile EEZ is regulated.

Recruit: An individual that has entered the fished component of the **stock**. Fish that are not recruited are either not catchable by the gear used (e.g., because they are too small) or live in areas that are not fished.

Recruited biomass: Refers to that portion of a **stock’s biomass** that is available to the fishery; also called **exploitable biomass** or **vulnerable biomass**.

Recruitment: The addition of new individuals to the fished component of a **stock**. This is determined by the size and age at which fish are first caught.

Reference Point: A benchmark against which the biomass or abundance of the **stock** or the **fishing mortality rate** (or **exploitation rate**) can be measured in order to determine its **status**. These reference points can be targets, thresholds or limits depending on their intended use.

RLWG: Rock Lobster (Science) Working Group.

SAMWG: Stock Assessment Methods (Science) Working Group.

S_{AV} : The average historical **spawning biomass**.

Selectivity ogive: Curve describing the relative vulnerability of fish of different ages or sizes to the fishing gear used.

SFWG: The Shellfish (Science) Working Group.

SINS: Southern Inshore (Science) Working Group.


Soft Limit: A **biomass** limit below which the requirement for a formal, time-constrained **rebuilding plan** is triggered.

Spawning biomass: The total weight of sexually mature fish in the **stock**. This quantity depends on the abundance of **year classes**, the **exploitation** pattern, the rate of growth, both fishing and **natural mortality rates**, the onset of sexual maturity, and environmental conditions. Same as **mature biomass**.

Spawning (biomass) Per Recruit or Spawning Potential Ratio (SPR): The expected lifetime contribution to the **spawning biomass** for the average recruit to the fishery. For a given exploitation pattern, rate of growth, maturity schedule and **natural mortality**, an **equilibrium** value of SPR can be calculated for any level of fishing mortality. SPR decreases monotonically with increasing fishing mortality.

Statistical area: See the map below for the official **Territorial Sea** and Exclusive Economic Zone (**EEZ**) statistical areas.

Crown Copyright Reserved



Steepness: A parameter of **stock-recruitment relationships** that determines how rapidly, or steeply, it rises from the origin, and therefore how resilient a stock is to rebounding from a depleted state. It equates to the proportion of virgin recruitment that corresponds to 20% B_0 . A steepness value greater than about 0.9 is considered to be high, while one less than about 0.6 is considered to be low. The minimum value is 0.2.

Stock: The term has different meanings. Under the Fisheries Act, it is defined with reference to units for the purpose of fisheries management (Fishstock). On the other hand, a biological stock is

a population of a given species that forms a reproductive unit and spawns little if at all with other units. However, there are many uncertainties in defining spatial and temporal geographical boundaries for such biological units that are compatible with established data collection systems. For this reason, the term “stock” is often synonymous with an assessment / management unit, even if there is migration or mixing of some components of the assessment/management unit between areas.

Stock assessment: The analysis of available data to determine stock status, usually through application of statistical and mathematical tools to relevant data in order to obtain a quantitative understanding of the **status** of the **stock** relative to defined management benchmarks or **reference points** (e.g. B_{MSY} and/or F_{MSY}).

Stock-recruitment relationship: An equation describing how the expected number of recruits to a stock varies as the **spawning biomass** changes. The most frequently used stock-recruitment relationship is the asymptotic Beverton-Holt equation, in which the expected number of recruits changes very slowly at high levels of spawning biomass.

Stock status: Refers to a determination made, on the basis of **stock assessment** results, about the current condition of the **stock**. Stock status is often expressed relative to management benchmarks and **biological reference points** such as B_{MSY} or B_0 or F_{MSY} or $F_{\%SPR}$. For example, the current biomass may be said to be above or below B_{MSY} or to be at some percentage of B_0 . Similarly, fishing mortality may be above or below F_{MSY} or $F_{\%SPR}$.

Stock structure: (1) Refers to the geographical boundaries of the **stocks** assumed for assessment and management purposes (e.g., albacore tuna may be assumed to be comprised of two separate **stocks** in the North Pacific and South Pacific), (2) Refers to boundaries that define self-contained **stocks** in a genetic sense, (3) refers to known, inferred or assumed patterns of residence and migration for stocks that mix with one another.

Surplus production: The amount of **biomass** produced by the **stock** (through growth and **recruitment**) over and above that which is required to maintain the [total stock] **biomass** at its current level. If the catch in each year is equal to the surplus production then the biomass will not change.

Sustainability: Pertains to the ability of a fish **stock** to persist in the long-term. Because fish **populations** exhibit natural variability, it is not possible to keep all fishery and **stock** attributes at a constant level simultaneously, thus sustainable fishing does not imply that the fishery and **stock** will persist in a constant **equilibrium** state. Because of natural variability, even if F_{MSY} could be achieved exactly each year, **catches** and **stock biomass** will oscillate around their average MSY and B_{MSY} levels, respectively. In a more general sense, sustainability refers to providing for the needs of the present generation while not compromising the ability of future generations to meet theirs.

TAC: Total Allowable Catch is the sum of the Total Allowable Commercial Catch (**TACC**) and the allowances for customary Maori interests, recreational fishery interests and other sources of fishing-related mortality that can be taken in a given period, usually a year.

TACC: Total Allowable Commercial Catch is the total regulated commercial catch from a **stock** in a given time period, usually a fishing year.

Target: Generally, a **biomass**, **fishing mortality** or **exploitation rate** level that management actions are designed to achieve with at least a 50% probability.

Threshold: Generally, a **biological reference point** that raises a “red flag” indicating that **biomass** has fallen below the **target**, or **fishing mortality** or **exploitation rate** has increased above its **target**, to the extent that additional management action may be required in order to prevent the stock from declining further and possibly breaching the **soft limit**.

TCEPR: Trawl Catch-Effort Processing Return.

TCER: Trawl Catch-Effort Return.

TLCER: Tuna Longline Catch-Effort Return.

TS: Territorial Sea: a belt of coastal waters extending at most 12 nautical miles (22.2 km; 13.8 mi) from the baseline (usually the mean low-water mark) of a coastal state.

U_{MSY} : The **exploitation rate** associated with the maximum sustainable yield.

$U_{40\%B_0}$: The **exploitation rate** associated with a biomass of 40% B_0 at equilibrium or on average.

von Bertalanffy equation: An equation describing how fish increase in length as they grow older. The mean length (L) at age a is

$$L = L_{\infty}(1 - e^{-k(a-t_0)})$$

where L_{∞} is the average length of the oldest fish, k is the average growth rate (Brody coefficient) and t_0 is a constant.

Vulnerable biomass: Refers to that portion of a **stock's biomass** that is available to the fishery. Also called **exploitable biomass** or **recruited biomass**.

Year class (cohort): Fish in a **stock** that were born in the same year. Occasionally, a **stock** produces a very small or very large year class which can be pivotal in determining **stock** abundance in later years.

Yield: Catch expressed in terms of weight.

Yield per Recruit (YPR): The expected lifetime **yield** for the average recruit. For a given **exploitation pattern**, rate of growth, and **natural mortality**, an **equilibrium** value of YPR can be calculated for each level of **fishing mortality**. YPR analyses may play an important role in advice for management, particularly as they relate to minimum size controls.

Z: Total mortality rate. The sum of **natural** and **fishing mortality rates**.

Terms of Reference for Fisheries Assessment Working Groups (FAWGs) in 2018

Overall purpose

The purpose of the FAWGs is to assess the status of fish stocks managed within the Quota Management System, as well as other important species of interest to New Zealand. Based on scientific information the FAWGs assess the current status of fish stocks or species relative to MSY-compatible reference points and other relevant indicators of stock status, conduct projections of stock size and status under alternative management scenarios, and review results from relevant research projects. They do not make management recommendations or decisions (this responsibility lies with Fisheries New Zealand (FNZ) fisheries managers and the Minister responsible for fisheries).

Preparatory tasks

1. Prior to the beginning of the main sessions of FAWG meetings (January to May and September to November), FNZ fisheries scientists will produce a list of stocks and issues for which new stock assessments or evaluations are likely to become available prior to the next scheduled sustainability rounds. This list will include stocks for which the fishing industry and others intend to directly purchase scientific analyses. It is therefore incumbent on those purchasing research to inform the relevant FAWG chair of their intentions at least three months prior to the start of the sustainability round. FAWG Chairs will determine the final timetables and agendas for each Working Group.
2. At least six months prior to the main sessions of FAWG meetings, FNZ fisheries managers will alert FNZ science managers and the Principal Advisor Fisheries Science to unscheduled special cases for which assessments or evaluations are urgently needed.

Technical objectives

3. To review new research information on stock structure, productivity, abundance and related topics for each fish stock/issue under the purview of individual FAWGs.
4. Where possible, to derive appropriate MSY-compatible reference points¹ for use as reference points for determining stock status, based on the Harvest Strategy Standard for New Zealand Fisheries² (the Harvest Strategy Standard).
5. To conduct stock assessments or evaluations for selected fish stocks in order to determine the status of the stocks relative to MSY-compatible reference points¹ and associated limits, based on the "Guide to Biological Reference Points for Fisheries Assessment Meetings", the Harvest Strategy Standard, and relevant management reference points and performance measures set by fisheries managers.
6. For stocks where the status is unknown, FAWGs should use existing data and analyses to draw logical conclusions about likely future trends in biomass levels and/or fishing mortality (or exploitation) rates if current catches and/or TACs/TACCs are maintained, or if fishers or fisheries managers are considering modifying them in other ways.

¹ MSY-compatible reference points include those related to stock biomass (i.e. B_{MSY}), fishing mortality (i.e. F_{MSY}) and catch (i.e. MSY itself), as well as analytical and conceptual proxies for each of the three of these quantities.

² Link to the Harvest Strategy Standard: <http://fs.fish.govt.nz/Page.aspx?pk=104>

7. Where appropriate and practical, to conduct projections of likely future stock status using alternative fishing mortality (or exploitation) rates or catches and other relevant management actions, based on the Harvest Strategy Standard and input from the FAWG and fisheries managers.
8. For stocks that are deemed to be depleted or collapsed, to develop alternative rebuilding scenarios based on the Harvest Strategy Standard and input from the FAWG and fisheries managers.
9. For fish stocks for which new stock assessments or analyses are not conducted in the current year, to review the existing Fisheries Assessment Plenary report text on the “Status of the Stocks” in order to determine whether the latest reported stock status summary is still relevant; else to revise the evaluations of stock status based on new data or analyses, or other relevant information.

Working Group reports

10. To include in the Working Group report information on commercial, Māori customary, non-commercial and recreational interests in the stock; as well as all other mortality to that stock caused by fishing, which might need to be allowed for in setting a TAC or TACC. Estimates of recreational harvest will normally be provided by the Marine Amateur Fisheries Working Group (MAFWG).
11. To provide information and advice on other management considerations (e.g. area boundaries, by-catch issues, effects of fishing on habitat, other sources of mortality, and input controls such as mesh sizes and minimum legal sizes) required for specifying sustainability measures. Sections of the Working Group reports related to bycatch and other environmental effects of fishing will be reviewed by the Aquatic Environment Working Group (AEWG) although the relevant FAWG is encouraged to identify to the AEWG Chair any major discrepancies between these sections and their understanding of the operation of relevant fisheries.
12. To summarise the stock assessment methods and results, along with estimates of MSY-compatible reference points and other metrics that may be used as benchmarks for assessing stock status.
13. To review, and update if necessary, the “Status of the Stocks” tables in the Fisheries Assessment Plenary report for all stocks under the purview of individual FAWGs (including those for which a full assessment has not been conducted in the current year) based on new data or analyses, or other relevant information.
14. For all important stocks, to complete (and/or update) the Status of Stocks tables using the template provided in the Introductory chapter of the most recent May and November Plenary reports.
15. It is desirable that full agreement amongst technical experts is achieved on the text of the FAWG reports, particularly the “Status of the Stocks” sections, noting that the AEWG will review sections on bycatch and other environmental effects of fishing, and the MAFWG will provide text on recreational harvests. If full agreement amongst technical experts cannot be reached, the Chair will determine how this will be depicted in the FAWG report, will document the extent to which agreement or consensus was achieved, and record and attribute any residual disagreement in the meeting notes.

Working Group input to the Plenary

16. To advise the Principal Advisor Fisheries Science about stocks requiring review by the Fisheries Assessment Plenary and those stocks that are not believed to warrant review by the Plenary. The general criteria for determining which stocks should be discussed by the Plenary are that (i) the assessment is controversial and Working Group members have had difficulty reaching consensus on one or more base cases, or (ii) the assessment is the first for a particular stock or the methodology has been substantially altered since the last assessment, or (iii) new data or analyses have become available that alter the previous assessment, particularly assessments of recent or current stock status, or projections of likely future stock status. Such information could include:
- new or revised estimates of MSY-compatible reference points, recent or current biomass, productivity or yield projections;
 - the development of a major trend in the catch or catch per unit effort; or
 - any new studies or data that extend understanding of stock structure, fishing patterns, or non-commercial activities, and result in a substantial effect on assessments of stock status.

Membership and Protocols for all Science Working Groups

17. FAWG members are bound by the Membership and Protocols required for all Science Working Group members.

Terms of Reference for the Aquatic Environment Working Group (AEWG) in 2018

Overall purpose

For all New Zealand fisheries in the New Zealand TS and EEZ as well as other important fisheries in which New Zealand engages:

to assess, based on scientific information, the effects of (and risks posed by) fishing, aquaculture, and enhancement on the aquatic environment, including:

- bycatch and unobserved mortality of protected species (e.g. seabirds and marine mammals), fish, and other marine life, and consequent impacts on populations;
- effects on benthic ecosystems, species, and habitat;
- effects on biodiversity, including genetic diversity;
- changes to ecosystem structure and function from fishing, including trophic effects; and
- effects of aquaculture and fishery enhancement on the environment and on fishing.

Where appropriate and feasible, such assessments should explore the implications of the effect, including with respect to government standards, other agreed reference points, or other relevant indicators of population or environmental status. Where possible, projections of future status under alternative management scenarios should be made.

AEWG does not make management recommendations or decisions (this responsibility lies with Fisheries New Zealand (FNZ) fisheries managers and the Minister responsible for Fisheries).

Fisheries New Zealand also convenes a Biodiversity Research Advisory Group (BRAG) which has a similar review function to the AEWG. Projects reviewed by BRAG and AEWG have some commonalities in that they relate to aspects of the marine environment. However, the key focus of projects considered by BRAG is on the functionality of the marine ecosystem and its productivity, whereas projects considered by AEWG more commonly focus on the direct effects of fishing, aquaculture or enhancement.

Preparatory tasks

1. Prior to the beginning of AEWG meetings each year, FNZ fisheries scientists will produce a list of issues for which new assessments or evaluations are likely to become available that year.
2. The Ministry's research planning processes should identify most information needs well in advance but, if urgent issues arise, FNZ-Fisheries or aquaculture staff will alert the relevant AEWG chair prior to the required meeting of items that could be added to the agenda. AEWG Chairs will determine the final timetables and agendas for meetings.

Technical objectives

3. To review any new research information on fisheries, aquaculture or enhancement impacts, including risks of impacts, and the relative or absolute sensitivity or susceptibility of potentially affected species, populations, habitats, and systems.
4. To estimate appropriate reference points for determining population, system, or environmental status, noting any draft or published Standards.

5. To conduct environmental assessments or evaluations for selected species, populations, habitats, or systems in order to determine their status relative to appropriate reference points and Standards, where such exist.
6. In addition to determining the status of the species, populations, habitats, and systems relative to reference points, and particularly where the status is unknown, AEWG should explore the potential for using existing data and analyses to draw conclusions about likely future trends in fishing effects or status if current fishing methods, effort, catches, and catch limits are maintained, or if fishers or fisheries managers are considering modifying them in other ways.
7. Where appropriate and practical, to conduct or request projections of likely future status using alternative management actions, based on input from AEWG, fisheries plan advisers and fisheries and standards managers, noting any draft or published Standards.
8. For species or populations deemed to be depleted or endangered, to develop ideas for alternative rebuilding scenarios to levels that are likely to ensure long-term viability based on input from AEWG, fisheries managers, noting any draft or published Standards.
9. To review and revise existing environmental and ecosystem consideration sections of Fisheries Assessment Plenary report text based on new data or analyses, or other relevant information.

Working Group input to annual Aquatic Environment and Biodiversity Review

10. To include in contributions to the Aquatic Environment and Biodiversity Review (AEBAR) summaries of information on selected issues that may relate to species, populations, habitats, or systems that may be affected by fishing, aquaculture or enhancement. These contributions are analogous to Working Group reports from the Fisheries Assessment Working Groups.
11. To provide information and scientific advice on management considerations (e.g. area boundaries, by-catch issues, effects of fishing on habitat, other sources of mortality, and input controls such as mesh sizes and minimum legal sizes) that may be relevant for setting sustainability measures.
12. To summarise the assessment methods and results, along with estimates of relevant standards, references points, or other metrics that may be used as benchmarks or to identify risks to the aquatic environment.
13. It is desirable that full agreement among technical experts is achieved on the text of contributions to the AEBAR. If full agreement among technical experts cannot be reached, the Chair will determine how this will be depicted in the AEBAR, will document the extent to which agreement or consensus was achieved, and record and attribute any residual disagreement in the meeting notes.
14. To advise the Principal Advisor Fisheries Science and Aquatic Environment manager about issues of particular importance that may require independent review or updating in the AEBAR. The general criterion for determining which issues should be discussed by a wider group or text changed in the AEBAR is that new data or analyses have become available that alter the previous assessment of an issue, particularly assessments of population status or projection results. Such information could include:
 - New or revised estimates of environmental reference points, recent or current population status, trend, or projections;
 - The development of a major trend in bycatch rates or amount;
 - Any new studies or data that extend understanding of population, system, or environmental susceptibility to an effect or its recoverability, fishing patterns, or mitigation measures that have

a substantial implications for a population, system, or environment or identify risks associated with fishing activity, aquaculture or enhancement; and

- Consistent performance outside accepted reference points or Standards.

Membership and Protocols for all Science Working Groups

15. The AEWG is bound by the same membership and protocols as other Science Working Groups (see separate document).

Terms of Reference for the Antarctic Working Group (ANTWG) in 2018

Overall purpose

The purpose of the ANTWG is to review science and research information intended for submission to or use by the Commission for the Conservation of Antarctic Marine Living Resources (CCAMLR). CCAMLR is an inter-governmental organisation that is committed to conserving the marine life of the Southern Ocean while allowing rational use of marine resources, including commercial fishing. The CCAMLR Convention requires that management considers the effects of fishing on dependent and associated species as well as on the target species. The area of jurisdiction of the CCAMLR Convention is approximately south of the circumpolar Antarctic Polar Front in the Southern Ocean. Science and research requested or used by CCAMLR may include, inter alia, fishery characterisations, abundance indices, catch-at-age or catch-at-length data, and stock assessment modelling to assess the status of fish stocks managed by CCAMLR; bycatch and unobserved mortality of protected species, fish, and other marine life; effects on biodiversity and benthic biodiversity, species, and habitat; and changes to ecosystem structure and function as a result of fishing, including trophic effects. The ANTWG also undertakes scientific review of documents and papers that may be submitted to the scientific working groups of CCAMLR to aid and inform its management. The ANTWG does not make management recommendations or decisions; these responsibilities lie with CCAMLR's Scientific Committee and the Commission.

Preparatory tasks

1. Prior to the first meeting of the ANTWG each year, the ANTWG Chair will produce a list of stocks/issues for which new stock assessments, evaluations, impact assessments, risk assessments, or other scientific analyses have been requested by the CCAMLR Scientific Committee or the Commission (including its contributing bodies), fishing industry, or other stakeholders. The ANTWG Chair will determine the final timetables and agendas of the working group each year, taking account of the available time and resources.

Technical objectives

2. To review new research information on stock structure, productivity, abundance and related topics for each fish stock or environmental issue under the purview of the ANTWG.
3. Where possible, to derive yields or reference points requested by CCAMLR's Scientific Committee or Commission related to fish stocks or environmental issues relevant to CCAMLR fisheries.
4. To conduct stock assessments or evaluations for selected stocks in order to determine the precautionary yields and status of the stocks relative to the requested reference points or, if no such reference points are specified by CCAMLR, MSY-compatible reference points and associated limits, based on the "Guide to Biological Reference Points for Fisheries Assessment Meetings" and New Zealand's Harvest Strategy Standard.
5. For stocks where the status is unknown, the ANTWG should, where possible, use any existing data and analyses to draw conclusions about likely future trends in biomass levels and/or fishing mortality (or exploitation) rates if current catches and/or TACs are maintained, or if fishers or CCAMLR are considering modifying them in other ways.
6. Where requested by the CCAMLR Scientific Committee or Commission, to conduct projections of likely future stock status using alternative fishing mortality (or exploitation) rates or catches

- and other relevant management actions, based on input from the ANTWG and any guidance from the CCAMLR Scientific Committee or Commission.
7. Where requested by the CCAMLR Scientific Committee or Commission, in relation to specified stocks, to develop and report on alternative rebuilding scenarios.
 8. To conduct environmental impact assessments and qualitative or quantitative risk assessments in relation to bycatch species, other species of concern, benthic systems, or vulnerable marine ecosystems to support the work of the CCAMLR Scientific Committee and Commission.

Working Group reports

9. To review, and update if necessary, the “Status of the Stocks” tables in the Fisheries Assessment Plenary report based on new data or analyses, or other relevant information.
10. To complete (and/or update) the Status of Stocks tables using the template provided in the Introductory chapter of the most recent May Plenary report.
11. It is desirable that full agreement amongst technical experts is achieved on the text of the ANTWG reports. If full agreement amongst technical experts cannot be reached, the Chair will determine how this will be depicted in the ANTWG report, will document the extent to which agreement or consensus was achieved, and record and attribute any residual disagreement in the meeting notes.

Papers and reports to CCAMLR

12. Papers and reports summarising work reviewed by the ANTWG are generally submitted to CCAMLR’s Scientific Committee, and their content varies widely. It is desirable that full agreement amongst technical experts is achieved on the content of such papers or reports, noting that deadlines for submission to CCAMLR may require the Chair to finalise text after a meeting of the ANTWG has considered and resolved scientific issues. If full agreement amongst technical experts cannot be reached, the Chair will determine how this will be depicted in the paper or report to be submitted to CCAMLR. In such cases, the Chair will also document the extent to which agreement or consensus was achieved and record and attribute any residual disagreement in the meeting notes.

Membership and Protocols for all Science Working Groups

13. ANTWG members are bound by the Membership and Protocols required for all Science Working Group members.

Terms of Reference for the Marine Amateur Fisheries Working Group (MAFWG) in 2018

Overall purpose

The purpose of the MAFWG is to assess the harvest of marine amateur fishers from fish stocks managed within or outside the Quota Management System and to review other scientific or research information relevant to the management of marine amateur fisheries. MAFWG does not make management recommendations or decisions; this responsibility lies with Fisheries New Zealand (FNZ) fisheries managers and the Minister responsible for fisheries.

Preparatory tasks

1. It is anticipated that marine amateur fisheries research will focus primarily on the estimation of amateur harvests of fish stocks based on corroborated off-site national surveys conducted about every 5 years. At least six months before any such survey is conducted, FNZ fisheries managers will alert FNZ science managers and the Principal Advisor Fisheries Science to their priority stocks for harvest estimation to facilitate good survey design. In years when national surveys are not being conducted, FNZ fisheries managers and fisheries scientists will work closely together to prioritise the meeting of other key information needs in relation to marine amateur fisheries.

Technical objectives

2. To review new research information on the harvest and harvesting patterns of marine amateur fishers using off-site and/or on-site methods, focussing primarily on priority non-commercial and shared stocks or fisheries identified by fisheries managers.
3. To develop methods for making reliable estimates of total catch by fish stock (finfish and shellfish); catch per unit of effort (CPUE); fish lengths and weights within the harvest; daily bag sizes in relation to limits; the spatial and temporal variability of fishing, CPUE, or harvest; and other information likely to inform fisheries management decisions, the development of environmental standards, or the formulation of relevant policy.

Working Group reports

4. In collaboration with relevant Stock Assessment Working Group Chairs, to provide timely and current information on marine amateur harvest for Working Group reports for non-commercial and shared stocks. MAFWG will also periodically review information on marine amateur harvest in Working Group reports to ensure accuracy and currency.
5. As necessary, provide information and advice on other management considerations for marine amateur fisheries (e.g. effects of fishing on habitat, other sources of mortality, and potential input controls such as bag limits, mesh sizes, and minimum legal sizes) required for specifying sustainability measures.
6. It is desirable that full agreement amongst technical experts is achieved on the information provided for Working Group reports on the harvest and other aspects of marine amateur fisheries. If full agreement amongst technical experts cannot be reached, the Chair will determine how this will be depicted in the Working Group report, will document the extent to which agreement or consensus was achieved, and record and attribute any residual disagreement in the meeting notes.

Membership and Protocols for all Science Working Groups

7. MAFWG members are bound by the Membership and Protocols required for all Science Working Group members.

Membership and Protocols for all Science Working Groups in 2018

This document summarises the protocols for membership and participation in all Science Working Groups including Fisheries Assessment Working Groups (FAWGs), the Aquatic Environment Working Group (AEWG), the Biodiversity Research Advisory Group (BRAG), the Highly Migratory Species Working Group (HMS), the South Pacific Working Group (SPACWG), the Antarctic Working Group (ANTWG), and the Marine Amateur Fisheries Working Group (MAFWG).

Working Group chairs

1. The Ministry will select and appoint the Chairs for Science Working Groups. The Chair will be a Fisheries New Zealand (FNZ) fisheries or marine scientist who is an active participant in the Working Group, providing technical input, rather than simply being a facilitator. Working Group Chairs will be responsible for:
 - ensuring that Working Group participants are aware of the Terms of Reference for the Working Group, and that the Terms of Reference are adhered to by all participants;
 - setting the rules of engagement, facilitating constructive questioning, and focussing on relevant issues;
 - ensuring that all peer review processes are conducted in accordance with the Research and Science Information Standard for New Zealand Fisheries³ (the Research Standard), and that research and science information is reviewed by the relevant Working Group against the P R I O R principles for science information quality (page 6) and the criteria for peer review (pages 12–16) in the Standard;
 - requesting and documenting the affiliations of participants at each Working Group meeting that have the potential to be, or to be perceived to be, a conflict of interest of relevance to the research under review (refer to page 15 of the Research Standard). Chairs are responsible for managing conflicts of interest, and ensuring that fisheries management implications do not jeopardise the objectivity of the review or result in biased interpretation of results;
 - ensuring that the quality of information that is intended or likely to inform fisheries management decisions, the development of environmental standards or the formulation of relevant fisheries policy is ranked in accordance with the information ranking guidelines in the Research Standard (page 21–23), and that resulting information quality ranks are appropriately documented in the Plenary and the Aquatic Environment and Biodiversity Annual Review (AEBAR);
 - striving for consensus while ensuring the transparency and integrity of research analyses, results, conclusions and final reports; and
 - reporting on Working Group recommendations, conclusions and action items; and ensuring follow-up and communication with the FNZ Principal Advisor Fisheries Science, relevant FNZ fisheries management staff, and other key stakeholders.

Working Group members

2. Membership of Science Working groups will be open to any participant with the agreement of the Working Group Chair.
3. Working Groups will consist of the following participants:

³ Link to the Research Standard: <http://www.fish.govt.nz/en-nz/Publications/Research+and+Science+Information+Standard.htm>

- Fisheries New Zealand fisheries science chair – required;
 - research providers – required (may be the primary researcher, or a designated substitute capable of presenting and discussing the agenda item);
 - other scientists not conducting the presented research to act in a peer review capacity;
 - representatives of relevant FNZ fisheries management teams; and
 - any interested party who agrees to the standards of participation below.
4. Working Group participants must commit to:
- participating appropriately in the discussion;
 - resolving issues;
 - following up on agreements and tasks;
 - maintaining confidentiality of Working Group discussions and deliberations (unless otherwise agreed in advance, and subject to the constraints of the Official Information Act);
 - adopting a constructive approach;
 - avoiding repetition of earlier deliberations, particularly where agreement has already been reached;
 - facilitating an atmosphere of honesty, openness and trust;
 - respecting the role of the Chair; and
 - listening to the views of others, and treating them with respect.
5. Participants in Working Group meetings will be expected to declare their sector affiliations and contractual relationships to the research under review, and to declare any substantial conflicts of interest related to any particular issue or scientific conclusion.
6. Working Group participants must adhere to the requirements of independence, impartiality and objectivity listed under the Peer Review Criteria in the Research Standard (pages 12–16). It is understood that Working Group participants will often be representing particular sectors and interest groups, and may be expressing the views of those groups. However, when participating in the review of science information, representatives are expected to step aside from their sector affiliations, and to ensure that individual and sector views do not result in bias in the science information and conclusions.
7. Participants in each Working Group will have access to the corresponding sections of the Science Working Group website including the Working Group papers and other information provided in those sections. Access to Science Working Group websites will generally be restricted to those who have a reasonable expectation of attending at least one meeting of a given Science Working Group each year.
8. Working Group members who do not adhere to the standards of participation (paragraph 4), or who use Working Group papers and related information inappropriately (see paragraph 10), may be requested by the Chair to leave a particular meeting or to refrain from attending one or more future meetings. In more serious instances, members may be removed from the Working Group membership and denied access to the Working Group website for a specified period of time.

Working Group papers and related information

9. Working Group papers will be posted on the FNZ -Fisheries website prior to meetings if they are available. As a general guide, PowerPoint presentations and draft or discussion papers should be available at least two working days before a meeting, and near-final papers should be available at least five working days before a meeting if the Working Group is expected to agree to the paper. However, it is also likely that some papers will be made available for the first time during the meeting due to time constraints. If a paper is not available for sufficient time before the meeting, the Chair may provide for additional time following the meeting for additional comments from Working Group members.
10. Working Group papers are “works in progress” intended to facilitate the discussion of analyses by the Working Groups. They often contain preliminary results that are receiving peer review for the first time and, as such, may contain errors or preliminary analyses that will be superseded by more rigorous work. **For these reasons, no-one may release the papers or any information contained in these papers to external parties. In general, Working Group papers should not be cited.** Exceptions may be made in rare instances by obtaining permission in writing from the Principal Advisor Fisheries Science, and the authors of the paper. It is also anticipated that Working Group participants who are representing others at a particular Working Group meeting or series of such meetings may wish to communicate preliminary results to the people they are representing. Participants, along with recipients of the information, are required to exercise discretion in doing this, and to guard against preliminary results being made public.
11. From time to time, FNZ commissions external reviews of analyses, models or issues. Terms of Reference for these reviews and the names of external reviewers may be provided to the Working Group for information or feedback. It is extremely important to the proper conduct of these reviews that all contact with the reviewers is through the Chair of the Working Group or the Principal Advisor Fisheries Science. Under no circumstances should Working Group members approach reviewers directly until after the final report of the review has been published.

Working Group meetings

12. Meetings will take place as required, generally January–April and July–November for FAWGs and throughout the year for other Working Groups (AEWG, BRAG, HMSWG, SPACWG, ANTWG and MAFWG).
13. A quorum will be reached when the Chair, the designated presenter, and at least three other technical experts are present. In the absence of a quorum, the Chair may decide to proceed as a sub-group, with outcomes being discussed with the wider Working group via email or taken forward to the next meeting at which a quorum is formed.
14. The Chair is responsible for deciding, with input from the entire Working Group, but focussing primarily on the technical discussion and the views of technical expert members:
 - the quality and acceptability of the information and analyses under review;
 - the way forward to address any deficiencies;
 - the need for any additional analyses;
 - contents of research reports, Working Group reports and AEBAR chapters;
 - choice of best models and sensitivity analyses to be presented; and
 - the status of the stocks, or the status/performance in relation to any relevant environmental standards or targets.
15. The Chair is responsible for facilitating a consultative and collaborative discussion.

16. Working Group meetings will be run formally, with agendas pre-circulated, and formal records kept of recommendations, conclusions and action items.
17. A record of recommendations, conclusions and action items will be posted on the FNZ - Fisheries website after each meeting has taken place.
18. Data upon which analyses presented to the Working Groups are based must be provided to FNZ in the appropriate format and level of detail in a timely manner (i.e. the data must be available and accessible to FNZ ; however, data confidentiality concerns mean that some data may not necessarily be made available to Working Group members).
19. Working Group processes will be evaluated periodically, with a view to identifying opportunities for improvement. Terms of Reference and the Membership and Protocols may be updated as part of this review.
20. FNZ fisheries scientists and science officers will provide administrative support to the Working Groups.

Information Quality Ranking

21. Science Working Groups are required to rank the quality of research and science information that is intended or likely to inform fisheries management decisions, in accordance with the science information quality ranking guidelines in the Research Standard (pages 21–23). Information quality rankings should be documented in Working Group reports and, where appropriate, in Status of Stock summary tables. Note that:
 - Working Groups are not required to rank all research projects and analyses, but key pieces of information that are expected or likely to inform fisheries management decisions, the development of environmental decisions or the formulation of relevant policy should receive a quality ranking;
 - explanations substantiating the quality rankings will be included in Working Group reports. In particular, the quality shortcomings and concerns for moderate/mixed and low quality information should be documented; and
 - the Chair, working with participants, will determine which pieces of information require a quality ranking. Not all information resulting from a particular research project would be expected to achieve the same quality rank, and different quality ranks may be assigned to different components, conclusions or pieces of information resulting from a particular piece of research.

Record-keeping

22. The overall responsibility for record-keeping rests with the Chair of the Working Group, and includes:
 - keeping notes on recommendations, conclusions and follow-up actions for all Working Group meetings, and to ensure that these are available to all members of the Working Group and the Principal Advisor Fisheries Science in a timely manner. If full agreement on the recommendations or conclusions cannot readily be reached amongst technical experts, then the Chair will document the extent to which agreement or consensus was achieved, and record and attribute any residual disagreement in the meeting notes; and
 - compiling a list of generic assessment issues and specific research needs for each stock, species or environmental issue under the purview of the Working Group, for use in subsequent research planning processes.

Fisheries Assessment Working Groups: Membership 2018

Northern and Southern Inshore Working Group

Convenor: Marc Griffiths

Members: John Annala, Suze Baird, Josh Barclay, Mike Beentjes, Nokome Bentley Anthony Brett, Bill Chisholm, Tom Clark, Phil Clow, Matt Dunn, Jack Fenaughty, Malcolm Francis, Allen Frazer, Laura Furneaux, Mark Geytenbeek, Bruce Hartill, George Harvey, Nicholas Hay, Jeremy Helson, Sonja Hempel, John Holdsworth, Rosie Hurst, John Jameson, Terese Kendrick, Brigid Kerrigan, Adam Langley, Laws Lawson, Greg Lydon, Dan MacGibbon, Andy McKay, Jeremy McKenzie, Alicia McKinnon, David Middleton, Connor Nielson, Richard O'Driscoll, Tom Oosting, Yvan Papa, Steve Parker, Darren Parsons, Nathan Reid, Peter Ritchie, Kirsten Rogers, Max Schofield, Carol Scott, Bill Smellie, Paul Starr, Kevin Sullivan, Ali Undorf-Lay, John Taunton-Clark, Peter van Kampen, Cameron Walsh, Tamara Wells, Oliver Wilson.

Species:	Anchovy	Jack Mackerel (JMA 1)	Rough Skate
	Barracouta (BAR 1)	John dory	School shark
	Bluenose	Kahawai	Sea perch (SPE1,2,8,9)
	Blue cod	Kingfish	Smooth Skate
	Blue mackerel (EMA 1&2)	Leatherjacket	Snapper
	Blue moki	Ling (LIN 1&2)	Spiny dogfish (SPD1,3,7,8)
	Blue warehou	Parore	Sprats
	Butterfish	Pilchard	Stargazer
	Elephant fish	Porae	Tarakihi
	Flatfish	Red cod	Trevally
	Gemfish (SKI 1&2)	Red gurnard	Trumpeter
	Garfish	Red snapper	Yellow-eyed mullet
	Grey mullet	Rig	
	Groper	Ribaldo (RIB 1, 2 & 9)	

Shellfish Working Group

Convenors: Julie Hills and Marine Pomarède

Members: John Annala, Matt Baird, Roger Belton, Katrin Berkenbusch, Anthony Brett, Donal Boyle, Barry Chandler, Bill Chisholm, Jeremy Cooper, Jack Fenaughty, Rich Ford, Allen Frazer, Laura Furneaux, Mark Geytenbeek, Nicholas Hay, Sonja Hempel, Brian Inns, Kath Large, John Leader, Craig Marsh, Tom McCowan, Andy McKenzie, David Middleton, Russell Millar, Kiri Morgan, Reyn Naylor, Phil Neubauer, Duncan Petrie, John Reid, Alan Riwaka, Adam Slater, Storm Stanley, Paul Starr, Ian Tuck, Peter van Kampen, Adam Watson, D'Arcy Webber, Richard Wells, Lindsey White, James Williams.

Species:	Cockles	Kina	Triangle shell
	Deepwater crab	Paddle crab	Ringed dosinia
	Dredge oysters	Paua	Fine (Silky) dosinia
	Deepwater (king) clam (Geoduc)	Pipi	Scallop
	Green-lipped mussel	Red crab	Scampi
	King crab	Queen scallop	Surf clam
	Friiled venus shell	Deepwater tuatua	Toheroa
	Knobbed whelk	Giant spider crab	Tuatua
	Sea cucumber	Trough shell	Horse mussel
		Large trough shell	

Antarctic Working Group

Convenor: Marine Pomarède

Members: David Bowden, Martin Cryer, Alistair Dunn, Regina Eisert, Jack Fenaughty, Debbie Freeman, Greg Funnell, Stuart Hanchet, Kath Large, Alexandra Macdonald, Sophie Mormede, Conor Neilson, Richard O'Driscoll, Steve Parker, Matt Pinkerton, Brodie Plum, Darryn Shaw, Andy Smith, Darren Stevens, Kalolaine Vaipuna, Barry Weeber, Bob Zuur.

Species: Antarctic toothfish

Deepwater Working Group

Convenors: Alistair Dunn

Members: Owen Anderson, Karen Baird, Suze Baird, Sira Ballara, Tiffany Bock, Malcolm Clark, George Clement, Patrick Cordue, Matt Dunn, Jack Fenaughty, David Foster, Sharleen Gargiulo, Cara Halford, Stuart Hanchet, Nicholas Hay, Lyndsey Holland, Peter Horn, Dean Jurasovich, Brigid Kerrigan, Daniel Kerrigan, David Kopp, Yoann Ladroit, Adam Langley, Kath Large, Laws Lawson, Mary Livingston, Greg Lydon, Pamela Mace, Dan MacGibbon, Craig Marsh, Michelle Masi, Andy McKay, Andy McKenzie, Peter McMillan, David Middleton, Sophie Mormede, Richard O'Driscoll, Graham Patchell, Jim Roberts, Tim Ryan, Alexandre Schimel, Andy Smith, Paul Starr, John Taunton-Clark, Rob Tilney, Geoff Tingley, Rob Tinkler, Di Tracey, Barry Weeber, Richard Wells, Inge Wisselink, Jingjing Zhang

Species:	Alfonsino	Ling
	Arrow squid	Lookdown dory
	Barracouta (BAR 4,5 & 7)	Orange roughy
	Black cardinalfish	Redbait
	Black oreo	Ribaldo (RIB 3 – 8)
	Blue mackerel (EMA 3&7)	Rubyfish
	Frostfish (FRO 3 – 9)	Sea perch (SPE 3 – 7)
	Gemfish (SKI 3&7)	Silver warehou
	Dark ghost shark (GSH 4 – 6)	Smooth oreo
	Pale ghost shark	Southern blue whiting
	Hake	Spiny dogfish (SPD 4&5)
	Hoki	White warehou
	Jack Mackerel (JMA 3&7)	

Eel Working Group

Convenor: Marc Griffiths

Members: Mike Beentjes, Jacques Boubée, Anthony Brett, Bill Chisholm, Shannan Crowe, Allen Frazer, Eric Graynoth, Nicholas Hay, Simon Hoyle, Mark James, John Jameson, Laws Lawson, Pamela Mace, Michael Martin, David Middleton,, Duncan Petrie, Paul Starr, John Taunton-Clark, Dave West, Erica Williams

Species: Freshwater eels

Stock Assessment Methods Working Group

Convenor: Pamela Mace

Members: John Annala, Paul Breen, Patrick Cordue, Ian Doonan, Alistair Dunn, Matt Dunn, Marc Griffiths, Julie Hills, Simon Hoyle, Adam Langley, Kath Large, Andy McKenzie, Jeremy McKenzie, Craig Marsh, David Middleton, Marine Pomerade, Sophie Mormede, Phil Neubauer, Richard O'Driscoll, Nokuthaba Sibanda, Paul Starr, Daryl Sykes, John Taunton-Clark, Geoff Tingley, D'Arcy Webber

Fisheries Data Working Group

Convenor: Kim George

Members: Cara Halford, Jeremy Helson, David Kopp, Adam Langley, Greg Lydon, Pamela Mace, Alicia McKinnon, David Middleton, Paul Starr, Daryl Sykes, John Taunton-Clark, Finlay Thompson

Marine Amateur Fisheries Working Group

Convenors: Martin Cryer

Members: Sonja Austin, Josh Barclay, Steve Beatson, Marty Bowers, Erin Breen, Paul Breen, Tom Clark, Mark Edwards, Laura Furneaux, Mark Geytenbeek, Cara Halford, Bruce Hartill, Nicholas Hay, Jeremy Helson, John Holdsworth, Brigid Kerrigan, David Kopp, Laws Lawson, Andy McKay, Alicia McKinnon, David Middleton, Trish Rea, Merrill Rudd, Carol Scott, Paul Starr, Daryl Sykes, John Taunton-Clark, Scott Tindale, D'Arcy Webber, Oliver Wilson.

Aquatic Environment Working Group

Convenors: Rich Ford and Nathan Walker

Members: Ed Abraham, Owen Anderson, Sonja Austin, Hilary Ayrton, Karen Baird, Suze Baird, Barry Baker, Scott Baker, Sira Ballara, Joshua Baller, Josh Barclay, Steve Beatson, Katrin Berkenbusch, Tiffany Bock, Lesley Bolton-Ritchie, Laura Boren, Erin Breen, Anthony Brett, Niall Broekhuizen, Ian Brown, Paul Breen, Tania Cameron, Simon Childerhouse, Bill Chisholm, Malcolm Clark, Tom Clark, Katie Clemens-Seely, Deanna Clement, George Clement, Igor Debski, Peter Dillingham, Matt Dunn, Charles Edwards, Mark Edwards, Jack Fenaughty, David Foster, Chris Francis, Malcolm Francis, Allen Frazer, Laura Furneaux, Sharleen Gargiulo, Mark Geytenbeek, William Gibson, Neil Gilbert, Kim Goetz, Cara Halford, Nicholas Hay, Jeremy Helson, Kristina Hillock, John Holdsworth, Lyndsey Holland, Brigid Kerrigan, Daniel Kerrigan, Kirstie Knowles, David Kopp, Jo Lambie, Todd Landers, Laws Lawson, Amanda Leathers, Mary Livingston, Carolyn Lundquist, Dave Lundquist, Greg Lydon, Darryl MacKenzie, Gemma McGrath, Andy McKay, Andy McKenzie, Alicia McKinnon, Peter McMillan, David Middleton, Janice Molloy, Kiri Morgan, Sophie Mormede, Phil Neubauer, Richard O'Driscoll, Jenny Oliver, Tracey Osborne, Enrique Pardo, Steve Parker, Darren Parsons, Mike Patrick, Johanna Pierre, Trish Rea, Yvan Richard, Peter Ritchie, Jim Roberts, Christine Rose, Charles Rowe, Carol Scott, Liz Slooten, Andy Smith, Paul Starr, John Taunton-Clark, David Thompson, Finlay Thompson, Rob Tilney, Geoff Tingley, Rob Tinkler, Di Tracey, Ian Tuck, Anton Van Helden, Adam Watson, D'Arcy Webber, Barry Weeber, Richard Wells, Tamar Wells, James Williams, Oliver Wilson, Inge Wisselink, Andrew Wright, Jingjing Zhang

Guide to Biological Reference Points for Fisheries Assessment Meetings

The Guide to Biological Reference Points was originally developed by a Stock Assessment Methods Working Group in 1988, with the aim of defining commonly used terms, explaining underlying assumptions, and describing the biological reference points used in fisheries assessment meetings and associated reports. However, this document has not been substantially revised since 1992 and the methods described herein, while still used in several assessments, have been replaced with other approaches in a number of cases. Some of the latter approaches are described in the Harvest Strategy Standard for New Zealand Fisheries and the associated Operational Guidelines, and are being further developed in various Fisheries Assessment Working Groups and the current Stock Assessment Methods Working Group.

Here, methods of estimation appropriate to various circumstances are given for two levels of yield: Maximum Constant Yield (**MCY**) and Current Annual Yield (**CAY**), both of which represent different forms of maximum sustainable yield (**MSY**). The relevance of these to the setting of Total Allowable Catches (TACs) is discussed.

Definitions of **MCY** and **CAY**

The Fisheries Act 1996 defines Total Allowable Catch in terms of maximum sustainable yield (**MSY**). The definitions of the biological reference points, **MCY** and **CAY**, derive from two ways of viewing **MSY**: a static interpretation and a dynamic interpretation. The former, associated with **MCY**, is based on the idea of taking the same catch from the fishery year after year. The latter interpretation, from which **CAY** is derived, recognises that fish populations fluctuate in size from year to year (for environmental and biological, as well as fishery, reasons) so that to get the best yield from a fishery it is necessary to alter the catch every year. This leads to the idea of maximum average yield (**MAY**) which is how fisheries scientists generally interpret **MSY** (Ricker 1975).

The definitions are:

MCY – Maximum Constant Yield

The maximum constant catch that is estimated to be sustainable, with an acceptable level of risk, at all probable future levels of biomass.

and

CAY – Current Annual Yield

The one-year catch calculated by applying a reference fishing mortality, F_{REF} , to an estimate of the fishable biomass present during the next fishing year. F_{REF} is the level of (instantaneous) fishing mortality that, if applied every year, would, within an acceptable level of risk, maximise the average catch from the fishery.

Note that **MCY** is dependent to a certain extent on the current state of the fish stock. If a stock is fished at the **MCY** level from a virgin state then over the years its biomass will fluctuate over a range of levels depending on environmental conditions, abundance of predators and prey, etc. For stock sizes within this range the **MCY** remains unchanged (though our estimates of it may well be refined). If the current state of the stock is below this range the **MCY** will be lower.

The strategy of applying a constant fishing mortality, F_{REF} , from which the **CAY** is derived each year is an approximation to a strategy which maximises the average yield over time. For the purposes of this document the **MAY** is the long-term average annual catch when the catch each year is the **CAY**. With perfect knowledge it would be possible to do better by varying the fishing mortality from year to year. Without perfect knowledge, adjusting catch levels by a **CAY** strategy as stock size varies is probably the best practical method of maximising average yield. Appropriate values for F_{REF} are discussed below.

What is meant by an “acceptable level of risk” for **MCYs** and **CAYs** is intentionally left undefined here. For most stocks our level of knowledge is inadequate to allow a meaningful quantitative assessment of risk. However, we have two qualitative sources of information on risk levels: the experience of fisheries

scientists and managers throughout the world, and the results of simulation exercises such as those of Mace (1988a). Information from these sources is incorporated, as much as is possible, in the methods given below for calculating *MCY* and *CAY*.

It is now well known that *MCY* is generally less than *MAY* (see, e.g., Doubleday 1976, Sissenwine 1978, Mace 1988a). This is because *CAY* will be larger than *MCY* in the majority of years. However, when fishable biomass becomes low (through overfishing, poor environmental conditions, or a combination of both), *CAY* will be less than *MCY*. This is true even if the estimates of *CAY* and *MCY* are exact. The following diagram shows the relationships between *CAY*, *MCY* and *MAY*.




Figure 1: Relationship between *CAY*, *MCY* and *MAY*.

In this example *CAY* represents a constant fraction of the fishable biomass, and so (if it is estimated and applied exactly) it will track the fish population exactly. *MAY* is the average over time of *CAY*. The reason *MCY* is less than *MAY* is that *MCY* must be low enough so that the fraction of the population removed does not constitute an unacceptable risk to the future viability of the population. With an *MCY* strategy, the fraction of a population that is removed by fishing increases with decreasing stock size. With a *CAY* strategy, the fraction removed remains constant. A constant catch strategy at a level equal to the *MAY*, would involve a high risk at low stock sizes.

Relationship Between *MCY*, *CAY*, TAC and Total Allowable Commercial Catch (TACC)

The TAC covers all mortality to a fish stock caused by human activity, whereas the TACC includes only commercial catch. *MCY* and *CAY* are reference points used to evaluate whether the current stock size can support the current TAC and/or TACC. It should not be assumed that the TAC and/or TACC will be equal to either one of these yields. There are both legal and practical reasons for this.

Legally, we are bound by the Fisheries Act 1996. In setting or varying any TACC for any quota management stock, 'the Minister shall have regard to the total allowable catch for that stock and shall allow for –

- (a) The following non-commercial fishing interests in that stock, namely –
 - (i) Maori customary non-commercial fishing interests; and
 - (ii) Recreational interests; and
- (b) All other mortality to that stock caused by fishing.

From a practical point of view it must be acknowledged that the concepts of *MCY* and *CAY* are directly applicable only in idealised management regimes. The *MCY* could be used in a regime where a catch level was to be set for once and for all; our system allows changes to be made if, the level is found to be too low or too high.

With a *CAY* strategy the yield would probably change every year. Even if there were no legal impediments to following a *CAY* strategy, the fishing industry's desire for stability may be a sufficient reason to make TACC changes only when the need is pressing.

Natural and Fishing Mortality

Before describing how to calculate **MCY** and **CAY** we must discuss natural and fishing mortality, which are used in these calculations. Both types of mortality are expressed as instantaneous rates (thus, over n years a total mortality Z will reduce a population of size B to size Be^{-nZ} , ignoring recruitment and growth). Units for mortalities are 1/year.

Natural mortality

Methods of estimating natural mortality, M , are reviewed by Vetter (1988). When a lack of data rules out more sophisticated methods, M may be estimated by the formula,

$$M = \frac{\log_e(p)}{A}$$

where p is the proportion of the population that reaches age A (or older) in an unexploited stock. p is often set to 0.01, when A is the "maximum age" observed. Other values for p may be chosen dependent on the fishing history of the stock. For example, in an exploited stock the maximum observed age may correspond to a value of $p = 0.05$, or higher. For a discussion of the method see Hoenig (1983).

Reference Fishing Mortalities

Reference fishing mortalities in widespread use include $F_{0.1}$, F_{MSY} , F_{MAX} , F_{MEY} , and M .

The most common reference fishing mortality used in the calculation of **CAY** (and, in some cases, **MCY**) is $F_{0.1}$ (pronounced 'F zero point one'). This is used as a basis for fisheries management decisions throughout the world and is widely believed to produce a high level of yield on a sustainable basis (Mace 1988b). It is estimated from a yield per recruit analysis as the level of fishing mortality at which the slope of the yield-per-recruit curve is 0.1 times the slope at $F = 0$. If an estimate of $F_{0.1}$ is not available an estimate of M may be substituted.

F_{MAX} , the fishing mortality that produces the maximum yield per recruit. It may be too high as a target fishing mortality because it does not account for recruitment effects (e.g. recruitment declining as stock size is reduced). However, it may be a valid reference point for those fisheries that have histories of sustainable fishing at this level.

F_{MSY} , the fishing mortality corresponding to the deterministic **MSY**, is another appropriate reference point. F_{MSY} may be estimated from a surplus production model, or a combination of yield per recruit and stock recruitment models.

When economic data are available it may be possible to calculate F_{MEY} the fishing mortality corresponding to the maximum (sustainable) economic yield.

Every reference fishing mortality corresponds to an equilibrium or long-run average stock biomass. This is the biomass which the stock will tend towards or randomly fluctuate around, when the reference fishing mortality is applied constantly. The fluctuations will be caused primarily by variable recruitment. It is necessary to examine the equilibrium stock biomass corresponding to any candidate reference fishing mortality.

A reference fishing mortality which corresponds to a low stock biomass may be undesirable if the low biomass would lead to an unacceptable risk of stock collapse. For fisheries where this applies a lower reference fishing mortality may be appropriate.

Natural Variability Factor

Fish populations are naturally variable in size because of environmental variability and associated fluctuations in the abundance of predators and food. Computer simulations (e.g., Mace 1988a) have shown that, all other things being equal, the **MCY** for a stock is inversely related to the degree of natural variability in its abundance. That is, the higher the natural variability, the lower the **MCY**.

The natural variability factor, **c**, provides a way of incorporating the natural variability of a stock's biomass into the calculation of **MCY**. It is used as a multiplying factor in method 5 below. The greater the variability in the stock, the lower is the value of **c**. Values for **c** should be taken from the table below and are based on the estimated mean natural mortality rate of the stock. It is assumed that because a stock with a higher natural mortality will have fewer age-classes it will also suffer greater fluctuations in biomass. The only stocks for which the table should be deviated from are those where there is evidence that recruitment variability is unusually high or unusually low.

Natural mortality rate <i>M</i>	Natural variability factor <i>c</i>
<0.05	1.0
0.05-0.15	0.9
0.16-0.25	0.8
0.26-0.35	0.7
>0.35	0.6

Methods of Estimating **MCY**

It should be possible to estimate **MCY** for most fish stocks (with varying degrees of confidence). For some stocks, only conservative estimates for **MCY** will be obtainable (e.g., some applications of Method 4) and this should be stated. For other stocks it may be impossible to estimate **MCY**. These stocks include situations in which: the fishery is very new; catch or effort data are unreliable; strong upwards or downwards trends in catch are not able to be explained by available data (e.g., by trawl survey data or by catch per unit effort data).

When catch data are used in estimating **MCY** all catches (commercial, illegal, and non-commercial) should be included if possible. If this is not possible and the excluded catch is thought to be a significant quantity, then this should be stated.

The following examples define **MCY** in an operational context with respect to the type, quality and quantity of data available. Knowledge about the accuracy or applicability of the data (e.g., reporting anomalies, atypical catches in anticipation of the introduction of the Quota Management System) should play a part in determining which data sets are to be included in the analysis.

As a general rule it is preferable to apply subjective judgements to input data rather than to the calculated **MCYs**. For example, rather than saying “with the official catch statistics the **MCY** is **X** tonnes, but we think this is too high because the catch statistics are wrong” it would be better to say “we believe (for reasons given) that the official statistics are wrong and the true catches were probably such and such, and the **MCY** based on these catches is **Y** tonnes”.

Background information on the rationale behind the following calculation methods can be found in Mace (1988a) and other scientific papers listed at the end of this document.

New fisheries

$$MCY = 0.25F_{0.1}B_0$$

where B_0 is an estimate of virgin recruited biomass. If there are insufficient data to conduct a yield per recruit analysis $F_{0.1}$ should be replaced with an estimate of natural mortality (M). Tables 1–3 in Mace (1988b) show that $F_{0.1}$ is usually similar to (or sometimes slightly greater than) M .

It may appear that the estimate of MCY for new fisheries is overly conservative, particularly when compared to the common approximation to MSY of $0.5MB_0$ (Gulland 1971). However various authors (including Beddington & Cooke 1983; Getz et al 1987; Mace 1988a) have shown that $0.5MB_0$ often overestimates MSY , particularly for a constant catch strategy or when recruitment declines with stock size. Moreover it has often been observed that the development of new fisheries (or the rapid expansion of existing fisheries) occurs when stock size is unusually large, and that catches plummet as the accumulated biomass is fished down.

It is preferable to estimate MCY from a stochastic population model (Method 5), if this is possible. The simulations of Mace (1988a) and Francis (1992) indicate that the appropriate factor to multiply $F_{0.1}B_0$ may be somewhat higher or somewhat lower than **0.25**. This depends primarily on the steepness of the assumed stock recruitment relationship (*see* Mace & Doonan 1988 for a definition of steepness).

New fisheries become developed fisheries once F has approximated or exceeded M for several successive years, depending on the lifespan of the species.

2. Developed fisheries with historical estimates of biomass

$$MCY = 0.5F_{0.1}B_{AV}$$

where B_{AV} is the average historical recruited biomass, and the fishery is believed to have been fully exploited (i.e., fishing mortality has been near the level that would produce MAY). This formulation assumes that $F_{0.1}$ approximates the average productivity of a stock.

As in the previous method an estimate of M can be substituted for $F_{0.1}$ if estimates of $F_{0.1}$ are not available.

3. Developed fisheries with adequate data to fit a population model

$$MCY = \frac{2}{3}MSY$$

where MSY is the deterministic maximum equilibrium yield.

This reference point is slightly more conservative than that adopted by several other stock assessment agencies (e.g. ICES, CAFSAC) that use as a reference point the equilibrium yield corresponding to 2/3 of the fishing effort (fishing mortality) associated with the deterministic equilibrium MSY .

If it is possible to estimate MSY then it is generally possible to estimate MCY from a stochastic population model (Method 5), which is the preferable method. The simulations of Mace (1988a) and Francis (1992) indicate that the appropriate factor to multiply MSY varies between about **0.6** and **0.9**. This depends on various parameters of which the steepness of the assumed stock recruitment relationship is the most important.

If the current biomass is less than the level required to sustain a yield of 2/3 MSY then

$$MCY = \frac{2}{3}CSP$$

where CSP is the deterministic current surplus production.

4. Catch data and information about fishing effort (and/or fishing mortality), either qualitative or quantitative, without a surplus production model

$$MCY = cY_{AV}$$

where c is the natural variability factor (defined above) and Y_{AV} is the average catch over an appropriate period.

If the catch data are from a period when the stock was fully exploited (i.e. fishing mortality near the level that would produce MAY), then the method should provide a good estimate of MCY . In this case, $Y_{AV} = MAY$. If the population was under-exploited the method gives a conservative estimate of MCY .

Familiarity with stock demographics and the history of the fishery is necessary for the determination of an appropriate period on which to base estimates of Y_{AV} . The period chosen to perform the averaging will depend on the behaviour of the fishing mortality or fishing effort time series, the prevailing management regime, the behaviour of the catch time series, and the lifespan of the species.

The period should be selected so that it contains no systematic changes in fishing mortality (or fishing effort, if this can be assumed to be proportional to fishing mortality). Note that for species such as orange roughy, where relatively static aggregations are fished, fishing mortality cannot be assumed to be proportional to effort. If catches during the period are constrained by a TACC then it is particularly important that the assumption of no systematic change in fishing mortality be adhered to. The existence of a TACC does not necessarily mean that the catch is constrained by it.

The period chosen should also contain no systematic changes in catch. If the period shows a systematic upward (or downward) trend in catches then the MCY will be under-estimated (over-estimated). It is desirable that the period be equal to at least half the exploited life span of the fish.

5. Sufficient information for a stochastic population model

This is the preferred method for estimating MCY but it is the method requiring the most information. It is the only method that allows some specification of the risk associated with an MCY .

The simulations in Mace (1988a) and Breen (1989) provide examples of the type of calculations necessary for this method. A trial and error procedure can be used to find the maximum constant catch that can be taken for a given level of risk. The level of risk may be expressed as the probability of stock collapse within a specified time period. At the moment the Ministry of Fisheries has no standards as to how stock collapse should be defined for this purpose, what time period to use, and what probability of collapse is acceptable. These will be developed as experience is gained with this method.

Methods of Estimating *CAY*

It is possible to estimate *CAY* only when there is adequate stock biomass data. In some instances relative stock biomass indices (e.g., catch per unit effort data) and relative fishing mortality data (e.g., effort data) may be sufficient. *CAY* calculated by method 1 includes non-commercial catch.

If method 2 is used and it is not possible to include a significant non-commercial catch, then this should be stated.

1. Where there is an estimate of current recruited stock biomass, *CAY* may be calculated from the appropriate catch equation. Which form of the catch equation should be used will depend on the way fishing mortality occurs during the year. For many fisheries it will be a reasonable approximation to assume that fishing is spread evenly throughout the year so that the Baranov catch equation is appropriate and *CAY* is given by

$$CAY = \frac{F_{ref}}{F_{ref} + M} (1 - e^{-(F_{ref}+M)}) B_{beg}$$

Where B_{BEG} is the projected stock biomass at the beginning of the fishing year for which the *CAY* is to be calculated and F_{REF} is the reference fishing mortality described above.

If most of the fishing mortality occurs over a short period each year it may be better to use one of the following equations:

$$CAY = (1 - e^{-F_{ref}}) B_{beg}$$

$$CAY = (1 - e^{-F_{ref}}) e^{-\frac{M}{2}} B_{beg}$$

$$CAY = (1 - e^{-F_{ref}}) e^{-M} B_{beg}$$

where the first equation is used when fishing occurs at the beginning of the fishing year, the second equation when fishing is in the middle of the year, and the third when fishing is at the end of the year.

It is important that the catch equation used to calculate *CAY* and the associated assumptions are the same as those used in any model employed to estimate stock biomass or to carry out yield per recruit analyses. Serious bias may result if this criterion is not adhered to. The assumptions and catch equations given here are by no means the only possibilities.

The risk associated with the use of a particular F_{REF} may be estimated using simulations.

2. Where information is limited but the current (possibly unknown) fishing mortality is thought to be near the optimum, there are various "status quo" methods which may be applied. Details are available in Shepherd (1984, 1991) and Pope (1983).

FOR FURTHER INFORMATION

- Beddington, J R; Cooke, J G (1983) The potential yield of fish stocks. *FAO Fisheries Technical Paper* No. 242, Rome. 47 p.
- Beddington, J R; May, R M (1977) Harvesting natural populations in a randomly fluctuating environment. *Science* 197: 463–465.
- Breen, P A (1989) Rock lobster stock assessment 1989. New Zealand Fisheries Assessment Research Document 1989/6. (Unpublished document held by NIWA library, Wellington.)
- Deriso, R B (1985) Risk adverse harvesting strategies. pp 65–73 in M. Mangel (Ed.): Resource Management. Lecture Notes in Biomathematics 61.
- Doubleday, W C (1976) Environmental fluctuations and fisheries management. *International Commission for the Northwest Atlantic Fisheries, Selected Papers* (1): 141–150.
- Francis, R I C C (1992) Recommendations concerning the calculation of maximum constant yield (MCY) and current annual yield (CAY). New Zealand Fisheries Assessment Research Document 1992/8. (Unpublished document held by NIWA library, Wellington.)
- Gatto, M; Rinaldi, S (1976) Mean value and variability of fish catches in fluctuating environments. *Journal of the Fisheries Research Board of Canada* 33: 189–193.
- Getz, W M; Francis, R C; Swartzman, G L (1987) On managing variable marine fisheries. *Canadian Journal of Fisheries and Aquatic Sciences* 44: 1370–1375.
- Gulland, J A (1971) (comp.) The Fish Resources of the Ocean. West Byfleet, Surrey, Fishing News (Books) Ltd., for FAO, 255pp. Rev. ed. of *FAO Fisheries Technical Paper* (97): 425p. (1970).
- Hoenig, J M (1983) Empirical use of longevity data to estimate mortality rates. *Fisheries Bulletin*. 81: 898–903.
- Kirkwood, G P (1981) Allowing for risks in setting catch limits based on *MSY*. *Mathematical Biosciences* (53): 119–129.
- Mace, P M (1988a) The relevance of *MSY* and other biological references points to stock assessments in New Zealand. New Zealand Fisheries Assessment Research Document 1988/30. 41 p. (Unpublished document held by NIWA library, Wellington.)
- Mace, P M (1988b) A survey of stock assessment methods and results. New Zealand Fisheries Assessment Research Document 1988/6. 35 p. (Unpublished document held by NIWA library, Wellington.)
- Mace, P M; Doonan, I J (1988) A generalised bioeconomic simulation model for fish population dynamics. New Zealand Fisheries Assessment Research Document 1988/4. 51 p. (Unpublished document held by NIWA library, Wellington.)
- May, R M; Beddington, J R; Horwood, J W; Shepherd, J G (1978) Exploiting natural populations in an uncertain world. *Mathematical Biosciences* 42: 219–252.
- New Zealand Ministry of Fisheries (2008) Harvest Strategy Standard for New Zealand fisheries. 25 p. Available at <http://fs.fish.govt.nz/Page.aspx?pk=61&tk=208&se=&sd=Asc&filSC=&filAny=False&filSrc=False&filLoaded=False&filDCG=9&filDC=0&filST=&filYr=0&filAutoRun=1>.
- New Zealand Ministry of Fisheries (2011) Operational Guidelines for New Zealand's Harvest Strategy Standard Revision 1. 78 p. Available at http://fs.fish.govt.nz/Doc/22847/Operational_Guidelines_for_HSS_rev_1_Jun_2011.pdf.ashx.
- Pope, J G (1983) Analogies to the status quo TACs: their nature and variance, pp. 99–113 in W.G. Doubleday and D. Rivard, Sampling commercial catches of marine fish and invertebrates. *Canadian Special Publication of Fisheries and Aquatic Sciences* 66.
- Reed, W J (1983) Recruitment variability and age structure in harvested animal populations. *Mathematical Biosciences* 65: 239–268.
- Reeves, J E (1974) Comparison of long-term yields from catch quotas and effort quotas under conditions of variable recruitment. ICNAF Res. Doc. 74/31.
- Ricker, W E (1975) Computation and Interpretation of Biological Statistics of Fish Populations. *Bulletin of the Fisheries Research Board of Canada*. 382 p.
- Shepherd, J G (1984) Status quo catch estimation and its use in fishery management. International Council for the Exploration of the Sea CM 1984/G: 5
- Shepherd, J G (1991) Simple methods for short-term forecasting of catch and biomass. *ICES Journal of Marine Science* 48: 67–78.
- Sissenwine, M P (1977) The effects of random fluctuations on a hypothetical fishery. ICNAF, Selected Papers (2): 137–144.
- Sissenwine, M P (1978) Is *MSY* an adequate foundation for optimum yield? *Fisheries* 3(6): 22–42.
- Vetter, E F (1988) Estimation of natural mortality in fish stocks: a review. *Fishery Bulletin* 86(1): 25–43.

Guidelines for Status of the Stocks Summary Tables

A new format for Status of the Stocks summaries was developed by the Stock Assessment Methods Working Group over the period February-April 2009. The purpose of this project was to provide more comprehensive and meaningful information for fisheries managers, stakeholders and other interested parties. Previously, Status of the Stocks summary sections had not reflected the full range of information of relevance to fisheries management contained in the earlier sections of Plenary reports, and were of variable utility for evaluating stock status and informing fisheries management decisions.

Status of the Stocks summary tables should be constructed for all stocks except those designated as “nominal”; e.g. those with administrative TACs or TACCs (generally less than 10–20 t) or those for which a commercial or non-commercial development potential has not currently been demonstrated. As of November 2014, there were a total of 292 stocks in this classification. The list of nominal stocks can be found at: <http://www.mpi.govt.nz/dmsdocument/19331-nz-nominal-fish-stocks-2017-report>.

In 2012 a number of changes were made to the format for the Status of the Stocks summary tables, primarily for the purpose of implementing the science information quality rankings required by the Research and Science Information Standard for New Zealand Fisheries that was approved in April 2011 (New Zealand Ministry of Fisheries 2011a). At the time, these changes were only applied for Status of Stocks tables updated in 2012. Subsequently, an attempt has been made to revise some of the older tables as well.

In 2013, the format was further modified to require Science Working Groups to make a determination about whether overfishing is occurring, and to further standardise and clarify the requirements for other parts of the table.

It is anticipated that the format of the Status of the Stocks tables will continue to be reviewed, standardised and modified in the future so that it remains relevant to fisheries management and other needs. New formats will be implemented each time stocks are reviewed and as time allows.

The table below provides a template for the Status of the Stocks summaries. The text following the template gives guidance on the contents of most of the fields in the table. Superscript numbers refer to the corresponding numbered paragraph in the following text. [Light blue](#) text provides an example of how the table might be completed.

STATUS OF THE STOCKS TEMPLATE¹

Stock Structure Assumptions²

<insert relevant text>

- Fishstock name³

Stock Status	
Year of Most Recent Assessment	2018
Assessment Runs Presented	Base case model only
Reference Points	Target: 40% B_0 Soft Limit: 20% B_0 Hard Limit: 10% B_0 Overfishing threshold: $F_{40\%B_0}$
Status in relation to Target	B_{2018} was estimated to be 50% B_0 ; Very Likely (> 90%) to be at or above the target
Status in relation to Limits	B_{2018} is Very Unlikely (< 10%) to be below both the soft and hard limits
Status in relation to Overfishing	The fishing intensity in 2014 was Very Unlikely (< 10%) to be above the overfishing threshold [or, Overfishing is Very Unlikely (<10%) to be occurring]

Historical Stock Status Trajectory and Current Status

<insert relevant graphs>

Fishery and Stock Trends

Recent Trend in Biomass or Proxy	Biomass reached its lowest point in 2001 and has since consistently increased.
Recent Trend in Fishing Intensity or Proxy	Fishing intensity reached a peak of $F=0.54$ in 1999, subsequently declining to less than $F=0.2$ since 2006.
Other Abundance Indices	-
Trends in Other Relevant Indicators or Variables	Recent recruitment (2005–2017) is estimated to be near the long-term average.

Projections and Prognosis

Stock Projections or Prognosis	Biomass is expected to stay steady over the next 5 years assuming current (2016–17) catch levels.
Probability of Current Catch or TACC causing Biomass to remain below or to decline below Limits	Soft Limit: Very Unlikely (< 10%) Hard Limit: Very Unlikely (< 10%)
Probability of Current Catch or TACC causing Overfishing to continue or to commence	Very Unlikely (< 10%)

Assessment Methodology and Evaluation

Assessment Type	Level 1 - Full Quantitative Stock Assessment	
Assessment Method	Age-structured CASAL model with Bayesian estimation of posterior distributions	
Assessment Dates	Latest assessment: 2018	Next assessment: 2019
Overall assessment quality rank	1 – High Quality	
Main data inputs (rank)	<ul style="list-style-type: none"> - Research time series of abundance indices (trawl and acoustic surveys) - Proportions at age data from the commercial fisheries and trawl surveys - Estimates of biological parameters 	1 – High Quality 1 – High Quality 1 – High Quality
Data not used (rank)	Commercial CPUE	3 – Low Quality: does not track stock biomass
Changes to Model Structure and Assumptions	None since the 2012 assessment	
Major sources of Uncertainty	<ul style="list-style-type: none"> - The base case model deals with the lack of older fish in commercial catches and surveys by estimating natural mortality at age which results in older fish suffering high natural mortality. However, there is no evidence to validate this outside the model estimates. - Aside from natural mortality, other major sources of uncertainty include stock structure and migration patterns, stock-recruit steepness and natal fidelity assumptions. Uncertainty about the size of recent year classes affects the reliability of stock projections. 	

Qualifying Comments
The impact of the current young age structure of the population on spawning success is unknown.

Fishery Interactions
Main bycatch species are hake, ling, silver warehou and spiny dogfish, with lesser bycatches of ghost sharks, white warehou, sea perch and stargazers. Incidental interactions and associated mortalities are noted for New Zealand fur seals and seabirds. Low productivity species taken in the fishery include basking sharks and deepsea skates.

Guidance on preparing the Status of the Stocks summary tables

1. Everything included in the Status of the Stocks summary table should be derived from earlier sections in the Working Group or Plenary report. No new information should be presented in the summary that was not encompassed in the main text of the Working Group or Plenary report.

Stock Structure Assumptions

2. The current assumptions regarding the stock structure and distribution of the stocks being reported on should be briefly summarised. Where the assessed stock distribution differs from the relevant QMA fishstock(s), an explanation must be provided of how the stock relates to the QMA fishstock(s) it includes.

Stock Status

3. One Status of the Stocks summary table should be completed for each assessed stock or stock complex.
4. Management targets for each stock will be established by fisheries managers. Where management targets have not been established, it is suggested that an interim target of 40% B_0 , or a related B_{MSY} -compatible target (or $F_{40\%}$, or a related target) should be assumed. In most cases, the soft and hard limits should be set at the default levels specified in the Harvest Strategy Standard (20% B_0 for the soft limit and 10% B_0 for the hard limit). Similarly, the overfishing threshold should be set at F_{MSY} , or a related F_{MSY} -compatible threshold. Overfishing thresholds can be expressed in terms of fishing mortality, exploitation rates, or other valid measures of fishing intensity. When agreed reference points have not been established, stock status may be reported against interim reference points.
5. Reporting stock status against reference points requires Working Group agreement on the model run to use as a base case for the assessment. The preference, wherever possible, is to report on the best estimates from a single base case, or to make a single statement that covers the results from a range of cases. In general, ranges or confidence intervals should not be included in the table. Only where more than one equally plausible model run exists, and agreement cannot be reached on a single base case, should multiple runs be reported. This should still be done simply and concisely (e.g. median results only).
6. Where probabilities are used in qualifying a statement regarding the status of the stock in relation to target, limit, or threshold reference levels, the following probability categories and associated verbal descriptions are to be used (IPCC, 2007):

Probability	Description
> 99 %	Virtually Certain
> 90 %	Very Likely
> 60 %	Likely
40–60 %	About as Likely as Not
< 40 %	Unlikely
< 10 %	Very Unlikely
< 1 %	Exceptionally Unlikely

Probability categories and associated descriptions should relate to the probability of being “at or above” biomass targets (or “at or below” fishing intensity targets if these are used), below biomass limits, and above overfishing thresholds. Note, however, that the descriptions and associated probabilities adopted need not correspond exactly to model outputs; rather they should be superimposed with the Working Group’s belief about the extent to which the model fully specifies the probabilities. This is particularly relevant for the “Virtually Certain” and “Exceptionally Unlikely” categories, which should be used sparingly.

7. The status in relation to overfishing can be expressed in terms of an explicit overfishing threshold, or it can simply be a statement about the Working Group’s belief, based on the evidence at hand, about the likelihood that overfishing is occurring (based on, for example, a stock abundance index exhibiting a pronounced recent increase or decline). The probability rankings in the IPCC (2007) table above should be used. Overfishing thresholds can be considered in terms of fishing mortality rates, exploitation rates, or other valid measures of fishing intensity.

Historical Stock Status Trajectory and Current Status

8. This heading should be changed to reflect the graphs that are available to illustrate trends in biomass or fishing intensity (or proxies) and the current stock or fishery status.

Recent Fishery and Stock Trends

9. Recent stock or fishery trends should be reported in terms of stock size and fishing intensity (or proxies for these), respectively. For full quantitative (Level 1) assessments, median results should be used when reporting biomass. Observed trends should be reported using descriptors such as increasing, decreasing, stable, or fluctuating without trend. Where it is considered relevant and important to fisheries management, mention could be made of whether the indicator is moving towards or away from a target, limit, threshold, or long term average.
10. Other Abundance Indices: This section is primarily intended for reporting of trends where a Level 2 (partial quantitative) evaluation has been conducted, and appropriate abundance indices (such as standardised CPUE or survey biomass) are available.
11. Other Relevant Indicators or Variables: This section is primarily intended for reporting of trends where only a Level 3 (qualitative) evaluation has been conducted. Potentially useful indicators might include trends in mean size, size or age composition, or recruitment indices. Catch trends vs TACC may be relevant here, provided these are qualified when other factors are known to have influenced the trends.

Projections and Prognosis

12. These sections should be used to report available information on likely future trends in biomass or fishing intensity or related variables under current (or a range of) catch levels over a period of approximately 3–5 years following the last year in the assessment. If a longer period is used, this must be stated.
13. When reporting probabilities of current catches or TACC levels causing declines below limits, the probability rankings in the IPCC (2007) table above should be used. Results should be reported separately (i.e. split into two rows) if the catch and TACC differ appreciably, resulting in differing conclusions for each level of removals, with the level of each specified. The

timeframe for the projections should be approximately 3–5 years following the last year in the assessment unless a longer period of time is required by fisheries managers.

Assessment Methodology and Evaluation

14. Assessment type: the envisaged Assessment Levels are:

- 1 – Full Quantitative Stock assessment: There is a reliable index of abundance and an assessment indicating status in relation to targets and limits.
- 2 – Partial Quantitative Stock Assessment: An evaluation of agreed abundance indices (e.g. standardised CPUE) or other appropriate fishery indicators (e.g. estimates of F (Z) based on catch-at-age) is available. Indices of abundance or fishing intensity have not been used in a full quantitative stock assessment to estimate stock or fishery status in relation to reference points.
- 3 – Qualitative Evaluation: A fishery characterisation with evaluation of fishery trends (e.g. catch, effort, unstandardised CPUE, or length-frequency information) has been conducted but there is no agreed index of abundance.
- 4 – Low Information Evaluation: There are only data on catch and TACC, with no other fishery indicators.

Management Procedure (MP) updates should be presented in a separate table. In years when an actual assessment is conducted for stocks under MPs, the MP update table should be preceded by a Status of the Stocks summary table.

Table content will vary for these different assessment levels.

Ranking of Science Information Quality

15. The Research and Science Information Standard for New Zealand Fisheries (2011a) specifies (pages 21–23) that the Ministry will implement processes that rank the quality of research and science information used in support of fisheries management decisions. The quality ranking system is:

- 1 – High Quality: information that has been subjected to rigorous science quality assurance and peer review processes as required by this Standard, and substantially meets the key principles for science information quality. Such information can confidently be accorded a high weight in fisheries management decisions. An explanation is not required in the table for high quality information.
- 2 – Medium or Mixed Quality: information that has been subjected to some level of peer review against the requirements of the Standard and has been found to have some shortcomings with regard to the key principles for science information quality, but is still useful for informing management decisions. Such information should be accompanied by a description of its shortcomings.
- 3 – Low Quality: information that has been subjected to peer review against the requirements of the Standard but has substantially failed to meet the key principles for science information quality. Such information should be accompanied by a description of its shortcomings and should not be used to inform management decisions.

One of the key purposes of the science information quality ranking system is to inform fisheries managers and stakeholders of those datasets, analyses or models that are of such poor quality that they should not be used to make fisheries management decisions (i.e. those ranked as “3”). Most other datasets, analyses or models that have been subjected to peer review or staged technical guidance in the Ministry’s Science Working Group processes and have been accepted by these processes should be given the highest score (ranked as “1”). Uncertainty, which is inherent in all fisheries science outputs, should not by itself be used as a reason to score down a research output, unless it has not been properly considered or analysed, or if the uncertainty is so large as to render the results and conclusions meaningless (in which case, the Working Group should consider rejecting the output altogether). A ranking of 2 (medium or

mixed quality) should only be used where there has been limited or inadequate peer review or the Working Group has mixed views on the validity of the outputs, but believes they are nevertheless of some use to fisheries management.

16. In most cases, the “Data not used” row can be filled in with “N/A”; it is primarily useful for specifying particular datasets that the Working Group considered but did not use in an assessment because they were of low quality and should not be used to inform fisheries management decisions.

Changes to Model Assumptions and Structure

17. The primary purpose of this section is to briefly identify only the most significant model changes that directly resulted in significant changes to results on the status of the stock concerned, and to briefly indicate the main effect of these changes. Details on model changes should be left in the main text of the report.

Qualifying Comments

18. The purpose of the “Qualifying Comments” section is to provide for any necessary explanations to avoid misinterpretation of information presented in the sections above. This section may also be used for brief further explanation considered important to understanding the status of the stock.

Fishery Interactions

19. The “Fishery Interactions” section should be used to simply list QMS by-catch species, non-QMS by-catch species and protected / endangered species interactions.


FOR FURTHER INFORMATION

IPCC (2007) Climate Change 2007: Synthesis Report. Contribution of Working Groups I, II and III to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change. [Core Writing Team, Pachauri, R K; Reisinger, A (eds.)]. IPCC, Geneva, Switzerland, 104 p.

New Zealand Ministry of Fisheries (2008) Harvest Strategy Standard for New Zealand fisheries. 25 p. Available at <http://fs.fish.govt.nz/Page.aspx?pk=61&tk=208&se=&sd=Asc&filSC=&filAny=False&filSrc=False&filLoaded=False&filDCG=9&filDC=0&filST=&filYr=0&filAutoRun=1>.

New Zealand Ministry of Fisheries (2011a) Research and Science Information Standard for New Zealand Fisheries. 31 p. Available at <http://www.fish.govt.nz/en-nz/Publications/Research+and+Science+Information+Standard.htm>.

New Zealand Ministry of Fisheries (2011b) Operational Guidelines for New Zealand’s Harvest Strategy Standard Revision 1. 78 p. Available at http://fs.fish.govt.nz/Doc/22847/Operational_Guidelines_for_HSS_rev_1_Jun_2011.pdf.ashx.



Fisheries New Zealand

Tini a Tangaroa

FNZ management teams and primary species managed

New Zealand Government

FISHERIES MANAGEMENT - INSHORE

Common name	Code	Stock
Anchovy	ANC	All
Barracouta	BAR	BAR1
Bladder kelp	KBB	All
Blue cod	BCO	All
Blue moki	MOK	All
Blue warehou	WAR	All
Bluenose	BNS	All
Butterfish	BUT	All
Cockle	COC	All
Deepwater (king) clam	PZL	All
Dredge oyster	OYS, OYU	All
Elephantfish	ELE	All
English mackerel	EMA	EMA1, 2
Flatfish	FLA	All
Freshwater eels (NI and SI)	ANG, LFE, SFE	All
Frostfish	FRO	FRO1, 2
Garfish	GAR	All
Gemfish	SKI	SKI1, 2
Ghost shark, dark	GSH	GSH1-3, 7-9
Greenlipped mussel	GLM	All
Grey mullet	GMU	All
Gurnard	GUR	All
Hapuka / bass	HPB	All
Horse mussel	HOR	All
Jack mackerel	JMA	JMA1
John dory	JDO	All
Kahawai	KAH	All
Kina	SUR	All
Kingfish	KIN	All
Knobbed whelk	KWH	All

FISHERIES MANAGEMENT - DEEPWATER

Common name	Code	Stock
Alfonsino	BYX	All
Barracouta	BAR	BAR4, 5, 7
Cardinalfish	CDL	All
Deepwater crabs (red crab, king crab, giant spider crab)	CHC, KIC, GSC	All
English mackerel	EMA	EMA3, 7
Frostfish	FRO	FRO3-9
Gemfish	SKI	SKI3, 7
Ghost shark, dark	GSH	GSH4-6
Ghost shark, pale	GSP	All
Hake	HAK	All
Hoki	HOK	All
Jack mackerel	JMA	JMA3, 7
Ling	LIN	LIN3-7
Lookdown dory	LDO	All
Orange roughy	ORH	All
Oreos	SSO, BOE, SOR, WOE, OEO	All
Patagonian toothfish	PTO	All
Prawnkiller	PRK	All
Redbait	RBT	All
Ribaldo	RIB	RIB3-8
Rubyfish	RBV	All
Scampi	SCI	All
Sea perch	SPE	SPE3-7
Silver warehou	SWA	All
Southern blue whiting	SBW	All
Spiny dogfish	SPD	SPD4, 5
Squid	SQU	All
White warehou	WWA	All

FISHERIES MANAGEMENT - HMS


Common name	Code	Stock
Albacore tuna *	ALB	All
Bigeye tuna	BIG	All
Blue shark	BWS	All
Mako shark	MAK	All
Moonfish	MOO	All
Pacific bluefin tuna	TOR	All
Portbeagle shark	POS	All
Ray's bream	RBM	All
Skipjack tuna *	SKJ	All
Southern bluefin tuna	STN	All
Swordfish	SWO	All
Yellowfin tuna	YFN	All

* non-QMS species

INTL POLICY - FISHERIES MGMT

Common name	RFMO
Antarctic toothfish,	CCAMLR
Patagonian toothfish	SPRFMO
Orange roughy	WCPFC
Pacific HMS species *	CCSBT
Southern bluefin tuna	
Regional Fisheries Management Organisations (RFMOs)	
CCAMLR - Commission for the Conservation of Antarctic Marine Living Resources	
SPRFMO - South Pacific Regional Fisheries Management Organisation	
WCPFC - Western and Central Pacific Fisheries Commission	
CCSBT - Commission for the Conservation of Southern Bluefin Tuna	

* primarily ALB, BIG, SKJ, SWO and YFN

ALFONSINO (BYX)*(Beryx splendens, B. decadactylus)***1. FISHERY SUMMARY**

Alfonsino was introduced into the Quota Management System (QMS) on 1 October 1986. Current allowances, TACCs and TACs are shown in Table 1.

Table 1: Recreational and Customary non-commercial allowances, TACCs and TACs for alfonsino by Fishstock for 2016–17.

Fishstock	Recreational Allowance	Customary non-commercial allowance	TACC	TAC
BYX 1	2	-	300	304
BYX 2	-	-	1 575	1 575
BYX 3	-	-	1 010	1 010
BYX 7	-	-	80.5	80.5
BYX 8	-	-	20	20
BYX 10	-	-	10	10

1.1 Commercial fisheries

Alfonsino has supported a major mid-water target trawl fishery off the east coast of the North Island since 1983 and is a minor bycatch of other trawl fisheries around New Zealand. The original gazetted TACs were based on the 1983–84 landings except for BYX 10 which was administratively set. Recent reported domestic landings and actual TACCs are shown in Table 2, while Figure 1 shows the historical landings and TACC values for the main BYX stocks.

Alfonsino landings in New Zealand consist almost entirely of one species, *Beryx splendens*: the other species, *B. decadactylus*, is thought to make up less than 1% of landings. Before 1983 alfonsino were virtually unfished, but two main fisheries now exist in New Zealand. The first to develop was the lower east coast North Island fishery (BYX 2), which developed in the mid-1980s. The other is the eastern Chatham Rise fishery (BYX 3), which developed in the mid-1990s. Alfonsino are caught throughout the New Zealand EEZ but only in small quantities outside of the east coast North Island and eastern Chatham Rise fisheries.

In BYX 1, alfonsino is mainly caught as a target species by bottom trawl within QMA 1. A smaller amount is taken as bycatch by bottom longline in the bluenose target fishery. The TACC for BYX 1 was increased for the 2001–02 fishing year from 31 t to 300 t when it was included in the adaptive management programme, and allocated 2 t for both customary and other mortality increasing the TAC to a total of 304 t. The new TACC was attained for the first time in 2004–05 and has been

ALFONSINO (BYX)

undercaught since then.

BYX 2 has historically been the major alfonsino fishery in the New Zealand EEZ. Prior to 1983, alfonsino was virtually an unfished resource. The domestic BYX 2 target fishery was developed during 1981, and was concentrated on the banks and seamount features off the east coast of the North Island, between Gisborne and Cape Palliser. Major fishing grounds included the Palliser Bank, Tuaheni Rise, Ritchie Banks and Paoanui Ridge. In more recent years, the alfonsino catch and effort has decreased from these areas, and an increasing proportion of the annual catch has been taken from the Madden Banks and Motukura Bank.

In BYX 3 catches of alfonsino were low in the early 1990s and were mainly bycatch of the hoki fishery. The TACC for BYX 3 was increased for the 1987–88 fishing year from 220 t to 1 000 t but annual landings remained low until 1993–94. However, the discovery of new grounds in the mid-1990s saw the rapid development of a target alfonsino fishery, most notably south-east of the Chatham Islands in Statistical Area 051. Annual landings are usually close to 1 000 t. The vast majority of the BYX 3 alfonsino catch is targeted now, followed by bycatch in fisheries for orange roughy, bluenose, hoki and hake. Catches are made all year round but decrease during the winter months. Catches of alfonsino in the Southland and Sub-Antarctic regions of BYX 3 are negligible.

Catches of alfonsino in BYX 7 are small. They are mainly taken by vessels midwater trawling for spawning hoki in Statistical Areas 034 and 035 in winter. There is essentially no targeting of alfonsino in BYX 7. The TACC was increased from 30 t to 80 t in 1989 but the TACC has never been caught. Annual landings are usually less than 30 t.

Landings have been reported from BYX 8 in only a few years. No targeting has ever been reported from this area. All catch has been from midwater trawls targeting jack mackerel and bottom longline targeting bluenose.

Catches of alfonsino from BYX 10 (Kermadec Region) are negligible. Apart from 1 t in 1989, and less than 1 t in each of 1992 and 1993, there have been no reported landings of alfonsino from this area.

Table 2: Reported domestic landings (t) of alfonsino by Fishstock from 1985–86 to 2016–17 and actual TACCs (t) from 1986–87 to 2016–17. QMS data from 1986–present. [Continued on next page].

Fishstock FMA (s)	BYX 1		BYX 2		BYX 3		BYX 7	
	1 & 9		2		3, 4, 5 & 6		7	
	Landings	TACC	Landings	TACC	Landings	TACC	Landings	TACC
1985–86*	11	-	1 454	-	3	-	1	-
1986–87	3	10	1 387	1 510	75	220	4	30
1987–88	8	27	1 252	1 511	101	1 000	2	30
1988–89	6	27	1 588	1 630	64	1 000	4	30
1989–90	24	31	1 496	1 274	147	1 007	21	80
1990–91	17	31	1 459	1 274	202	1 007	26	81
1991–92	7	31	1 368	1 499	264	1 007	2	81
1992–93	6	31	1 649	1 504	113	1 007	12	81
1993–94	7	31	1 688	1 569	275	1 007	31	81
1994–95	11	31	1 670	1 569	482	1 010	59	81
1995–96	11	31	1 868	1 569	961	1 010	66	81
1996–97	39	31	1 854	1 575	983	1 010	77	81
1997–98	14	31	1 652	1 575	1 164	1 010	67	81
1998–99	37	31	1 658	1 575	912	1 010	13	81
1999–00	25	31	1 856	1 575	743	1 010	24	81
2000–01	25	31	1 665	1 575	890	1 010	21	81
2001–02	123	300	1 574	1 575	1 197	1 010	10	81
2002–03	136	300	1 665	1 575	1 118	1 010	7	81
2003–04	219	300	1 468	1 575	884	1 010	11	81
2004–05	300	300	1 669	1 575	1 067	1 010	14	81

Table 2 [Continued]

Fishstock FMA (s)	BYX 1		BYX 2		BYX 3		BYX 7	
	1 & 9		2		3, 4, 5 & 6		7	
	Landings	TACC	Landings	TACC	Landings	TACC	Landings	TACC
2005–06	195	300	1 633	1 575	1 068	1 010	7	81
2006–07	66	300	1 644	1 575	945	1 010	21	81
2007–08	154	300	1 532	1 575	1 030	1 010	32	81
2008–09	172	300	1 589	1 575	895	1 010	18	81
2009–10	185	300	1 643	1 575	1 016	1 010	21	81
2010–11	48	300	1 686	1 575	1 084	1 010	17	81
2011–12	45	300	1 603	1 575	1 037	1 010	14	81
2012–13	22	300	1 605	1 575	1 013	1 010	39	81
2013–14	29	300	1 551	1 575	930	1 010	58	81
2014–15	53	300	1 617	1 575	997	1 010	26	81
2015–16	24	300	1 573	1 575	1 104	1 010	27	81
2016–17	22	300	1 611	1 575	991	1 010	29	81

Fishstock FMA (s)	BYX 10			
	10		Total	
	Landings	TACC	Landings	TACC
1985–86*	0	-	1 469	-
1986–87	0	10	1 470	1 800
1987–88	0	10	1 364	2 598
1988–89	1	10	1 663	2 717
1989–90	0	10	1 688	2 422
1990–91	0	10	1 664	2 423
1991–92	< 1	10	1 641‡	2 648
1992–93	< 1	10	1 780‡	2 653
1993–94	0	10	2 001‡	2 718
1994–95	0	10	2 223‡	2 721
1995–96	0	10	2 906‡	2 721
1996–97	0	10	2 953‡	2 727
1997–98	0	10	2 898‡	2 727
1998–99	0	10	2 624‡	2 727
1999–00	0	10	2 648‡	2 727
2000–01	0	10	2 601‡	2 727
2001–02	0	10	2 904‡	2 925
2002–03	0	10	2 927‡	2 925
2003–04	0	10	2 584‡	2 925
2004–05	0	10	3 052‡	2 925
2005–06	0	10	2 903‡	2 925
2006–07	0	10	2 677‡	2 925
2007–08	0	10	2 748‡	3 000
2008–09	0	10	2 674‡	3 000
2009–10	0	10	2 865‡	3 000
2010–11	0	10	2 836‡	2 996
2011–12	0	10	2 699‡	2 996
2012–13	0	10	2 679‡	2 996
2013–14	0	10	2 568‡	2 996
2014–15	0	10	2 693‡	2 996
2015–16	0	10	2 729‡	2 996
2016–17	0	10	2 653‡	2 996

*FSU data.

‡ Excludes catches taken outside the New Zealand EEZ.

ALFONSINO (BYX)




Figure 1: Reported commercial landings and TACC for the four main BYX stocks. Above: BYX 1 (Auckland) BYX 2 (Central East), BYX 3 (South East Coast, South East Chatham Rise, Sub Antarctic, Southland) Note that these figures do not show data prior to entry into the QMS. [Continued on next page].




Figure 1 [Continued]: Reported commercial landings and TACC for the four main BYX stocks, BYX 7 (Challenger). Note that these figures do not show data prior to entry into the QMS.

1.2 Recreational fisheries

Occasional catches of alfonsino have been recorded from recreational fishers.

1.3 Customary non-commercial fisheries

No quantitative information on the level of customary non-commercial catch is available.

1.4 Illegal catch

No quantitative information on the level of illegal alfonsino catch is available.

1.5 Other sources of mortality

No qualitative information is available.

2. BIOLOGY

In New Zealand waters, most “alfonsino” landings are alfonsino *B. Splendens*, with landings of the red bream *B. decadactylus* accounting for less than 1% of the catch. These species are primarily associated with undersea structures such as the seamounts that occur off the east coast of the North Island and on the Chatham Rise, in depths from 300–600 m. They can be found all around New Zealand waters but occur in greatest numbers along the lower east coast North Island and south-east Chatham Rise. These two areas are essentially where the commercial fisheries for alfonsino in New Zealand are confined.

Alfonsino are widespread in tropical, subtropical and temperate waters from the Atlantic, Pacific, and Indian Oceans (Busakhin 1982). They have been recorded in depths ranging from 10–1200 m but are most commonly found at 200–800 m, on or close to the seabed, often in association with seamounts and other underwater features (Maul 1981, Vinnichenko 1997a, Vinnichenko 1997b).

Stock structure is not currently known for New Zealand alfonsino. Horn & Massey (1989) found substantial differences in length frequency distributions between commercially-caught alfonsino from the Palliser bank compared with those from other locations on the east coast North Island. These differences suggest that there may be some age-specific migration occurring.

It has been suggested that alfonsino could comprise widespread populations in large oceanic eddy systems (Alekseev et al 1986). If New Zealand alfonsino form part of such a system then the east coast North Island may be a vegetative, non-reproductive zone where fish grow and mature before leaving for a possible reproductive zone further east of the mainland (Horn & Massey 1989).

ALFONSINO (BYX)

Alfonsino from Japan, northwest of Hawaii, and in the northeast of the Atlantic are known to spawn from August to October (Masuzawa et al 1975, Uchida & Uchihama 1986). In the southeast Atlantic, alfonsino spawn from January to March (Alekseev et al 1986) and from November to February in New Caledonian waters (Lehoday & Grandperrin 1994, Lehoday et al 1997). In New Zealand waters it has been suggested that alfonsino spawn from July to August (Horn & Massey 1989). This was based on observations of fish caught commercially from the lower east coast North Island that were ripening to spawn. However it is not known when and where spawning of alfonsino occurs in New Zealand waters. No running ripe fish were observed in regular samples taken over a 14-month period off the lower Wairarapa coast (Horn & Massey 1989).

Masuzawa et al (1975) estimated that the fecundity of a 40 cm female alfonsino from Japan to be 300 000–500 000 eggs. The fecundity of New Zealand alfonsino however has not been established because a full size range of ripening fish has not been observed (Horn & Massey 1989). Because of this the size and age at maturity cannot be determined precisely for either sex.

Tagging has been unsuccessful for alfonsino (Horn 1989). Being a moderately deepwater fish means that bringing them to the surface is not a viable option due to sudden and usually fatal changes in temperature, light, and particularly pressure. Horn (1989) evaluated the use of detachable hook tags using drop lines to tag alfonsino without bringing them to the surface. Only a small proportion of alfonsino tags were returned by commercial fishermen. This was thought to be due to a combination of low numbers being tagged to begin with (the tagging programme essentially targeted bluenose), low recapture rates, the loss of tags (either before or during capture by commercial fishermen), and possibly low rates of observation by fishermen.

Massey & Horn (1990) examined otoliths from commercially caught alfonsino from various alfonsino fishing grounds of the lower east coast of the North Island (BYX 2) from November 1985 to December 1986. They found evidence that one opaque and one hyaline zone (one 'ring') were formed annually (as did Lehodey & Grandperrin (1996)). They investigated the validity of zone counts by measuring the position of each ring and comparing it to the position of successive ring groups. They calculated the 'marginal index' of each otolith which was defined as the distance from the outer edge of the last hyaline ring to the otolith edge divided by the width of the last complete opaque and hyaline ring. They plotted the mean marginal indices of fish for each month over the study period and found that the index in every fishing ground dropped dramatically from June to December. This drop in mean marginal index meant that for most fish opaque material has started forming in June, and that the hyaline margin is probably laid down from March to May for most fish. Subsequent ageing has also shown the progression of relatively strong year classes between consecutive years of sampling, thus providing further support for the ageing method.

Massey & Horn (1990) observed very few fish younger than three years of age, and believed that full recruitment to the commercial fishery probably occurs at around five years of age. Size-at-sexual maturity is probably about 30 cm fork length (FL) at 4 to 5 years of age. Juvenile fish have been recorded in the pelagic and epipelagic zones in the North Pacific and Indian Oceans. Alfonsino less than 20 cm FL are seldom recorded in New Zealand waters. Differences in length-frequency distributions between fishing grounds off the east coast North Island suggest that some age-specific migration occurs. Fish probably recruit to these grounds at 28–31 cm FL.

Von Bertalanffy growth parameters were derived for alfonsino from BYX 2 by Stocker & Blackwell (1991) (Table 3). They found that females attain a larger size than males and are also larger at corresponding ages. Massey & Horn (1990) presented von Bertalanffy parameters separately by sex for three fishing grounds off lower east coast North Island.

Stocker & Blackwell (1991) used the equation $M = \log_e 100/\text{maximum age}$, where maximum age is the age to which 1% of the population survives in an unexploited stock. Using a maximum age of 20 years, they estimated M for both sexes as 0.23 for BYX 2.

Length-weight relationships are presented in Table 3. Parameters for the Chatham Rise are those reported by O'Driscoll et al (2011) for all fish from the summer Chatham Rise trawl survey time series from 1992–2010.

Horn et al (2010) examined stomach contents from *Beryx splendens* caught on three consecutive summer trawl surveys of the Chatham Rise (2005–2007). They found that alfonsino were moderately selective feeders that fed primarily in the mesopelagic layers. The most common prey items were crustaceans and mesopelagic fishes. By mass, the most important were prawns from the genus *Sergestes*, followed by the myctophid fish *Lampanyctodes hectoris*, and then prawns from the genus *Pasiphaea*.

Smaller crustaceans such as euphysiids and amphipods are most important in the diet of smaller alfonsino (17–26.5 cm fork length). Larger prawn species and mesopelagic fishes were more important for larger alfonsino (27–42 cm fork length). Horn et al (2010) postulated that they are selective feeders based on the observation that prey items such as squid and salps would be relatively abundant where alfonsino feed on the Chatham Rise, but are rarely taken.

Table 3: Estimates of biological parameters for alfonsino.

Fishstock			Estimate	Source
<u>1. Natural mortality (<i>M</i>)</u>				
BYX 2			0.23	Stocker & Blackwell (1991)
<u>2. Weight = a(length)^b (Weight in g, length in cm fork length).</u>				
			<u>Both Sexes</u>	
			a	b
BYX 2			0.0226	3.018
BYX 3			0.019	3.049
Stocker & Blackwell (1991)				
O'Driscoll et al (2011)				
<u>3. Von Bertalanffy growth parameters</u>				
			<u>Females</u>	
			<u><i>L</i>_∞</u>	<u><i>k</i></u>
			<u><i>t</i>₀</u>	
BYX 2			57.5	0.08
			-4.10	
			<u>Males</u>	
			<u><i>L</i>_∞</u>	<u><i>k</i></u>
			<u><i>t</i>₀</u>	
BYX 2			51.1	0.11
			-3.56	
Stocker & Blackwell (1991)				

3. STOCKS AND AREAS

No information is available as to whether alfonsino is a single stock in New Zealand waters. Overseas data on alfonsino stock distributions suggest that New Zealand fish could form part of a widely distributed South Pacific stock.

4. STOCK ASSESSMENT

4.1 Estimates of fishery parameters and abundance

i) BYX 1

Starr et al (2010) presented CPUE analyses from the bycatch of alfonsino in the east Northland and Bay of Plenty target longline fisheries for bluenose and hapuku. The two series showed no sign of decline up to 2007–08, but the indices were based on only 12% of the BYX catch from the area. The analyses have not been updated, and the catch of BYX has decreased to below 50 t for the last five years.

ii) BYX 2

A biomass index derived from a standardised CPUE (log linear, kg/day) analysis of the target trawl fishery represented by seven core vessels (Blackwell 2000) was calculated for BYX 2. However, the analysis was very uncertain, and the model accounted for only 25% of the variance in catch rates. The results of the standardised analysis were not accepted by the Inshore WG as indices of abundance.

The age composition of the commercial landings in BYX 2 was determined in 1998–99, 1999–00, and 2000–01 and 2002–03, 2003–04 and 2004–05. The commercial catch is dominated by 5–11 year

old fish. Without linking age structure to specific fishing grounds the age structure of the catch is unlikely to monitor changes in the population.

iii) BYX 3

The potential to monitor trends in abundance using catch and effort data from the target BYX 3 fishery was investigated by Langley & Walker (2002b). However, it was concluded that the high variation in catch rates, the relatively small number of catch and effort records, and the complex nature of the fishery precluded the development of a reliable CPUE index.

4.2 Biomass estimates

Estimates of current biomass are not available.

4.3 Yield estimates and projections

4.3.1 Other yield estimates and stock assessment factors

Long-term sustainable yield using an $F_{0.1}$ fishing strategy was estimated for BYX 2 using the simulation model with alternative estimates of M . $F_{0.1}$ has been estimated as 0.25 and 0.32 for $M = 0.2$ and $M = 0.23$, respectively, for both sexes combined in BYX 2 (Stocker & Blackwell 1991). The biomass at this long-term equilibrium yield is about 35% B_0 and the $F_{0.1}$ yield is about 8–9% B_0 .

4.4 Other factors

The most recent assessment for BYX 2 was based upon the historical fishery areas. In recent years the fishery has expanded to new areas not previously fished. Subsequent CPUE analyses have been rejected by Working Groups and it is no longer thought possible to monitor abundance in BYX 2 using trawl CPUE.

Current data on alfonsino movements are inconclusive. It is not known whether the fish on the east coast of the North Island spend some part of their life cycle in other New Zealand waters, or whether the east coast-Chatham Rise region is just one of several pre-reproductive regions. It is possible that the domestic trawl fishery may be exploiting part of a wider South Pacific stock. Catches may be maintained due to the discovery of new grounds. However, the potential for increased catches may be constrained by the availability of BNS 3 quota to cover likely bluenose bycatch.

5. STATUS OF THE STOCKS

Stock Structure Assumptions

No information is available as to whether alfonsino is a single stock in New Zealand fishery waters. Overseas data on alfonsino stock distributions suggest that New Zealand fish could form part of a widely distributed South Pacific stock. In addition to alfonsino (*Beryx splendens*) the BYX Fishstock includes landings of the red bream (*B. decadactylus*), however, red bream makes up less than 1% of the total landings.

BYX 1

Under the adaptive management programme the TACC was increased to 300 t in 2001–02, and catches increased for the next 9 years in the target trawl fishery. However, catches have been below 50 t since 2010–11 as target fishing in this fishery has waned.

BYX 2

Annual landings from 1986 to 2014–15 have remained reasonably stable at or above the level of the TACC. However, as the fishing grounds have extended throughout this time, it is not known if the recent catch levels or the current TACCs are sustainable.

BYX 3

Alfonsino on the Chatham Rise (BYX 3) were lightly fished prior to 1995–96 when catches increased to near the TACC, due to the development of new fishing grounds. Catch has fluctuated around the

TACC since then. It is not known if the recent catch levels or the current TACCs are sustainable.

Table 4: Summary of TACCs (t) and reported landings (t) for alfonsino for the most recent fishing year.

Fishstock	FMAs	2016–17	2016–17
		Actual TACC	Reported landings
BYX 1 Auckland (East) (West)	1 & 9	300	22
BYX 2 Central (East)	2	1 575	1 611
BYX 3 South-East (Coast)	3, 4, 5,	1 010	991
Southland & Sub-Antarctic	& 6		
BYX 7 Challenger	7	81	29
BYX 8 Central (West)	8	20	<1
BYX 10 Kermadec	10	10	0
Total		2 996	2 653

6. FUTURE RESEARCH CONSIDERATIONS

Neither CPUE nor trawl surveys are likely to provide an index of alfonsino abundance. The best method to determine the status of the stocks and to continue monitoring is likely to be a catch-at-age sampling programme. A large proportion of the alfonsino catch from the two main fisheries is still landed green which would allow for a land-based shed sampling programme for either area, although at-sea observer-based sampling would allow for the detection of any differences in sub-regions within the main fishery areas.

7. FOR FURTHER INFORMATION


- Alekseev, F E; Alekseeva, E I; Trunov, I A; Shlibanov, V I (1986) Macroscale water circulation, ontogenetic geographical differentiation and population structure of alfonsino, *Beryx splendens* Lowe, in the Atlantic Ocean. *International Council for the Exploration of the Sea [ICES]* No 10. 16 p.
- Blackwell, R (2000) Alfonsino (*Beryx splendens*) abundance indices from standardised catch per unit effort (CPUE) analysis for the east coast North Island (BYX 2) midwater trawl fishery 1989–90 to 1997–98. *New Zealand Fisheries Assessment Report 2000/53*. 40 p.
- Busakhin, S V (1982) Systematics and distribution of the family Berycidae (Osteichthyes) in the World Ocean. *Journal of Ichthyology* (22): 1–21.
- Horn, P L (1988) Alfonsino. New Zealand Fisheries Assessment Research Document 1988/7. 21 p. (Unpublished document held by NIWA library, Wellington.)
- Horn, P L (1989) An evaluation of the technique of tagging alfonsino and bluenose with detachable hook tags. *N.Z. Fisheries Technical Report No. 16*. 15 p.
- Horn, P L; Forman, J; Dunn, M R (2010) Feeding habits of alfonsino *Beryx splendens*. *Journal of Fish Biology* 76: 2382–2400.
- Horn, P L; Massey, B R (1989) Biology and abundance of alfonsino and bluenose off the lower east coast, North Island, New Zealand. *New Zealand Fisheries Technical Report No. 15*. 32 p.
- Langley, A D (1995) Analysis of commercial catch and effort data from the QMA 2 alfonsino-bluenose trawl fishery 1989–94. New Zealand Fisheries Assessment Research Document 1995/18. 12 p. (Unpublished document held by NIWA library, Wellington.)
- Langley, A D; Walker, N (2002a) Characterisation of the alfonsino (*Beryx splendens*) fishery in BYX 3. *New Zealand Fisheries Assessment Report 2002/29*. 49 p.
- Langley, A D; Walker, N (2002b) CPUE analysis of the target BYX 3 alfonsino fishery and associated bluenose catch. *New Zealand Fisheries Assessment Report 2002/24*. 45 p.
- Lehodey, P; Grandperrin, R (1994) A study of the fishery and biology of *Beryx splendens* (alfonsino) in New Caledonia. *SPC Fisheries Newsletter* 71: 30–36.
- Lehodey, P; Grandperrin, R; Marchal, P (1997) Reproductive biology and ecology of a deep demersal fish, alfonsino *Beryx splendens*, over the seamounts off New Caledonia. *Marine Biology* 128(1): 17–27.
- MacGibbon, D G (2015) Fishery characterisation and standardised CPUE analyses for alfonsino, *Beryx splendens*, 1989–90 to 2013–14. *New Zealand Fisheries Assessment Report 2015/78*. 238 p.
- Massey, B R; Horn, P L (1990) Growth and age structure of alfonsino (*Beryx splendens*) from the lower east coast, North Island, New Zealand. *New Zealand Journal of Marine and Freshwater Research* 24: 121–136.
- Masuzawa, T; Kurata, Y; Onishi, K (1975) Results of group study on a population of demersal fishes in waters from Sagami Bay to Southern Izu Islands: population ecology of Japanese alfonsino and other demersal fishes. *Japan Aquatic Resources Conservation Association Fishery Research Paper No 28*. 105 p.
- Maul, G E (1981) Berycidae. In: Fiches FAO d'Identification des espèces pour les besoins de la pêche. Atlantique centre-est. Fonds D'épôt-FAO. Ottawa, Canada.
- Northern Inshore Fisheries Company Ltd. (2001) Proposal to increase the TACC for BYX 1 -- final 30/04/01.
- O'Driscoll, R L; MacGibbon, D; Fu, D; Lyon, W; Stevens, D W (2011) A review of hoki and middle-depth trawl surveys of the Chatham Rise, January 1992–2010. *New Zealand Fisheries Assessment Report 2011/47*. 814 p.
- Seafood Industry Council (SeaFIC) (2003) BYX 1 performance report to the 2003 Adaptive Management Programme Working Group.
- Seafood Industry Council (SeaFIC) (2004) Report to the Adaptive Management Fishery Assessment Working Group: Performance of the BYX 1 Adaptive Management Programme. AMP-WG-2004/06. (Unpublished report held by FNZ.)
- Seafood Industry Council (SeaFIC) (2005) Report to the Adaptive Management Fishery Assessment Working Group: Performance of the BYX 1 Logbook Programme. AMP-WG-2005/05. (Unpublished report held by FNZ.)

ALFONSINO (BYX)

- Starr, P J; Kendrick, T H; Bentley, N (2010) Report to the Adaptive Management Programme Fishery Assessment Working Group: Characterisation, CPUE analysis and logbook data for BYX 1. Document 2010/04-v2, 86 p. (Unpublished report held byFNZ) (<http://cs.fish.govt.nz/forums/thread/3874.aspx>)
- Starr, P J; Kendrick, T H; Lydon, G J (2006) 2006 Report to the Adaptive Management Programme Fishery Assessment Working Group: Full Term Review of the BNS 1 Adaptive Management Programme. AMP-WG-2006/03. 59 p. (Unpublished report available from Seafood New Zealand, Wellington.)
- Stocker, M; Blackwell, R (1991) Biomass and yield estimates for alfonsino in BYX 2 for the 1991–92 fishing year. New Zealand Fisheries Assessment Research Document 1991/12. 12 p. (Unpublished document held in NIWA library, Wellington.)
- Uchida, R N; Uchihama, J H (1986) Fishery atlas of the northwestern Hawaiian Islands. *NOAA Technical Report NAPS No 38*. 142 p.
- Vinnichenko, V I (1997a) Russian investigations and deep water fishery on the Corner Rising Seamount in Subarea 6. *NAFO Scientific Council Studies* (30): 41–49.
- Vinnichenko, V I (1997b) Vertical daily migrations of the slender alfonsino *Beryx splendens* (Berycidae) at the underwater rises of the open North Atlantic. *Journal of Ichthyology* (37): 438–444.

ANCHOVY (ANC)

(*Engraulis australis*)
Kokowhaawhaa



1. FISHERY SUMMARY

Anchovy were introduced into the QMS on 1 October 2002, with allowances, TACCs and TACs in Table 1. These have not changed.

Table 1: Recreational and Customary non-commercial allowances, TACCs and TACs for anchovy by Fishstock.

Fishstock	Recreational Allowance	Customary non-commercial allowance	TACC	TAC
ANC 1	10	5	200	215
ANC 2	10	5	100	115
ANC 3	2	1	50	53
ANC 4	3	2	10	15
ANC 7	10	5	100	115
ANC 8	10	5	100	115
ANC 10	0	0	0	0

1.1 Commercial fisheries

There is no information on catches or landings of anchovy prior to 1990, although sporadic catches were made in some years during exploratory fishing projects for small pelagic species, in the 1960s and 1970s. It is thought that anchovy were caught in most years, but were either not reported, reported as “bait”, or included in the category “mixed species”. Reported annual landings have fluctuated from less than 1 t to 21 t since 1990–91 (Table 2). Under-reporting is likely to have occurred due to misidentification of anchovy in pilchard and other mixed catches and the low value of the species.

Historically most landings have been reported from northeastern New Zealand, ANC 1, with occasional small landings in ANC 3 and 8.

The most consistent (though small) catches have been taken by purse seine. Very few catches have been reported as targeted; most anchovy appear to have been taken as non-target catch in the pilchard fishery. Up to four vessels reported a catch or landing in any one year.

ANCHOVY (ANC)

Table 2: Reported catches or landings (t) of anchovy by fishstock from 1990–91 to 2016–17 (prior to 2002–03 reported by FMA). MHR data from 2001–02 - present.

Fishstock FMA	ANC 1 1	ANC 2 2	ANC 3 3,5&6	ANC 4 4	ANC 7 7	ANC 8 8&9	ANC 10 10	Total
1990–91†	< 1	0	0	0	< 1	0	0	< 1
1991–92†	1	0	1	0	< 1	0	0	2
1992–93†	21	0	0	0	0	0	0	21
1993–94†	< 1	0	0	0	0	0	0	< 1
1994–95†	< 1	0	0	0	< 1	0	0	< 1
1995–96†	1	0	0	0	0	0	0	1
1996–97†	2	0	0	0	0	0	0	2
1997–98†	1	0	0	0	0	0	0	1
1998–99†	4	0	2	0	0	0	0	6
1999–00†	3	0	0	0	0	0	0	3
2000–01†	10	0	0	0	0	0	0	10
2001–02	7	0	0	0	0	0	0	7
2002–03	8	0	0	0	0	0	0	8
2003–04	4	0	0	0	0	10	0	15
2004–05	< 1	0	0	0	0	12	0	12
2005–06	10	0	0	0	0	< 1	0	10
2006–07	< 1	0	0	0	0	2	0	3
2007–08	< 1	0	0	0	< 1	< 1	0	< 1
2008–09	< 1	0	0	0	< 1	< 1	0	2
2009–10	6	0	0	0	6	0	0	12
2010–11	1	0	< 1	0	< 1	< 1	0	1
2011–12	< 1	0	0	0	0	0	0	< 1
2012–13	0	0	< 1	0	< 1	< 1	0	< 1
2013–14	2	0	< 1	0	< 1	< 1	0	2
2014–15	1	0	< 1	0	0	< 1	0	< 1
2015–16	< 1	0	0	0	11	0	0	11
2016–17	< 1	0	0	0	5	0	0	5

† CELR

1.2 Recreational fisheries

There is no known recreational fishery, but small numbers are caught in small-mesh setnets and beach seines. An estimate of the recreational harvest is not available.

1.3 Customary non-commercial fisheries

An estimate of the customary non-commercial catch is not available.

1.4 Illegal catch

There is no known illegal catch of anchovies.

1.5 Other sources of mortality

Some accidental captures of anchovy by vessels purse seining for other small pelagic species may be discarded if no market is available.

2. BIOLOGY

The single anchovy species, *Engraulis australis*, found in New Zealand also occurs around much of the Australian coast. In New Zealand, it occurs around most of the coastline, but is absent between Banks Peninsula and Foveaux Strait. It is found mostly inshore, particularly in gulfs, bays, harbours, and some large estuaries. In Australia it tends to move seaward in winter, returning closer inshore during spring and the same pattern is likely to occur in New Zealand. Its vertical distribution in the water column is not known, but it seems likely that it occurs at all depths between the surface and the coastal seafloor.

Anchovy are planktivorous, feeding mainly on copepods. They form compact schools, particularly during the warmer months and larger fishes, seabirds, and marine mammals prey heavily upon these schools. Although they generally form single-species schools, anchovies are closely associated with other small pelagic fishes, particularly pilchard and sprats.

The reproductive cycle is not well known. The main spawning season appears to be spring-summer, but in northern regions spawning may occur through much of the year. Spawning grounds extend from shallow water out to mid-shelf. The eggs are pelagic.

No reliable ageing work has been undertaken in New Zealand, but some information is available for this species in Australia where it reaches 16 cm at age 6, and matures at age 1. In northeastern New Zealand, the main size range of anchovy is 8–14 cm, which are likely to be 2–5 year old fish.

There have been no biological studies that are directly relevant to the recognition of separate stocks, or to yield estimates. Consequently no estimates of biological parameters are available. There is extensive international literature on similar species of anchovy, but the relevance of this to the New Zealand species is unknown.

3. STOCKS AND AREAS

No biological information is available on which to make an assessment on whether separate anchovy stocks exist in New Zealand. If spawning is as widespread as the fragmentary accounts suggest and if there is limited migration between regions, there is potential for localised depletion.

Anchovy and pilchard are often caught together. Anchovy fishstock boundaries are fully aligned with those for pilchard.

4. STOCK ASSESSMENT

There have been no stock assessments of New Zealand anchovy.

4.1 Estimates of fishery parameters and abundance

No fishery parameters are available.

4.2 Biomass estimates

No estimates of biomass are available.

4.3 Yield estimates and projections

MCY cannot be determined.

Current biomass cannot be estimated, so *CAY* cannot be determined.

4.4 Other yield estimates and stock assessment results

No information is available.

4.5 Other factors

Ichthyoplankton surveys show anchovy to be locally abundant. However, it is unlikely that the biomass is comparable to the very large stocks of anchovy in some oceans where strong upwelling promotes high productivity. It is more likely that New Zealand anchovy comprise abundant but localised coastal populations.

It is not known whether the biomass of anchovy is stable or variable, but the latter is considered more likely.

In some localities anchovy are a major food source for many fish, seabirds, and marine mammals (e.g., a major component of fur seal diet in May–August at Cape Foulwind). Excessive localised harvesting may disrupt ecosystems.

5. STATUS OF THE STOCKS

No estimates of current biomass are available. At the present level of minimal catches, stocks should be at or close to their natural level. This is nominally a virgin biomass, but not necessarily a stable one. It is not yet possible to estimate a long-term sustainable yield for anchovy.

TACCs and reported landings for the 2016–17 fishing year are summarised in Table 3.

Table 3: Summary of TACCs (t) and reported landings (t) of anchovy for the most recent fishing year.


Fishstock		FMA	2016–17 Actual TACC	2016–17 Reported landings
ANC 1	Auckland (East)	1	200	<1
ANC 2	Central (East)	2	100	0
ANC 3	South-east (Coast), Southland & sub-Antarctic	3, 5 & 6	50	0
ANC 4	South-east (Chatham)	4	10	0
ANC 7	Challenger	7	100	5
ANC 8	Central (West), Auckland (West)	8 & 9	100	0
ANC 10	Kermadec	10	0	0
Total			560	5

6. FOR FURTHER INFORMATION

- Cole, R G (1987) Distribution and abundance of clupeoid larvae in the Hauraki Gulf. MSc. thesis, Department of Zoology, University of Auckland.
- Crossland, J (1981) Fish eggs and larvae of the Hauraki Gulf, New Zealand. *Fisheries Research Division Bulletin* 23. 61 p.
- Crossland, J (1982) Distribution and abundance of fish eggs and larvae from the spring and summer plankton of north-east New Zealand, 1976–78. *Fisheries Research Division Bulletin* 24. 59 p.
- McKnee, A (1993) Anchovy. In: Kailola, P J et al (comps). 1993. Australian fisheries resources, pp 205–206. Bureau of Resource Sciences and the Fisheries Research and Development Corporation. Canberra. 422 p.
- Robertson, D A (1978) Blue mackerel, pilchard, anchovy, sprat, saury, and lanternfish. In: Habib, G; Roberts, P E (comps) Proceedings of the Pelagic Fisheries Conference July 1977. *Fisheries Research Division Occasional Publication No. 15*: 85–89
- Tricklebank, K A; Jacoby, C A; Montgomery, J C (1992) Composition, distribution and abundance of neustonic ichthyoplankton off northeastern New Zealand. *Estuarine, Coastal and Shelf Science* 34 (3): 263–275.

ARROW SQUID (SQU)

(*Nototodarus gouldi*, *N. sloanii*)
 Wheketere



1. FISHERY SUMMARY

1.1 Commercial fisheries

The New Zealand arrow squid fishery is based on two related species. *Nototodarus gouldi* is found around mainland New Zealand north of the Subtropical Convergence, whereas *N. sloanii* is found in and to the south of the convergence zone.

Except for the Southern Islands fishery, for which a separate TACC is set, the two species are managed as a single fishery within an overall TACC. The Southern Islands fishery (SQU 6T) is almost entirely a trawl fishery. Although the species (*N. sloanii*) is the same as that found around the south of the South Island, there is evidence to suggest that the Auckland Island shelf stock is different from the mainland stocks. Because the Auckland Island shelf squid are readily accessible to trawlers, and because they can be caught with little finfish bycatch and are therefore an attractive resource for trawlers, a quota has been set separately for the Southern Islands. Total reported landings and TACCs for each stock are shown in Table 1, while historical landings and TACC are depicted in Figure 1.

The New Zealand squid fishery began in the late 1970s and reached a peak in the early 1980s when over 200 squid jigging vessels came to fish in the New Zealand EEZ. The discovery and exploitation of the large squid stocks in the southwest Atlantic substantially increased the supply of squid to the Asian markets causing the price to fall. In the early 1980s, Japanese squid jiggers would fish in New Zealand for a short time before continuing on to the southwest Atlantic. In the late 1980s, the jiggers stopped transit fishing in New Zealand and the number of jiggers fishing declined from over 200 during the 1983–84 fishing year to around 15 in 1994–95. The jig catch in SQU 1J declined from a peak of 53 872 t in 1988–89 to under 1 000 t per year by 2012–13. In 2016–17 the TACC was reduced from 50 212 t to 5 000 t to reflect these changes within this fishery.

From 1987 to 1998 the trawl catch fluctuated between about 30 000–70 000 t, but in SQU 6T the impact of management measures to protect the Hooker's sea lion (*Phocarctos hookeri*) restricted the total catch in some years between 1999 and 2005.

Catch and effort data from the SQU 1T fishery show that the catch occurs between December and May, with peak harvest from January to April. The catch has been taken from the Snares shelf on the south coast of the South Island right through to the Mernoo Bank (east coast), but Statistical Area 028 (Snares shelf and Snares Island region) has accounted for over 77% of the total in recent years. Based on

ARROW SQUID (SQU)

Observer data, squid accounts for 67% of the total catch in the target trawl fishery, with bycatch principally of barracouta, jack mackerel, silver warehou and spiny dogfish.

For 2005–06 a 10% in-season increase to the SQU 1T TACC was approved by the Minister of Fisheries. The catch for December–March was 40% higher than the average over the previous eight years and catch rates were double the average, indicating an increased abundance of squid. Previously, in 2003–04, a 30% in-season increase to the TACC was agreed, but catches did not reach the higher limit. Note that the TACC automatically reverts to the original value at the end of the fishing year.

Table 1: Reported catches (t) and TACCs (t) of arrow squid from 1986–87 to 2016–17. Source - QMS.

Fishstock	SQU		SQU 1T*		SQU		SQU		Total	
	Landings	TACC	Landings	TACC	Landings	TACC	Landings	TACC	Landings	TACC
1986–87	32 394	57 705	25 621	30 962	16 025	32 333	0	10	74 040	121 010
1987–88	40 312	57 705	21 983	30 962	7 021	32 333	0	10	69 316	121 010
1988–89	53 872	62 996	26 825	36 081	33 462	35 933	0	10	114 160	135 080
1989–90	13 895	76 136	13 161	47 986	19 859	42 118	0	10	46 915	166 250
1990–91	11 562	46 087	18 680	42 284	10 658	30 190	0	10	40 900	118 571
1991–92	12 985	45 766	36 653	42 284	10 861	30 190	0	10	60 509	118 571
1992–93	4 865	49 891	30 862	42 615	1 551	30 369	0	10	37 278	122 875
1993–94	6 524	49 891	33 434	42 615	34 534	30 369	0	10	74 492	122 875
1994–95	33 615	49 891	35 017	42 741	30 683	30 369	0	10	99 315	123 011
1995–96	30 805	49 891	17 823	42 741	14 041	30 369	0	10	62 668	123 011
1996–97	20 792	50 212	24 769	42 741	19 843	30 369	0	10	65 403	123 332
1997–98	9 329	50 212	28 687	44 741	7 344	32 369	0	10	45 362	127 332
1998–99	3 240	50 212	23 362	44 741	950	32 369	0	10	27 553	127 332
1999–00	1457	50 212	13 049	44 741	6 241	32 369	0	10	20 747	127 332
2000–01	521	50 212	31 297	44 741	3 254	32 369	< 1	10	35 071	127 332
2001–02	799	50 212	35 872	44 741	11 502	32 369	0	10	48 173	127 332
2002–03	2 896	50 212	33 936	44 741	6 887	32 369	0	10	43 720	127 332
2003–04	2 267	50 212	48 060	#58 163	34 635	32 369	0	10	84 962	127 332
2004–05	8 981	50 212	49 780	44 741	27 314	32 369	0	10	86 075	127 332
2005–06	5 844	50 212	49 149	#49 215	17 425	32 369	0	10	72 418	127 332
2006–07	2 278	50 212	49 495	44 741	18 479	32 369	0	10	70 253	127 332
2007–08	1 371	50 212	36 171	44 741	18 493	32 369	0	10	56 035	127 332
2008–09	1 032	50 212	16 407	44 741	28 872	32 369	0	10	46 311	127 332
2009–10	891	50 212	16 759	44 741	14 786	32 369	0	10	32 436	127 332
2010–11	1 414	50 212	14 957	44 741	20 934	32 369	0	10	37 304	127 332
2011–12	1 811	50 212	18 969	44 741	14 427	32 369	0	10	35 207	127 332
2012–13	741	50 212	13 951	44 741	9 944	32 369	0	10	24 637	127 332
2013–14	167	50 212	7 483	44 741	7 403	32 369	0	10	15 053	127 332
2014–15	513	50 212	9 668	44 741	6 127	32 369	0	10	16 310	127 332
2015–16	937	50 212	17 018	44 741	25 172	32 369	<1	10	43 127	127 332
2016–17	1	5 000	7 735	44 741	10 726	32 369	0	10	18 462	82 120

* All areas except Southern Islands and Kermadec.

† Southern Islands.

‡ Kermadec.

In season increase of 30% for 2003–04 and 10% for 2005–06

1.2 Recreational fisheries

The amount of arrow squid caught by recreational fishers is not known.

1.3 Customary non-commercial fisheries

No quantitative information is available on the current level of customary non-commercial take.

1.4 Illegal catch

There is no quantitative information available on the level of illegal catch.

1.5 Other sources of mortality

No information is available on other sources of mortality.

2. BIOLOGY

Two species of arrow squid are caught in the New Zealand fishery. Both species are found over the continental shelf in water up to 500 m depth, though they are most prevalent in water less than 300 m depth. Both species are sexually dimorphic, though similar in biology and appearance. Individuals can be identified to species level based on sucker counts on Arm I and differences in the hectocotylized arm of males.




Figure 1: Reported commercial landings and TACC for the three main SQU stocks. Top to bottom: SQU 1J (All Waters Except 10T and 6T, Jigging), SQU 1T (All Waters Except 10T and 6T, All Other Methods), and SQU 6T (Southern Islands, All Methods). Note that these figures do not show data prior to entry into the QMS.

ARROW SQUID (SQU)

Recent work on the banding of statoliths from *N. sloanii* suggests that the animals live for around one year. Growth is rapid. Modal analysis of research data has shown increases of 3.0–4.5 cm per month for Gould's arrow squid measuring between 10 and 34 cm Dorsal Mantle Length (DML).

Estimated ages suggest that *N. sloanii* hatches in July and August, with spawning occurring in June and July. It also appears that *N. gouldi* may spawn one to two months before *N. sloanii*, although there are some indications that *N. sloanii* spawns at other times of the year. The squid taken by the fishery do not appear to have spawned.

Tagging experiments indicate that arrow squid can travel on average about 1.1 km per day with a range of 0.14–5.6 km per day.

Biological parameters relevant to stock assessment are shown in Table 2.

Table 2: Estimates of biological parameters.

Fishstock			Estimate	Source
<u>1. Weight = a (length)^b (Weight in g, length in cm dorsal length)</u>				
		a	b	
<i>N. gouldi</i>	≤ 12 cm DML	0.0738	2.63	Mattlin et al (1985)
<i>N. sloanii</i>	≥ 12 cm DML	0.029	3	
<u>2. von Bertalanffy growth parameters</u>				
	<i>K</i>	<i>t</i> ₀	<i>L</i> _∞	
<i>N. gouldi</i>	2.1–3.6	0	35	Gibson & Jones (1993)
<i>N. sloanii</i>	2.0–2.8	0	35	

3. STOCKS AND AREAS

There are no new data which would alter the stock boundaries given in previous assessment documents. It is assumed that the stock of *N. gouldi* (the northern species) is a single stock, and that *N. sloanii* around the mainland comprises a unit stock for management purposes, though the detailed structure of these stocks is not fully understood. The distribution of the two species is largely geographically separate but those occurring around the mainland are combined for management purposes. The Auckland Islands Shelf stock of *N. sloanii* appears to be different from the mainland stock and is managed separately.

4. ENVIRONMENTAL AND ECOSYSTEM CONSIDERATIONS

This section was last reviewed by the Aquatic Environment Working Group for the May 2016 Fishery Assessment Plenary and has been updated in 2018. This summary is from the perspective of the squid trawl fishery; a more detailed summary from an issue by issue perspective is available in the 2017 Aquatic Environment & Biodiversity Annual Review MPI 2017, (<https://www.mpi.govt.nz/dmsdocument/27471-aquatic-environment-and-biodiversity-annual-review-aebar-2017-a-summary-of-environmental-interactions-between-the-seafood-sector-and-the-aquatic-environment>).

4.1 Role in the ecosystem

Arrow squid are short-lived and highly variable between years (see Biology section). Hurst et al (2012) reviewed the literature and noted that arrow squid are an important part of the diet for many species. Stevens et al (2012) reported that, between 1960 and 2000, squids (including arrow squid) were important in the diet of banded stargazer (59% of non-empty stomachs), bluenose (26%), giant stargazer (34%), gemfish (43%), and hapuku (21%), and arrow squid were specifically recorded in the diets of alfonsino, barracouta, hake, hoki, ling, red cod, red gurnard, sea perch, and southern blue whiting. In a detailed study on the Chatham Rise (Dunn et al 2009), cephalopods were identified as prey of almost all demersal fish species, and arrow squid were identified in the diet of hake, hoki, ling, Ray's bream, shovelnose spiny dogfish, sea perch, smooth skate, giant stargazer and silver warehou, and was a significant component (over 10% prey weight) of the diet of barracouta and spiny dogfish.

Arrow squid have been recorded as important in the diet of marine mammals such as NZ fur seals and NZ sea lions, particularly during summer and autumn (Fea et al 1999, Harcourt et al. 2002, Chilvers 2008, Boren 2008) and in the diet of common dolphins (Meynier et al 2008, Stockin 2008). They are also important in the diet of seabirds such as shy albatross in Australia (Hedd & Gales 2001) and Buller's albatross at the Snares and Solander Islands (James & Stahl 2000). Cephalopods in general are important in the diet of a wide range of Australasian albatrosses, petrels and penguins (Marchant & Higgins 2004).

Arrow squid in New Zealand waters have been reported to feed on myctophids, sprats, pilchards, barracouta, euphausiids, mysids, isopods and squid, probably other arrow squid (Yatsu 1986, Uozumi 1998). Uozumi found that the importance of various food items changed between years, and the percentage of empty stomachs was influenced by area, season, size, maturation, and time of day. In Australia, *N. gouldi* was found to feed mostly on pilchard, barracouta, and crustaceans (O'Sullivan & Cullen 1983). Cannibalism was also recorded.

4.2 Bycatch (fish and invertebrates)

Based on models using observer and fisher-reported data, total bycatch in the arrow squid trawl fishery ranged from 4500 to 25 000 t per year between 1991 and 2010–11 (Anderson 2013). Over that time period arrow squid comprised about 80% of the total estimated catch recorded by observers in this fishery (Figure 2). The remainder of the observed catch comprised mainly the commercial fish species barracouta (8.5%), spiny dogfish (1.7%), and jack mackerel (1.1%). Invertebrate species made up a much smaller fraction of the bycatch overall (about 1%), but crabs (0.8%), especially the smooth red swimming crab (*Nectocarcinus bennetti*, 0.5%), were frequently caught.




Figure 2: Percentage of the total catch contributed by the main bycatch species (those representing 0.05% or more of the total catch) in the observed portion of the arrow squid fishery, and the percentage discarded. The Other category is the sum of all bycatch species representing less than 0.05% of the total catch (Anderson 2013).

ARROW SQUID (SQU)

Estimated total annual discards ranged from just over 200 t in 1995–96 to about 5500 in 2001–02 and, like bycatch, peaked in the early 1990s and were at relatively low levels after 2006–07 (Anderson 2013). Most discards were QMS species (about 62% for all years), followed by non-QMS species (19%), invertebrate species (11%), and arrow squid (7%). Absolute levels of discards increased in all categories over the 21-year period; this increase was strongly significant for non-QMS species and total discards, and also marginally significant for QMS species and invertebrates. The species discarded in the greatest amounts were spiny dogfish, redbait, rattails, and silver dory. Discards peaked at 0.13 kg of discarded fish for every 1 kg of arrow squid caught in the early 1990s and declined to 0.02–0.07 kg after 2002–03.

4.3 Incidental Capture of Protected Species (seabirds, mammals, and protected fish)

For protected species, capture estimates presented here include all animals recovered to the deck (alive, injured or dead) of fishing vessels but do not include any cryptic mortality (e.g., seabirds struck by a warp but not brought onboard the vessel, Middleton & Abraham 2007).

4.3.1 NZ sea lion interactions

The New Zealand sea lion (rāpoka) *Phocartos hookeri*, is the rarest sea lion in the world. The estimated total population of around 11 800 sea lions in 2015 was previously classified by the Department of Conservation as ‘Nationally Critical’ under the New Zealand Threat Classification System (Baker et al 2016); the NZ threat status will be updated in 2018. New Zealand sea lions were classified in 2016 as ‘Endangered’ by the International Union for Conservation of Nature (IUCN) on the basis of a projected ongoing decline in pup production of 4% per year at the largest breeding colonies on the Auckland Islands. Pup production at the main Auckland Island rookeries showed a steady decline between 1998 and 2009 and has subsequently stabilised (details can be found in the Aquatic Environment and Biodiversity Annual Review, MPI 2017).

Sea lions forage to depths of up to 600 m and overlap with trawling at up to 500 m depth for arrow squid. Sea lions interact with some trawl fisheries which can result in incidental capture and subsequent drowning (Smith & Baird 2005, 2007a & b, Thompson & Abraham 2010a, Thompson & Abraham 2012, Abraham & Thompson 2011, Abraham et al 2016). Since 1988, incidental captures of sea lions have been monitored by government observers on-board an increasing proportion of the fishing fleet. Since the 2012–13 fishing year, more than 80% of fishing trawls in the SQU 6T fishery have been observed each year.

Efforts to mitigate incidental captures in fisheries have focused on the SQU 6T fishery. From 2017, advice to manage sea lion interactions in this fishery has been developed in consultation with the Squid 6T Operational Plan Technical Advisory Group, including representatives from government and stakeholder groups as well as technical experts and advisors. Under the present Operational Plan, adopted in December 2017, MPI sets a fishing-related mortality limit (FRML) for sea lions in the Auckland Islands squid trawl fishery (SQU 6T) based on estimation of a Population Sustainability Threshold (PST) using a Bayesian population dynamic model (Roberts & Doonan 2016). The PST represents the maximum number of anthropogenic mortalities that the population can sustain while still achieving a defined population objective. For the Auckland Islands sea lion population, the choice of population objective underlying the current PST is as follows: ‘Fisheries mortalities will be limited to ensure that the impacted population is no more than 5% lower than it would otherwise be in the absence of fishing mortality, with 90% confidence, over five years’.

SLEDs were first utilised on some vessels in the SQU 6T fishing fleet in 2001–02. SLED use increased in subsequent years. The use of SLEDs is not mandatory, but use of a certified SLED is required by the current industry body (the Deepwater Group) and is necessary to receive the ‘Discount Rate’ relative to the tow limit applied by MPI). For these reasons, from 2006–07 a standardised model Mark 13/3 SLED has been universally employed by all vessels in the SQU 6T fleet. SLED deployment is monitored and audited by MPI observers.

In 1992, the Ministry adopted a fisheries-related mortality limit (FRML; previously referred to as a maximum allowable level of fisheries-related mortality or MALFiRM) to set an upper limit on the number of New Zealand sea lions that can be incidentally killed each year in the SQU 6T trawl fishery (Chilvers 2008). If this limit is reached, the fishery will be closed for the remainder of the season.

The original 'MALFiRM' was calculated using the potential biological removal approach (PBR; Wade 1998) and was used from 1992–93 to 2003–04 (Smith & Baird 2007a). Since 2003–04 the FRML has been translated into a maximum permitted number of tows calculated from assumed interaction and SLED efficacy rates, regardless of the number of observed New Zealand sea lion captures. This approach was taken because since the introduction of SLEDs, observed sea lion captures are no longer a reliable index of the number of sea lions interacting with the net, and there is uncertainty about the survival rate of sea lions exiting the net via the SLED ('SLED efficacy'); for this reason the number of sea lion deaths from fishery interactions cannot be observed directly. Instead, a management setting meant to approximate the interaction rate, i.e., the 'Strike Rate' is set by MPI and multiplied by a second setting, the 'Discount Rate' representing SLED efficacy, to inform a proxy estimate of potential sea lion fatalities per 100 tows. This proxy estimate is then used to set an effort limit on the operation of the fishery, to ensure that estimated sea lion mortalities remain below the FRML.

Since the introduction of SLEDs, observed capture rates have declined substantially and observer coverage has increased in the SQU 6T fishery (Table 4). Subsequently, statistical models formerly used to estimate interaction rates and SLED efficacy rates (Abraham et al 2016) became increasingly uncertain, because these rates are inversely correlated and, since the introduction of SLEDs, are no longer informed by observed captures data. For this reason Fisheries New Zealand no longer estimates interaction rates, and is progressing research to inform the direct estimation of cryptic mortalities (i.e. un-observable deaths) as a function of observed captures.

Table 3: Fisheries-related mortality limit (FRML) from 1991 to 2015 (♀ = females; numbers in parentheses are FRMLs modified in-season). Direct comparisons among years are not useful because the assumptions underlying the FRML changed over time.

Year	FRML	Discount rate	Management actions
1991–92	16 (♀)		
1992–93	63		
1993–94	63		
1994–95	69		
1995–96	73		Fishery closed by MFiSh (4 May)
1996–97	79		Fishery closed by MFiSh (28 Mar)
1997–98	63		Fishery closed by MFiSh (27 Mar)
1998–99	64		
1999–00	65		Fishery closed by MFiSh (8 Mar)
2000–01	75		Voluntary withdrawal by industry
2001–02	79		Fishery closed by MFiSh (13 Apr)
2002–03	70		Fishery closed by MFiSh (29 Mar), overturned by High Court
2003–04	62 (124)	20%	Fishery closed by MFiSh (22 Mar), overturned by High Court
2004–05	115	20%	Voluntary withdrawal by industry on reaching the FRML
2005–06	97 (150)	20%	FRML increased in mid-March due to abundance of squid
2006–07	93	20%	
2007–08	81	35%	
2008–09	113 (95)	35%	Lower interim limit agreed following decrease in pup numbers
2009–10	76	35%	
2010–11	68	35%	
2011–12	68	35%	
2012–13	68	82%	
2013–14	68	82%	
2014–15	68	82%	
2015–16	68	82%	
2016–17	68	82%	
2017–18	38	75%	

ARROW SQUID (SQU)

Table 4: Annual trawl effort, observer coverage, observed numbers of sea lions captured, observed capture rate (sea lions per 100 trawls), estimated sea lion captures, interactions, and the estimated strike or capture rate (with 95% confidence intervals) for the squid trawl fisheries operating in SQU 6T (Auckland Islands). Estimates are based on methods described in Abraham et al (2016) and available via <https://data.dragonfly.co.nz/psc>. Data for 1995–96 to 2014–15 are based on data version 2016v01.

Year	Tow	Obs. captures			Est. captures		Est. interactions		Est. Interaction rate	
		% obs.	No.	Rate	Mean	95% c.i.	Mean	95% c.i.	Mean	95% c.i.
1995–96	4 468	12.5	13	2.3	130	69–223	129	69–223	2.9	1.5–5
1996–97	3 721	19.8	28	3.8	140	92–208	140	90–211	3.8	2.4–5.7
1997–98	1 442	23.2	15	4.5	59	32–101	59	31–102	4.1	2.1–7.1
1998–99	403	38.7	5	3.2	14	7–26	14	5–27	3.5	1.2–6.7
1999–00	1 206	36.3	25	5.7	69	45–105	69	44–107	5.7	3.6–8.9
2000–01	583	99.1	39	6.7	39	39–40	62	41–85	10.6	7–14.6
2001–02	1 647	34.2	21	3.7	42	29–63	73	44–114	4.4	2.7–6.9
2002–03	1 466	28.4	11	2.6	18	12–28	47	25–79	3.2	1.7–5.4
2003–04	2 594	30.6	16	2	39	26–59	206	104–383	7.9	4–14.8
2004–05	2 693	29.9	9	1.1	30	16–49	167	76–323	6.2	2.8–12
2005–06	2 459	22.4	10	1.8	26	15–43	153	65–306	6.2	2.6–12.4
2006–07	1 317	40.7	7	1.3	15	9–25	93	33–216	7.1	2.5–16.4
2007–08	1 265	46.7	5	0.8	12	6–22	160	24–804	12.6	1.9–63.6
2008–09	1 925	39.6	2	0.3	7	2–15	134	14–672	7	0.7–34.9
2009–10	1 188	25.5	3	1	12	5–26	165	22–818	13.9	1.9–68.9
2010–11	1 583	34.6	0	0	3	0–10	90	5–501	5.7	0.3–31.6
2011–12	1 281	44.6	0	0	2	0–6	60	3–319	4.7	0.2–24.9
2012–13	1 027	86.2	3	0.3	4	3–6	73	8–384	7.1	0.8–37.4
2013–14	737	84.4	2	0.3	2	2–4	47	5–231	6.4	0.7–31.3
2014–15	633	88.3	1	0.2	1	1–3	44	3–236	7	0.5–37.3
2015–16	1 367	92.2	0	0						
2016–17	1 280	70.4	3	0.3						

* SLEDs were introduced. ^ SLEDs were standardised and in widespread use.

Observed sea lion captures in the squid fishery on the Stewart Snares shelf are low (less than one observed capture per year), with high observer coverage (Table 5). In choosing management settings for the SQU 6T fishery, the FRML is reduced by 1 to account for one potential sea lion mortality per year occurring in the SQU 1T fishery.

Table 5: Number of tows by fishing year and observed NZ sea lion captures in squid trawl fisheries on the Stewart-Snares shelf, 2002–03 to 2016–17. No. obs, number of observed tows; % obs, percentage of tows observed; Rate, number of captures per 100 observed tows. Estimates are based on methods described in Abraham et al (2016) and available via <https://data.dragonfly.co.nz/psc>. Data for 2002–03 to 2014–15 are based on data version 2016v1.

	Fishing effort			Observed captures		Estimated interactions	
	Tows	No. obs	% obs	Captures	Rate	Mean	95% c.i.
2002–03	3 281	506	15.4	0	0	2	0–5
2003–04	4 534	957	21.1	1	0.1	3	1–7
2004–05	5 861	1 582	27	3	0.19	6	3–10
2005–06	4 481	537	12	1	0.19	3	1–7
2006–07	2 925	706	24.1	1	0.14	2	1–5
2007–08	2 412	866	35.9	0	0	1	0–3
2008–09	1 808	532	29.4	0	0	1	0–3
2009–10	2 258	765	33.9	1	0.13	2	1–4
2010–11	2 176	685	31.5	0	0	1	0–3
2011–12	1 981	798	40.3	0	0	1	0–2
2012–13	1 528	1 342	87.8	0	0	0	0–1
2013–14	1 222	1 081	88.5	0	0	0	0–1
2014–15	1 116	1 047	93.8	1	0.1	0	0–1
2015–16	988	923	93.4	0	0		
2016–17	1 115	906	81.3	0	0		

A quantitative risk assessment of all threats to the New Zealand sea lion was undertaken to inform the development of a Threat Management Plan for the species. The risk assessment process used for the development of the TMP aimed to quantify which threats pose most risk to the population, and inform the prioritisation of management actions that would meet the management goals of the TMP. The

approach involved the development of demographic models, compilation of data on threats, a risk triage process and detailed modelling of key threats where sufficient data was available. A panel of national and international experts was convened to guide and review the process and provide opinion-based input where data availability was poor. For the Auckland Islands, the greatest risks identified from the triage were; *Klebsiella* disease, commercial trawl fishing, male aggression, trophic effects/prey availability, hookworm disease and wallows.

As the base of the risk assessment, a demographic assessment model were developed for females at the Auckland Islands (where the major squid trawl fishery 6T operates adjacent to), integrating information from mark-recapture observations, pup census and the estimated age distribution of lactating females. Good fits were obtained to all three types of observation and the model structure and parameter estimates appeared to be a good representation of demographic processes that have affected population decline there (primarily low pup survival and low adult survival) (Roberts & Doonan 2016).

Best-estimate projections were undertaken for commercial trawl related mortality, *Klebsiella pneumoniae*-related mortality of pups, trophic effects (food limitation), pups drowning in wallows, male aggression and hookworm mortality and these were compared with the base run – a continuation of demographic rates since 2005 ($\lambda_{2037} = 0.961$, 95% CI 0.890–1.020). A positive growth rate was obtained only with the alleviation of *Klebsiella* ($\lambda_{2037} = 1.005$, 95% CI 0.926–1.069). When assuming the most pessimistic view of cryptic mortality (all interactions resulted in mortality and associated death of pups), alleviating the effects of commercial trawl-related mortality resulted in an increased population growth rate relative to the base run, but did not reverse the declining trend ($\lambda_{2037} = 0.977$, 95% CI 0.902–1.036). The alleviation of trophic effects (food limitation) had the next greatest effect ($\lambda_{2037} = 0.974$, 95% CI 0.905–1.038) and all other threats had a minor effect relative to the base run projection (increase in λ_{2037} of less than 0.01) (Roberts & Doonan 2016).

Results from the risk assessment at the Auckland Islands indicated that alleviation of any one threat will not result in an increasing population. Similarly none of the major threats assessed were sufficient alone to explain the observed decline in pup production at the Auckland Islands. Clearly multiple factors were acting on the population, and for management to recover the species a holistic view must be adopted. Further studies will be needed to fully understand, and development management options for some of the key threats, such as trophic effects and *Klebsiella* disease.

4.3.2 NZ fur seal interactions

The New Zealand fur seal was classified in 2008 as “Least Concern” by IUCN and in 2010 as “Not Threatened” under the NZ Threat Classification System.

Vessels targeting arrow squid incidentally catch fur seals (Baird & Smith 2007a, Smith & Baird 2009, Thompson & Abraham 2010b, Baird 2011, Abraham et al 2016), mostly off the east coast South Island, on the Stewart-Snares shelf, and close to the Auckland Islands. In the 2016–17 fishing year there were 17 observed captures of New Zealand fur seal in squid trawl fisheries. The rate of capture over the period 2002–03 and 2016–17 varied from 0.1 to 1.1 captures per hundred tows without obvious trend (Table 6). Estimated capture rates from Abraham et al (2016) (available via <https://data.dragonfly.co.nz/psc>) are not reproduced here pending resolution of identified structural issues in the model related to the partition between model strata with contrasting capture rates, resulting in implausibly high estimates of uncertainty despite high observer coverage.

ARROW SQUID (SQU)

Table 6: Number of tows by fishing year and observed total NZ fur seal captures in squid trawl fisheries, 2002–03 to 2016–17.

	<u>Fishing effort</u>			<u>Observed</u>	
	Tows	No.	%	Capture	Rate
2002–03	8 410	1 308	15.6	8	0.6
2003–04	8 336	1 771	21.2	16	0.9
2004–05	10 489	2 512	23.9	15	0.6
2005–06	8 576	1 103	12.9	4	0.4
2006–07	5 906	1 289	21.8	9	0.7
2007–08	4 236	1 459	34.4	6	0.4
2008–09	3 867	1 299	33.6	1	0.1
2009–10	3 789	1 071	28.3	8	0.7
2010–11	4 214	1 263	30	8	0.6
2011–12	3 505	1 380	39.4	8	0.6
2012–13	2 646	2 273	85.9	7	0.3
2013–14	2 051	1 787	87.1	10	0.6
2014–15	1 950	1 694	86.9	19	1.1
2015–16	2 896	2 363	81.6	10	0.4
2016–17	2 594	1 934	74.6	17	0.9

4.3.3 Seabird interactions

Vessels targeting arrow squid incidentally catch seabirds. Baird (2005a) summarised observed seabird captures in the arrow squid target fishery for the fishing years 1998–99 to 2002–03 and calculated total seabird captures for the areas with adequate observer coverage using ratio based estimations. Baird & Smith (2007b, 2008) summarised observed seabird captures and used both ratio-based and model-based predictions to estimate the total seabird captures for 2003–04, 2004–05 and 2005–06. Abraham & Thompson (2011) summarised captures of protected species and used model and ratio-based predictions of the total seabird captures for 1989–90 and 2008–09.

A consistent modelling framework was developed to estimate the captures for ten species (and species groups), using hierarchical mixed-effects generalised linear model (GLM), fitted using Bayesian methods (Abraham et al 2016, Abraham & Richard 2017, 2018).

In the 2015–16 fishing year there were 302 observed captures of birds in squid trawl fisheries, and 361 estimated captures (95% c.i.: 324–441), with the estimates made using a statistical model (Table 7, Abraham et al 2016). In the 2016–17, there were 261 observed captures of seabirds in squid trawl fisheries, however estimates of total captures are not available (Table 7).

Table 7: Number of tows by fishing year and observed and model-estimated total bird captures in squid trawl fisheries, 2002–03 to 2016–17. No. obs, number of observed tows; % obs, percentage of tows observed; Rate, number of captures per 100 observed tows. Estimates are based on methods described in Abraham et al (2016) and Abraham & Richard (2017, 2018) and are available via <https://data.dragonfly.co.nz/psc>. Estimates from 2002–03 to 2015–16 are based on data version 2017v1.

	Tows	<u>Observed</u>				<u>Estimated</u>	
		No. obs	% obs	Captures	Rate	Captures	95% c.i.
2002–03	8 410	1 308	15.6	154	11.8	954	748–1 219
2003–04	8 336	1 771	21.2	194	11	885	712–1 106
2004–05	10 489	2 512	23.9	351	14	1 338	1 122–1 596
2005–06	8 576	1 103	12.9	195	17.7	1 213	954–1 538
2006–07	5 906	1 289	21.8	126	9.8	596	456–796
2007–08	4 236	1 459	34.4	162	11.1	492	391–627
2008–09	3 867	1 299	33.6	259	19.9	661	549–806
2009–10	3 789	1 071	28.3	92	8.6	422	322–561
2010–11	4 214	1 263	30	166	13.1	588	463–753
2011–12	3 505	1 380	39.4	106	7.7	350	272–452
2012–13	2 646	2 273	85.9	458	20.1	521	486–578
2013–14	2 051	1 787	87.1	200	11.2	237	214–274
2014–15	1 950	1 694	86.9	384	22.7	428	396–489
2015–16	2 896	2 363	81.6	302	12.8	361	324–441
2016–17	2 594	1 934	74.6	261	13.5		

Total estimated seabird captures in squid trawl fisheries varied from 237 to 1338 between 2002–03 and 2015–16 at a rate of 7.7 to 22.7 captures per hundred tows without obvious trend (Table 7). These estimates include all bird species and should be interpreted with caution because trends by species can be masked. The average capture rate in squid trawl fisheries over the last thirteen years is about 13.79 birds per 100 tows, a high rate relative to trawl fisheries for scampi (4.43 birds per 100 tows) and hoki (2.32 birds per 100 tows) over the same years.

Observed seabird captures since 2002–03 have been dominated by four species: white-capped and southern Buller’s albatrosses make up 83% and 13% of the albatrosses captured, respectively; and white-chinned petrels and sooty shearwaters make up 56% and 41% of other birds, respectively, the total and fishery risk ratios presented in Table 8. Most captures occur on the Stewart-Snares shelf (63%) or close to the Auckland Islands (36%). These numbers should be regarded as only a general guide on the distribution of captures because observer coverage is not uniform across areas and may not be representative.

The squid target fishery contributes to the total risk posed by New Zealand commercial fishing to seabirds. The two species to which the fishery poses the most risk are Southern Buller’s albatross and New Zealand white-capped albatross, with this target fishery poses 0.048 and 0.028 of PST (Table 9). Southern Buller’s albatross and New Zealand white-capped albatross were both assessed at high risk (Richard et al 2017).

Mitigation methods such as streamer (tori) lines, Brady bird bafflers, warp deflectors, and offal management are used in the squid trawl fishery. Warp mitigation was voluntarily introduced from about 2004 and made mandatory in April 2006 (Ministry of Fisheries 2006). The 2006 notice mandated that all trawlers over 28 m in length use a seabird scaring device while trawling (being “paired streamer lines”, “bird baffle” or “warp deflector” as defined in the notice). During the 2005–06 fishing year a large trial of mitigation devices was conducted in the squid fishery (Middleton & Abraham 2007). Eighteen vessels were involved in the trial which used observations of seabird heavily contacting the trawl warps (‘warp strikes’) to quantify the effect of using three mitigation devices; paired streamer/tori lines, four boom bird bafflers and warp scarers. Few warp strikes occurred in the absence of offal discharge. When offal was present the tori lines were most effective at reducing warp strikes. All mitigation devices were more effective for reducing large bird warp strikes than small bird. There were, however, about as many bird strikes on the tori lines as the number of strikes on unmitigated warps. The effect of these strikes has not been assessed (Middleton & Abraham 2007).

The three year average warp capture rate of white-capped albatross (84% of albatross observed caught in this fishery) before warp mitigation was made mandatory at the start of the 2005-06 fishing year was higher than 3 per 100 tows in hoki target trawls until the three year period from 2003-04 to 2005-06. Since 2005-06 to 2007-08, the three year warp capture rate has decreased to below 1 per 100 tows. For this same species the three year average capture rates from nets has fluctuated over this time period (Figure 3).

Table 8: Number of observed seabird captures in squid trawl fisheries, 2002–03 to 2016–17, by species and area. The risk category is an estimate of aggregate potential fatalities across trawl and longline fisheries relative to the Population Sustainability Threshold, PST (from Richard et al 2017, where full details of the risk assessment approach can be found). It is not an estimate of the risk posed by trawl fishing for squid alone

	Risk category	Auckland Islands	Chatham Rise	East Coast South Island	Fiordland	Stewart Snares Shelf	Subantarctic	Total
New Zealand white-capped albatross	High	399		3	11	525		938
Southern Buller's albatross	High	46			8	98		152
Salvin's albatross	High	1		4		17	1	23
Southern Royal albatross	Negligible					6		6
Campbell black-browed albatross	Low	1						1
Albatross spp.	-	4				1		5
Black-browed albatross	-	1						1
Buller's albatross	-				1			1
Royal albatross spp.	-					1		1
Total albatrosses		452	0	7	20	648	1	1128
White-chinned petrel	Negligible	493				633	2	1128
Sooty shearwater	Negligible	177		22	5	618		822
Antarctic prion	Negligible	34						34
Common diving petrel	Negligible	6				3		9
Cape petrel	Negligible				1	1		2
Fairy prion	Negligible	2						2
Black-bellied storm petrel	Negligible	1						1
Grey petrel	Negligible			1				1
New Zealand white-faced storm petrel	Negligible					1		1
White-headed petrel	Negligible	1						1
mid-sized petrels & shearwaters	-	8				1		9
Giant petrel spp.	-					7		7
Grey-backed storm petrel	-	3						3
Gadfly petrels	-	1						1
Prion spp.	-	1						1
Seabirds	-					1		1
Total other birds		727	0	23	6	1265	2	2023

Table 9: Risk ratio of seabirds predicted by the level two risk assessment for the squid target trawl fishery and all fisheries included in the level two risk assessment, 2006–07 to 2016–17, showing seabird species with a risk ratio of at least 0.001 of Population Sustainability Threshold, PST (from Richard et al 2017, where full details of the risk assessment approach can be found). The risk ratio is an estimate of aggregate potential fatalities across trawl and longline fisheries relative to the PST. The DOC threat classifications are shown (Robertson et al 2017 at <http://www.doc.govt.nz/Documents/science-and-technical/nztcsl9entire.pdf>).

Species name	PST (mean)	Risk ratio		Risk category	DOC Threat Classification
		Squid target trawl	TOTAL		
Southern Buller's albatross	1368.4	0.048	0.392	High	At Risk: Naturally Uncommon
New Zealand white-capped albatross	10900.3	0.028	0.353	High	At Risk: Declining
White-chinned petrel	25614.6	0.009	0.055	Negligible	At Risk: Declining
Salvin's albatross	3599.5	0.002	0.780	High	Threatened: Nationally Critical
Northern royal albatross	715.1	0.001	0.043	Low	At Risk: Naturally Uncommon




Figure 3: Three year rolling average of capture rates of white-capped albatross in squid trawl fisheries for warp and net captures.

4.4 Benthic interactions

Between 1989–90 and 2004–05, 131 973 trawl tows for squid on or within 1 m of the seabed were reported, comprising 13.7% of all trawl tows on or within 1 m of the seabed reported on TCEPR forms in those years (range 8–23% by year, Baird et al 2011). Black et al (2013) estimated that hoki arrow squid has accounted for 13.5% of all tows reported on TCEPR forms since 1989–90. Between 2006–07 and 2010–11, 95% of arrow squid catch was reported on TCEPR forms. The great majority of tows are conducted on the Stewart-Snares shelf or north and east of the Auckland Islands, with smaller numbers off the east coast of the South Island and the Chatham Rise. Tows were located in Benthic Optimised Marine Environment Classification (BOMECE, Leathwick et al 2009) classes E (outer shelf), F, H (upper slope), I, J, L, and M (mid-slope) (Baird & Wood 2012), and 92% were between 100 and 300 m depth (Baird et al 2011). Tables 4–7 show that the number of trawl tows for squid varies between years, largely without trend and presumably in response to variations in the abundance of squid and management measures to limit the number of sea lions caught. The average duration of trawls has increased over this time so the trend in aggregate swept area will not be the same.

Bottom trawling for squid, like trawling for other species, is likely to have effects on benthic community structure and function (e.g., see Rice 2006 for an international review) and there may be consequences for benthic productivity (e.g., Jennings et al 2001, Hermesen et al 2003, Hiddink et al 2006, Reiss et al 2009). These are not considered in detail here but are discussed in the 2012 Aquatic Environment and Biodiversity Annual Review.

4.5 Other considerations

A substantial decline in the west coast jig fishery for squid will have reduced any trophic implications of that fishery.

5. STOCK ASSESSMENT

Arrow squid live for one year, spawn once then die. Every squid fishing season is therefore based on what amounts to a new stock. It is not possible to calculate reliable yield estimates from historical catch and effort data for a resource which has not yet hatched, even when including data which are just one year old. Furthermore, because of the short life span and rapid growth of arrow squid, it is not possible to estimate the biomass prior to the fishing season. Moreover, the biomass increases rapidly during the season and then decreases to low levels as the animals spawn and die.

5.1 Estimates of fishery parameters and abundance

No estimates are available.

5.2 Biomass estimates

Biomass estimates are not available for squid.

5.3 Yield estimates and projections

It is not possible to estimate *MCY*.

It is not possible to estimate *CAY*.

5.4 Other yield estimates and stock assessment results

There are no other yield estimates of stock assessment results available for arrow squid.

5.5 Other factors

N. gouldi spawns one to two months before *N. sloanii*. This means that at any given time *N. gouldi* is older and larger than *N. sloanii*. The annual squid jigging fishery begins on *N. gouldi* and at some time during the season the biomass of *N. sloanii* will exceed that of *N. gouldi* and the fleet will move south. If *N. sloanii* are abundant the fleet will remain in the south fishing for *N. sloanii*. If *N. sloanii* are less abundant the fleet will return north and resume fishing *N. gouldi*.

6. STATUS OF THE STOCKS

No estimates of current and reference biomass are available. There is also no proven method at this time to estimate yields from the squid fishery before a fishing season begins based on biomass estimates or CPUE data.

Because squid live for about one year, spawn and then die, and because the fishery is so variable, it is not practical to predict future stock size in advance of the fishing season. As a consequence, it is not possible to estimate a long-term sustainable yield for squid, nor determine if recent catch levels or the current TACC will allow the stock to move towards a size that will support the *MSY*. There will be some years in which economic or other factors will prevent the TACC from being fully taken, while in other years the TACC may be lower than the potential yield. It is not known whether New Zealand squid stocks have ever been stressed through fishing mortality.

TACCs and reported landings for the 2016–17 fishing year are summarised in Table 9.

Table 9: Summary of TACCs (t) and reported landings (t) of arrow squid for the most recent fishing year.

Fishstock	2016–17 Actual TACC	2016–17 Reported landings
SQU 1J	5 000	1
SQU 1T	44 741	7 735
SQU 6T	32 369	10 726
SQU 10T	10	0
Total	82 120	18 462

7. FOR FURTHER INFORMATION

- Abraham, E.R.; Berkenbusch, K.; Richard, Y.; Thompson, F. (2016) Summary of the capture of seabirds, mammals, and turtles in New Zealand commercial fisheries, 2002–03 to 2012–13. *New Zealand Aquatic Environment and Biodiversity Report No.169*. 205 p.
- Abraham, E R (2011) Probability of Mild Traumatic Brain Injury for sea lions interacting with SLEDs. Final Research Report for project SRP2011-03 (Unpublished report held by FNZ, Wellington). 21 p.
- Abraham, E R; Richard, Y (2017) Summary of the capture of seabirds in New Zealand commercial fisheries, 2002–03 to 2013–14. *New Zealand Aquatic Environment and Biodiversity Report No. 184*. 88 p.
- Abraham, E R; Richard, Y (2018) Estimated capture of seabirds in New Zealand trawl and longline fisheries, 2002–03 to 2014–15. *New Zealand Aquatic Environment and Biodiversity Report No. 197*. 97 p.
- Abraham, E R; Thompson, F N (2011) Summary of the capture of seabirds, marine mammals, and turtles in New Zealand commercial fisheries, 1998–99 to 2008–09. *New Zealand Aquatic Environment and Biodiversity Report No. 80* 172 p.
- Abraham, E R; Thompson, F N; Berkenbusch, K (2013) Estimated capture of seabirds in New Zealand trawl and longline fisheries, 2002–03 to 2010–11. Final Research Report for Ministry for Primary Industries project PRO2010-01 (Unpublished report held by FNZ, Wellington).
- Abraham, E R; Thompson, F N; Oliver, M D (2010) Summary of the capture of seabirds, mammals, and turtles in New Zealand commercial fisheries, 1998–99 to 2007–08. *New Zealand Aquatic Environment and Biodiversity Report No.45*. 149 p.
- Anderson, O F (2004) Fish discards and non-target fish catch in the trawl fisheries for arrow squid, jack mackerel, and scampi in New Zealand waters. *New Zealand Fisheries Assessment Report. 2004/10*. 61 p.
- Anderson, O F (2013) Fish and invertebrate bycatch and discards in New Zealand arrow squid fisheries from 1990–91 until 2010–11. *New Zealand Aquatic Environment and Biodiversity Report No. 112*. 62 p.
- Baker, C.S.; Chilvers, B.L.; Childerhouse, S.; Constantine, R.; Currey, R.; Mattlin, R.; van Helden, A.; Hitchmough, R.; Rolfe, J. 2016: Conservation status of New Zealand marine mammals, 2013. New Zealand Threat Classification Series 14. Department of Conservation, Wellington. 18 p.
- Baird, S J (2005a) Incidental capture of seabird species in commercial fisheries in New Zealand waters, 2002–03. *New Zealand Fisheries Assessment Report 2005/2*. 50 p.
- Baird, S J (2005b) Incidental capture of *Phocarctos hookeri* (New Zealand sea lions) in New Zealand commercial fisheries, 2001–02. *New Zealand Fisheries Assessment Report. 2005/08*.
- Baird, S J (2005c) Incidental capture of *Phocarctos hookeri* (New Zealand sea lions) in New Zealand commercial fisheries, 2002–03. *New Zealand Fisheries Assessment Report. 2005/09*.
- Baird, S J (2011) New Zealand fur seals – summary of current knowledge. *New Zealand Aquatic Environment and Biodiversity Report No. 72*. 50 p.
- Baird, S J; Doonan, I J (2005) *Phocarctos hookeri* (New Zealand sea lions): incidental captures in New Zealand commercial fisheries during 2000–01 and in-season estimates of captures during squid trawling in SQU 6T in 2002. *New Zealand Fisheries Assessment Report, 2005/17*.
- Baird, S J; Smith, M H (2007a) Incidental capture of New Zealand fur seals (*Arctocephalus forsteri*) in commercial fisheries in New Zealand waters, 2003–04 to 2004–05. *New Zealand Aquatic Environment and Biodiversity Report No. 14*. 98 p.
- Baird, S J; Smith, M H (2007b) Incidental capture of seabird species in commercial fisheries in New Zealand waters, 2003–04 and 2004–05. *New Zealand Aquatic Environment and Biodiversity Report No. 9*. 108 p.
- Baird, S J; Smith, M H (2008) Incidental capture of seabird species in commercial fisheries in New Zealand waters, 2005–06. *New Zealand Aquatic Environment and Biodiversity Report No. 18*. 124 p.
- Baird, S J; Wood, B A (2012) Extent of coverage of 15 environmental classes within the New Zealand EEZ by commercial trawling with seafloor contact. *New Zealand Aquatic Environment and Biodiversity Report No. 89*.
- Baird, S J; Wood, B A; Bagley, N W (2011) Nature and extent of commercial fishing effort on or near the seafloor within the New Zealand 200 n. mile Exclusive Economic Zone, 1989–90 to 2004–05. *New Zealand Aquatic Environment and Biodiversity Report No. 73*.
- Baker, C S; Chilvers, B L; Constantine, R; DuFresne, S; Mattlin, R H; van Helden, A; Hitchmough, R (2010) Conservation status of New Zealand marine mammals (suborders Cetacea and Pinnipedia), 2009. *New Zealand Journal of Marine and Freshwater Research* 44: 101–115.
- Baker, C S; Chilvers, B L; Childerhouse, S; Constantine, R; Currey, R; Mattlin, R; van Helden, A; Hitchmough, R; Rolfe, J (2016) Conservation status of New Zealand marine mammals, 2013. *New Zealand Threat Classification Series 14. Department of Conservation, Wellington*. 18 p
- Ballara, S L; Anderson, O F (2009) Fish discards and non-target fish catch in the trawl fisheries for arrow squid and scampi in New Zealand waters. *New Zealand Aquatic Environment and Biodiversity Report No. 38*. 102 p.
- Black, J; Wood, R; Berthelsen, T; Tilney, R (2013) Monitoring New Zealand's trawl footprint for deepwater fisheries: 1989–1990 to 2009–2010. *New Zealand Aquatic Environment and Biodiversity Report No. 110*. 57 p.
- Boren, L (2008) Diet of New Zealand fur seals (*Arctocephalus forsteri*): a summary. *DOC Research & Development Series 319*. 19 p.
- Bradford, E (2002) Estimation of the variance of mean catch rates and total catches of non-target species in New Zealand fisheries. *New Zealand Fisheries Assessment Report 2002/54*. 60 p.
- Breen, P A; Fu, D; Gilbert, D J (2010) Sea lion population modelling and management procedure evaluations: Report for Project SAP2008/14, Objective 2. Final Research Report for Ministry of Fisheries project SAP2008/14, Objective 2. (Unpublished report held by FNZ, Wellington.)
- Brothers, N; Duckworth, A R; Safina, C; Gilman, E L (2010) Seabird bycatch in pelagic longline fisheries is grossly underestimated when using only haul data. *PloS One* 5: e12491. doi: 10.1371/journal.pone.001249
- Chilvers, B L (2008) New Zealand sea lions *Phocarctos hookeri* and squid trawl fisheries: bycatch problems and management options. *Endangered Species Research* 5: 193–204.
- Doonan, I J (1995) Incidental catch of Hooker's sea lion in the southern trawl fishery for squid, summer 1994. Fisheries Assessment Research Document. 1995/22. (Unpublished document held by NIWA library, Wellington.)
- Doonan, I J (2000) Estimation of Hooker's sea lion *Phocarctos hookeri* captures in the southern squid trawl fisheries, 2000. *New Zealand Fisheries Assessment Report. 2000/41*.
- Doonan, I J (2001) Estimation of Hooker's sea lion *Phocarctos hookeri* captures in the southern squid trawl fisheries, 2001. *New Zealand Fisheries Assessment Report. 2001/67*.
- Dunn, M (2009) Feeding habits of the ommastrephid squid *Nototodarus sloanii* on the Chatham Rise, New Zealand. *New Zealand Journal of Marine and Freshwater Research*, 43(5): 1103–1113.
- Dunn, M., Horn, P, Connell, A, Stevens, D, Forman, J, Pinkerton, M, Griggs, L, Notman, P, Wood, B. (2009) Ecosystem-scale trophic relationships: diet composition and guild structure of middle-depth fish on the Chatham Rise. Final Research Report for Ministry of Fisheries Project, ZBD2004-02. (Unpublished report held by FNZ, Wellington.)

ARROW SQUID (SQU)


- Fea, N I; Harcourt, R; Lalas, C (1999) Seasonal variation in the diet of New Zealand fur seals (*Arctocephalus forsteri*) at Otago Peninsula, New Zealand. *Wildlife Research* 26(2): 147–160.
- Förch, E C (1983) Squid - current research. In: Taylor, J L; Baird, G G (eds.) New Zealand finfish fisheries: the resources and their management, pp. 33–34. Trade Publications Ltd., Auckland.
- Gibson, D; Jones, J B (1993) Fed up with parasites? — old fish are. *Marine Biology* 117: 495–500.
- Gibson, D J M (1995) The New Zealand Squid Fishery, 1979–93. *MAF Fisheries Technical Report No 42*. 43 p.
- Harcourt, R G; Bradshaw, C J A; Dickson, K; Davis, L S (2002) Foraging ecology of a generalist predator, the female New Zealand fur seal. *Marine Ecology Progress Series* 227: 11–24.
- Hedd, H; Gales, R (2001) The diet of shy albatrosses (*Thalassarche cauta*) at Albatross Island, Tasmania. *Journal of Zoology* 253(1): 69–90.
- Hermesen, J M; Collie, J S; Valentine, P C (2003) Mobile fishing gear reduces benthic megafaunal production on Georges Bank. *Marine Ecology Progress Series* 260: 97–108.
- Hiddink, J G; Jennings, S; Kaiser, M J; Queiros, A M; Duplisea, D E; Piet, G J (2006) Cumulative impacts of seabed trawl disturbance on benthic biomass, production, and species richness in different habitats. *Canadian Journal of Fisheries and Aquatic Sciences* 63:721–36.
- Hurst, R J; Ballara, S L; MacGibbon, D; Triantafillos, L (2012) Fishery characterisation and standardised CPUE analyses for arrow squid (*Nototodarus gouldi* and *N. sloanii*), 1989–90 to 2007–08, and potential management approaches for southern fisheries. *New Zealand Fisheries Assessment Report 2012/47*. 303 p.
- James, G D; Stahl, J C (2000) Diet of the Buller's albatross (*Diomedea bulleri bulleri*) and the importance of fishery discards during chick rearing. *New Zealand Journal of Marine and Freshwater Research* 34: 435–454.
- Jennings, S; Dinmore, T A; Duplisea, D E; Warr, K J; Lancaster, J E (2001) Trawling disturbance can modify benthic production processes. *Journal of Animal Ecology* 70: 459–475.
- Jones, M R L (2007) Historic trawl data and recent information infers temporal change in the occurrence of squid in the diet of orange roughy (*Hoplostethus atlanticus* Collett) in New Zealand. *Reviews in Fish Biology and Fisheries* 17(2–3): 493–499.
- Langley, A D (2001) Summary of catch and effort data from the SQU 1J, SQU 1T, and SQU 6T fisheries for 1989–90 to 1999–2000. *New Zealand Fisheries Assessment Report*. 2001/51.
- Leathwick, J R; Rowden, A; Nodder, S; Gorman, R; Bardsley, S; Pinkerton, M; Baird, S J; Hadfield, M; Currie, K; Goh, A (2009) Benthic-optimised marine environment classification for New Zealand waters. Final Research Report project for Ministry of Fisheries project BEN2006/01. 52 p. (Unpublished report held by FNZ, Wellington.)
- McKenzie, D; Fletcher, D (2006) Characterisation of seabird captures in commercial trawl and longline fisheries in New Zealand 1997/98 to 2003/04. Final Research Report for Ministry of Fisheries project ENV2004/04. (Unpublished report held by FNZ, Wellington.) 102 p.
- Marchant, S; Higgins, P J (2004) (Co-ordinators). Handbook of Australian, New Zealand and Antarctic birds. Volume 1 Ratites to Ducks. Oxford University Press, Melbourne, Australia. 735 p.
- Mattlin, R H (1983) Squid. In: Taylor, J. L.; Baird, G. G. (eds.) New Zealand finfish fisheries: the resources and their management, pp. 30–32. Trade Publications Ltd., Auckland.
- Mattlin, R H; Colman, J A (1988) Arrow squid. New Zealand Fisheries Assessment Research Document 88/34. 16 p. (Unpublished document held in NIWA library, Wellington).
- Mattlin, R H; Scheibling, R E; Förch, E C (1985) Distribution, abundance and size structure of arrow squid (*Nototodarus* sp.) off New Zealand. *Northwest Atlantic Fisheries Organisation Scientific Council Studies* 9: 39–45.
- Meynier, L; Mackenzie, D D S; Duignan, P J; Chilvers, B L; Morel, P C H (2009) Variability in the diet of New Zealand sea lion (*Phocarcos hookeri*) at the Auckland Islands, New Zealand. *Marine Mammal Science* 25(2): 302–326.
- Meynier, L; Stockin, K A; Bando, M K H; Duignan, P J (2008) Stomach contents of common dolphin (*Delphinus* sp.) from New Zealand waters. *New Zealand Journal of Marine and Freshwater Research* 42: 257–268.
- Middleton, D A J; Abraham, E R (2007) The efficacy of warp strike mitigation devices: Trials in the 2006 squid fishery. Final Research Report for Ministry of Fisheries project IPA2006/02. (Unpublished report held by FNZ, Wellington).
- Ministry of Fisheries (2006) Fisheries (Commercial Fishing) Amendment Regulations 2006. Seabird Scaring Devices – Circular Issued under Authority of the Fisheries (Commercial Fishing) Amendment Regulations 2006 (No. F361). *New Zealand Gazette No. 33*, 6 April 2006. 1 p.
- MPI (2016) Aquatic Environment and Biodiversity Annual Review 2016. Compiled by the Fisheries Management Science. Ministry for Primary Industries, Wellington, New Zealand. 790 p.
- MPI (2017) Operational Plan to Manage the Incidental Capture of New Zealand Sea Lions 2016–2017. Ministry for Primary Industries, Wellington, New Zealand. 10 p. <http://www.mpi.govt.nz/document-vault/16564>
- O'Sullivan, D; Cullen, J M (1983) Food of the squid *Nototodarus gouldi* in Bass Strait. *Australian Journal of Marine and Freshwater Research* 34: 261–285.
- Reiss, H; Greenstreet, S P R; Sieben, K; Ehrich, S; Piet, G J; Quirijns, F; Robinson, L; Wolff, W J; Kröncke, I (2009) Effects of fishing disturbance on benthic communities and secondary production within an intensively fished area. *Marine Ecology Progress Series* 394: 201–213.
- Rice, J (2006) Impacts of Mobile Bottom Gears on Seafloor Habitats, Species, and Communities: A Review and Synthesis of Selected International Reviews. *Canadian Science Advisory Secretariat Research Document* 2006/057. 35 p. (available from http://www.dfo-mpo.gc.ca/CSAS/Csas/DocREC/2006/RES2006_057_e.pdf).
- Richard, Y; Abraham, E R (2013) Risk of commercial fisheries to New Zealand seabird populations. *New Zealand Aquatic Environment and Biodiversity Report No. 109*. 58 p.
- Richard, Y.; Abraham, E.R. (2015). Assessment of the risk of commercial fisheries to New Zealand seabirds, 2006–07 to 2012–13. *New Zealand Aquatic Environment and Biodiversity Report* 162. 85 p
- Richard, Y.; Abraham, E R; Berkenbusch, K (2017) Assessment of the risk of commercial fisheries to New Zealand seabirds, 2006–07 to 2014–15. *New Zealand Aquatic Environment and Biodiversity Report No. 191*. 133 p.
- Roberts, J; Doonan, I (2016) Quantitative Risk Assessment of Threats to New Zealand Sea Lions. *New Zealand Aquatic Environment and Biodiversity Report No. 166*. 111 p.
- Robertson, H A; Baird, K; Dowding, J E; Elliott, G P; Hitchmough, R A; Miskelly, C M; McArthur, N; O'Donnell, C F J.; Sagar, P M; Scofield, R P; Taylor, G A (2017) Conservation status of New Zealand birds, 2016. *New Zealand Threat Classification Series 19*. Department of Conservation, Wellington. 23 p.
- Smith, M H; Baird, S J (2005a) Representativeness of past observer coverage, and future coverage required for estimation of New Zealand sea lion (*Phocarcos hookeri*) captures in the SQU 6T fishery. *New Zealand Fisheries Assessment Report*. 2005/05.
- Smith, M H; Baird, S J (2005b) Factors that may influence the level of incidental mortality of New Zealand sea lions (*Phocarcos hookeri*) in the squid (*Notodarus* spp.) trawl fishery in SQU 6T. *New Zealand Fisheries Assessment Report* 2005/20.
- Smith, M H; Baird, S J (2007a) Estimation of incidental captures of New Zealand sea lions (*Phocarcos hookeri*) in New Zealand fisheries in 2003–04, with particular reference to the SQU 6T squid trawl fishery. *New Zealand Fisheries Assessment Report* 2007/7. 32 p.

- Smith, M H; Baird, S J (2007b) Estimation of incidental captures of New Zealand sea lions (*Phocartos hookeri*) in New Zealand fisheries in 2004–05, with particular reference to the SQU 6T squid trawl fishery. *New Zealand Aquatic Environment and Biodiversity Report No. 12*. 31 p.
- Smith, M H; Baird, S J (2009) Model-based estimation of New Zealand fur seal (*Arctocephalus forsteri*) incidental captures and strike rates for trawl fishing in New Zealand waters for the years 1994–95 to 2005–06. *New Zealand Aquatic Environment and Biodiversity Report No. 40*. 92 p.
- Smith, P J; Mattlin, R H; Roeleveld, M A; Okutani, T (1987) Arrow squids of the genus *Nototodarus* in New Zealand waters: systematics, biology and fisheries. *New Zealand Journal of Marine and Freshwater Research* 21: 315–326.
- Stevens, D W; Hurst, R J; Bagley, N W (2012) Feeding habits of New Zealand fishes; a literature review and summary of research trawl database records 1960 to 2000. *New Zealand Aquatic Environment and Biodiversity Report No. 85*.
- Stockin, K A (2008) The New Zealand common dolphin (*Delphinus* sp.): identity, ecology and conservation : a thesis submitted in partial fulfilment of the requirements for the degree of Doctor of Philosophy in Zoology, Massey University, Auckland, New Zealand.
- Thompson, D R (2008) Autopsy report for seabirds killed and returned from New Zealand fisheries, 1 October 2006 to 30 September 2007. Report prepared for the Conservation Services Programme, Department of Conservation: Contract INT2006/02. Available on www.doc.govt.nz
- Thompson, D R (2009) Seabird Autopsy Project: Summary Report for the 2007–08 Fishing Year. Report prepared for the Conservation Services Programme, Department of Conservation: Contract INT2007/02. Available on www.doc.govt.nz
- Thompson, F N; Abraham, E R (2010a) Estimation of the capture of New Zealand sea lions (*Phocartos hookeri*) in trawl fisheries, from 1995–96 to 2008–09. *New Zealand Aquatic Environment and Biodiversity Report* 66.
- Thompson, F N; Abraham, E R (2010b) Estimation of fur seal (*Arctocephalus forsteri*) bycatch in New Zealand trawl fisheries, 2002–03 to 2008–09. *New Zealand Aquatic Environment and Biodiversity Report No. 61*. 37 p.
- Thompson, F N; Abraham, E R (2012) Captures of New Zealand sea lion in squid trawl fisheries, during the 2008/2009 fishing year. Retrieved from <http://bycatch.dragonfly.co.nz/v20120315/new-zealand-sea-lion/squid-trawl/all-vessels/eez/2009/>, Apr 27, 2012
- Thompson, F N; Abraham, E R; Berkenbusch, K (2011) Marine mammal bycatch in New Zealand trawl fisheries, 1995–96 to 2009–10. Final Research Report for Ministry of Fisheries project PRO2010-01 (Unpublished report held by FNZ, Wellington). 80 p.
- Thompson, F N; Abraham, E R; Oliver, M D (2010a) Estimation of fur seal bycatch in New Zealand sea lions trawl fisheries, 2002–03 to 2007–08. *New Zealand Aquatic Environment and Biodiversity Report No. 56*.
- Thompson, F N; Berkenbusch, K; Abraham, E R (2013) Marine mammal bycatch in New Zealand trawl fisheries, 1995–96 to 2010–11. *New Zealand Aquatic Environment and Biodiversity Report No. 105*. 73 p.
- Thompson, F N; Oliver, M D; Abraham, E R (2010b) Estimation of the capture of New Zealand sea lions (*Phocartos hookeri*) in trawl fisheries, from 1995–96 to 2007–08. *New Zealand Aquatic Environment and Biodiversity Report No. 52*.
- Uozumi, Y (1998) Fishery biology of arrow squids, *Nototodarus gouldi* and *sloanii*, in New Zealand waters. *Bulletin of the National Institute of Far Seas Fisheries* 35. 111 p.
- Uozumi, Y; Ohara, H (1992) Age and growth of *Nototodarus sloanii* (Cephalopoda: Oegopsida) based on daily increment counts in statoliths. *Nippon Suisan Gakkaishi* 59: 1469–1477.
- Wade, P R (1998) Calculating limits to the allowable human-caused mortality of cetaceans and pinnipeds. *Marine Mammal Science* 14(1):1–37.
- Yatsu, A (1986) Feeding habit of *N. sloanii* caught by bottom trawl. *Japanese Marine Fishery Resource research Centre* 30: 45–52.

BARRACOUTA (BAR)

(Thyrsites atun)

Manga, maka



1. FISHERY SUMMARY

1.1 Commercial fisheries

Barracouta are caught in coastal waters around mainland New Zealand, The Snares and Chatham Islands, down to about 400 m and have been managed under the Quota Management System since 1 October 1986. Historical catch summaries are given in Tables 1 and 2. Catches by New Zealand vessels increased significantly in the late 1960s and total annual catch peaked at about 47 000 t in 1977, with the addition of foreign vessels around New Zealand. Between 1983–84 and 2013–14, catches fluctuated between 18 000 and 29 000 t per annum (Table 3), at an average 25 000 t. Figure 1 shows the historical landings and TACC values for the main BAR stocks.

Table 1: Reported landings (t) for the main QMAs from 1931 to 1982.

Year	BAR 1	BAR 4	BAR 5	BAR 7	Year	BAR 1	BAR 4	BAR 5	BAR 7
1931–32	4	0	0	0	1957	163	0	20	80
1932–33	55	0	0	77	1958	146	0	15	78
1933–34	5	0	1	0	1959	139	0	18	71
1934–35	36	0	0	52	1960	117	0	13	90
1935–36	1	0	0	0	1961	187	0	22	68
1936–37	26	0	0	35	1962	104	0	25	44
1937–38	21	0	0	26	1963	63	0	4	20
1938–39	91	0	22	55	1964	66	0	4	21
1939–40	107	0	27	50	1965	111	0	1	76
1940–41	153	0	53	30	1966	62	0	1	116
1941–42	212	0	86	17	1967	53	0	1	178
1942–43	371	0	151	20	1968	10 113	0	3	1 196
1943–44	192	0	79	7	1969	8 499	0	2	5 756
1944	247	0	97	50	1970	12 984	0	2	3 960
1945	306	0	114	32	1971	11 327	0	191	4 006
1946	391	0	125	63	1972	29 307	2	86	3 487
1947	590	0	213	45	1973	14 856	0	79	4 698
1948	466	0	172	27	1974	23 420	0	106	9 028
1949	425	0	169	40	1975	8 985	0	855	6 257
1950	430	0	153	76	1976	19 124	5	495	6 795
1951	266	0	95	47	1977	69 81	9 095	2 041	33 266
1952	190	0	56	68	1978	6 833	17	1 162	6 918
1953	202	0	41	77	1979	6 474	4 057	3 380	5 263
1954	166	0	35	38	1980	5 649	1 854	7 867	5 146
1955	139	0	14	58	1981	6 993	2 030	8 311	11 141
1956	165	0	16	45	1982	5 393	787	6 909	7 064

Notes:

1. The 1931–1943 years are April–March but from 1944 onwards are calendar years.
2. Data up to 1985 are from fishing returns: Data from 1986 to 1990 are from Quota Management Reports.
3. Data for the period 1931 to 1982 are based on reported landings by harbour and are likely to be underestimated as a result of under-reporting and discarding practices. Data includes both foreign and domestic landings. Data were aggregated to FMA using methods and assumptions described by Francis & Paul (2013).

BARRACOUTA (BAR)

Table 2: Reported landings (t) by nationality from 1977 to 1987–88.

Fishing Year	New Zealand		Foreign			Total	
	Domestic	Chartered	Japan	Korea	USSR	(FSU)	(QMS)
1977	4 697	0	34 357	8 109	0	47 163	-
1978–79	5 335	58	4 781	2 481	0	12 655	-
1979–80	7 748	6 679	4 339	3 879	47	22 922	-
1980–81	10 058	4 995	4 227	15	60	19 355	-
1981–82	12 055	11 077	2 813	373	0	26 328	-
1982–83	10 814	7 110	1 746	1 888	31	21 589	-
1983–83*	7 763	2 961	803	1 115	0	12 642	-
1983–84	12 390	10 226	1 786	4 355	0	28 757	-
1984–85	7 869	10 425	1 430	5 252	0	24 976	-
1985–86	8 427	7 865	1 371	815	0	18 478	-
1986–87	9 829	13 732	1 575	742	0	25 878	27 660†
1987–88	9 335	12 077	896	609	0	22 971	26 607†

* 6 month changeover in fishing years.

† The discrepancies between QMS and FSU total landings are due to under-reporting to the FSU.

Over 99% of the recorded catch is taken by trawlers. Major target fisheries have been developed on spring spawning aggregations (Chatham Islands, Stewart Island, west coast South Island and northern and central east coast South Island) as well as on summer feeding aggregations, particularly around The Snares and on the east coast of the South Island. Barracouta also comprise a significant proportion of the bycatch in the west coast North Island jack mackerel fishery, The Snares squid fishery, and the east coast South Island red cod and tarakihi fisheries. Catches have increased in recent years in BAR 1 to the level of the TACC, but have dropped in BAR 4 in the last three years. The TACC in BAR 5 was reduced from 9282 t to 7470 t on 1 October 1998 with a 2 t customary and 3 t recreational allocation and a TAC of 7475 t. It was increased to 8200 t in 2015. In BAR 7 the catch limit was exceeded from 2004–05 to 2006–07 (catches nearly reached 15 000 t in 2006–07), but catch has decreased since, to well below the TACC.

Table 3: Reported landings (t) of barracouta by Fishstock from 1983–84 to 2016–17 and actual TACCs (t) from 1986–87 to 2016–17. QMS data from 1986-present. [Continued on next page]

Fishstock FMAs	BAR 1 1, 2, 3		BAR 4 4		BAR 5 5 & 6		BAR 7 7, 8, 9	
	Landings	TACC	Landing	TACC	Landings	TACC	Landings	TACC
1983–84*	7 805	-	1 743	-	11 291	-	7 222	-
1984–85*	5 442	-	1 909	-	12 487	-	4 425	-
1985–86*	5 395	-	1 509	-	6 380	-	4 536	-
1986–87	8 877	8 510	3 084	3 010	7 653	9 010	8 046	10 510
1987–88	9 256	8 837	1 775	3 010	6 457	9 011	9 117	10 603
1988–89	5 838	9 426	946	3 010	5 323	9 011	8 071	10 702
1989–90	9 209	9 841	1 349	3 016	5 960	9 282	7 050	10 925
1990–91	9 401	9 957	1 399	3 016	8 817	9 282	7 138	10 925
1991–92	6 733	9 957	1 156	3 016	6 897	9 282	7 326	10 925
1992–93	9 032	9 969	2 251	3 016	7 019	9 282	10 141	10 925
1993–94	7 299	9 969	606	3 016	3 410	9 282	8 030	10 925
1994–95	10 023	9 969	331	3 016	2 645	9 282	9 345	10 925
1995–96	11 252	9 969	2 234	3 016	4 255	9 282	8 593	10 925
1996–97	11 873	11 000	1 081	3 016	2 839	9 282	10 203	10 925
1997–98	11 543	11 000	1 966	3 016	6 167	9 282	8 717	10 925
1998–99	9 229	11 000	459	3 016	7 302	7 470	4 427	10 925
1999–00	10 032	11 000	1 911	3 016	6 205	7 470	3 288	10 925
2000–01	7 118	11 000	2 122	3 016	6 101	7 470	6 890	10 925
2001–02	6 900	11 000	1 160	3 019	5 883	7 470	7 655	11 173
2002–03	7 595	11 000	573	3 019	7 843	7 470	9 025	11 173
2003–04	5 949	11 000	477	3 019	6 919	7 470	9 114	11 173
2004–05	6 085	11 000	98	3 019	8 593	7 470	12 156	11 173
2005–06	7 030	11 000	687	3 019	9 479	7 470	10 685	11 173
2006–07	5 351	11 000	3 233	3 019	6 334	7 470	14 699	11 173
2007–08	5 987	11 000	2 975	3 019	8 561	7 470	10 451	11 173
2008–09	8 861	11 000	968	3 019	7 659	7 470	8 955	11 173
2009–10	10 635	11 000	1 223	3 019	6 951	7 470	9 642	11 173
2010–11	11 420	11 000	1 190	3 019	8 201	7 470	6 129	11 173
2011–12	9 305	11 000	1 423	3 019	7 071	7 470	8 643	11 173
2012–13	9 740	11 000	706	3 019	7 931	7 470	6 897	11 173
2013–14	11 309	11 000	1 482	3 019	6 886	7 470	6 637	11 173
2014–15	6 902	11 000	3 671	3 019	6 779	7 470	6 974	11 173
2015–16	5 568	11 000	2 893	3 019	7 558	8 200	5 493	11 173
2016–17	9 520	11 000	2 606	3 019	8 916	8 200	7 127	11 173

Table 3 Continued: Reported landings (t) of barracouta by Fishstock from 1983–84 to 2016–17 and actual TACCs (t) from 1986–87 to 2016–17. QMS data from 1986–present.

Fishstock FMAs	BAR 10		Total	
	Landings	TACC	Landings	TACC
1983–84*	0	-	28 061	-
1984–85*	0	-	24 263	-
1985–86*	0	-	17 820	-
1986–87	0	10	27 660	31 050
1987–88	0	10	26 605	31 471
1988–89	0	10	20 178	32 159
1989–90	0	10	23 568	33 073
1990–91	0	10	26 755	33 190
1991–92	0	10	22 212	33 190
1992–93	<1	10	28 443	33 202
1993–94	0	10	19 345	33 202
1994–95	0	10	22 345	33 202
1995–96	0	10	26 334	33 202
1996–97	0	10	25 996	34 233
1997–98	0	10	28 393	34 233
1998–99	0	10	21 417	32 421
1999–00	0	10	21 436	32 421
2000–01	0	10	22 231	32 421
2001–02	0	10	21 598	32 672
2002–03	0	10	25 036	32 672
2003–04	0	10	22 459	32 672
2004–05	0	10	26 919	32 672
2005–06	0	10	27 881	32 672
2006–07	0	10	29 617	32 672
2007–08	0	10	27 968	32 672
2008–09	0	10	26 443	32 672
2009–10	0	10	28 451	32 672
2010–11	0	10	26 937	32 672
2011–12	0	10	26 442	32 672
2012–13	0	10	24 973	32 672
2013–14	0	10	26 313	32 672
2014–15	0	10	24 327	32 672
2015–16	0	10	21 511	33 403
2016–17	0	10	28 169	33 403

* FSU data.




Figure 1: Reported commercial landings and TACC for the four main BAR stocks. BAR 1 (Auckland East), [Continued on next page].

BARRACOUTA (BAR)




Figure 1: [Continued] Reported commercial landings and TACC for the four main BAR stocks. From top to bottom: BAR 4 (Chatham Rise), BAR 5 (Southland), and BAR 7 (Challenger).

1.2 Recreational fisheries

Barracouta are commonly encountered by recreational fishers in New Zealand, more frequently in the southern half of BAR 7 and BAR 1. Barracouta are typically harvested as bait for other fishing rather than for consumption. They are predominantly taken on rod and reel (97.9%) with a small proportion taken by net methods (1.7%). The catch is taken predominantly from boat (95.5%) with a small proportion from land based fishers (4.5%).

1.2.1 Management controls

The main method used to manage recreational harvests of barracouta is daily bag limits. General spatial and method restrictions also apply. Fishers can take up to 30 barracouta as part of their combined daily bag limit in the Fiordland and Southland Fishery Management Areas. There is currently no bag limit in place in the other Fishery Management Areas.

1.2.2 Estimates of recreational harvest

There are two broad approaches to estimating recreational fisheries harvest: the use of onsite or access point methods where fishers are surveyed or counted at the point of fishing or access to their fishing activity; and offsite methods where some form of post-event interview and/or diary are used to collect data from fishers.

The first estimates of recreational harvest for barracouta were calculated using an offsite approach, the offsite regional telephone and diary survey approach. Estimates for 1996 came from a national telephone and diary survey (Bradford 1998). Another national telephone and diary survey was carried out in 2000 (Boyd & Reilly 2002). The harvest estimates provided by these telephone diary surveys (Table 4) are no longer considered reliable.

In response to the cost and scale challenges associated with onsite methods, in particular the difficulties in sampling other than trailer boat fisheries, offsite approaches to estimating recreational fisheries harvest have been revisited. This led to the development and implementation of a national panel survey for the 2011–12 fishing year (Wynne-Jones et al 2014). The panel survey used face-to-face interviews of a random sample of New Zealand households to recruit a panel of fishers and non-fishers for a full year. The panel members were contacted regularly about their fishing activities and catch information collected in standardised phone interviews. Note that the national panel survey estimate does not include recreational harvest taken under s111 general approvals. Recreational catch estimates from the national panel survey are given in Table 4.

Table 4: Recreational harvest estimates for barracouta stocks. Early surveys were carried out in different years in the regions: South in 1991–92, Central in 1992–93, and North in 1993–94 (Teirney et al 1997). The estimated Fishstock harvest is indicative in these surveys and made by combining estimates from the different years. Some early survey harvests are presented as a range to reflect the considerable uncertainty in the estimates. The telephone/diary surveys ran from December to November but are denoted by the January calendar year. The national panel survey ran through the October to September fishing year but is denoted by the January calendar year. A mean weight of 2.14kg was used for the national panel survey.

Fishstock		Survey	Total		
			Number	CV	Survey harvest (t)
BAR 1	1992	South	27 000	47%	30–90
BAR 7	1992	South	2 100	44%	-
BAR 1	1993	Central	17 000	22%	25–35
BAR 7	1993	Central	15 600	24%	25–35
BAR 1	1996	National	68 000	8%	160–190
BAR 7	1996	National	74 000	15%	160–220
BAR 1	2000	National	156 000	35%	182–377
BAR 5	2000	National	2 000	51%	2–7
BAR 7	2000	National	35 000	28%	68–120
BAR 1	2012	Panel survey	22 224		47.7
BAR 5	2012	Panel survey	666		1.4
BAR 7	2012	Panel survey	16 743		35.9
All combined	2012	Panel survey	39 652	18%	85.05

BARRACOUTA (BAR)

1.3 Customary non-commercial fisheries

Quantitative information on the current level of customary non-commercial take is not available.

1.4 Illegal catch

Quantitative information on the level of illegal catch is not available.

1.5 Other sources of mortality

There may have been considerable amounts of barracouta discarded prior to the QMS, either because of quota restrictions under the deepwater policy, low value, or undesirable small size fish. There is also likely to be some mortality associated with escapement from trawl nets. Some discarding may also have occurred in BAR 1 because of the lack of quota availability and the high deemed value in relation to the low value of the fish.

2. BIOLOGY

Barracouta spawn mainly in late-winter/spring (August–September) on the east and west coasts of both of the main islands, and in late spring (November–December) in Southland and in the Chatham Islands. Some spawning activity may also extend into summer/autumn. Sexual maturity is reached at about 50–60 cm fork length (FL) at about 2–3 years of age.

Juvenile barracouta have been recorded from inshore areas (less than 100 m) all around New Zealand and the Chatham Islands, although they appear to be less common on the west coast of the South Island. Adult fish are found down to about 400 m depth. Tagging experiments indicated that mature fish from the east coast South Island waters migrate after June to northern waters off the east coast North Island to spawn during August–September; research survey results and commercial fishing patterns show some consistency with this movement (see Hurst et al 2012).

No age data are available for the period prior to the onset of commercial fishing, which developed rapidly from 1968. Ageing studies carried out in the mid-1970s showed that the maximum age rarely exceeded 10 years.

M was estimated using the equation $M = \log_e 100/\text{maximum age}$, where maximum age is the age to which 1% of the population survives in an unexploited stock. Using 10 years for the maximum age suggests an M of up to 0.46. The effect of fishing on age structure prior to the mid-1970s is unknown, but M is unlikely to be less than 0.3, which has been assumed in previous stock assessments.

Biological parameters relevant to the stock assessment are shown in Table 5.

Table 5: Estimates of biological parameters.

Fishstock		Estimate		Source	
<u>1. Natural mortality (<i>M</i>)</u>				Hurst (unpub. data)	
All-both sexes		Less than 0.46 <i>M</i> = 0.30 considered best estimate for all areas for both sexes			
<u>2. Weight = a(length)^b (Weight in g, length in cm fork length).</u>					
	Females		Males		
	a	b	a	b	
BAR 4	0.0074	2.94	0.0117	2.82	Hurst & Bagley (1992)
BAR 5	0.0075	2090	0.0075	2.90	Hurst & Bagley (1992)
<u>3. Von Bertalanffy growth parameters</u>					
	Both sexes				
	<i>K</i>	<i>t</i> ₀	<i>L</i> _∞		
Tasmania	0.45	0.166	91.17	(unconstrained)	Grant et al (1978)
	0.42	-0.25	91.01	(constrained, <i>t</i> ₀ fixed)	
Southland	0.336	-0.35	81.1	Male	Horn (2002)
	0.259	-0.60	89.3	Female	Horn (2002)

3. STOCKS AND AREAS

There are thought to be at least four main stocks, based on known spawning locations and movements. Stock boundaries are not well understood, but the Chatham Islands stock is probably separate. There may be some overlap between mainland stock management areas as currently defined from analysis of tagging data, commercial fishery data, biological data (i.e., length frequencies, otoliths, parasites, spawning areas and seasons) and from seasonal relative biomass estimates. In particular, it appears that there is considerable overlap of Southland fish with other areas, probably the west coast of the South Island and possibly the east coast as well. However, there are not enough data at this stage to alter the existing stock boundaries.

4. STOCK ASSESSMENT

Recent stock assessments have been conducted for BAR 1 and BAR 5. Hurst et al (2012) provided a comprehensive characterisation of all barracouta stocks and provided CPUE indices for BAR 1 (east coast South Island), BAR 4 (west coast South Island), and BAR 5 for 1989–90 to 2007–08. McGregor (in prep.) characterised the fisheries and estimated CPUE indices for the fisheries on the WCNI and WCSI (BAR 7) and the southern Snares fishery (BAR 5). Marsh (in prep.) updated CPUE indices for BAR 5 to 2015. In BAR 4 the fishery has been highly variable and no standardised analysis is possible.

A time series of trawl surveys was carried out in the Southland area (QMA 5) in February–March from 1993 to 1996 using *Tangaroa* (Table 6). Trawl surveys on the east and west coasts of the South Island in autumn using *Kaharoa* may help interpretation of trends in biomass around the South Island. The long time series of trawl surveys on the Chatham Rise (deeper than 200 m) and Sub-Antarctic (deeper than 300 m) using *Tangaroa* are not considered to adequately survey the preferred depth range of barracouta.

4.1 BAR 1 Auckland (E), Central (E), South-East (Coast)

4.1.1 Estimates of fishery parameters and abundance

The results from trawl surveys carried out during the mid 1980s (sometimes from a variety of different vessels) were used to provide an approximate estimate of minimum absolute biomass. This approach required an assumption about catchability to convert the trawl survey catches to estimates of absolute biomass. This method is now considered obsolete and the estimates of absolute biomass have not been included.

4.1.2 Biomass estimates

There is no trawl survey series for BAR 1 off the east coast of the North Island. The trawl survey information discussed below is for the east coast of the South Island.

The ECSI winter surveys from 1991 to 1996 in 30–400 m were replaced by summer trawl surveys (1996–97 to 2000–01) which also included the 10–30 m depth range, but these were discontinued after the fifth in the annual time series because of the extreme fluctuations in catchability between surveys (Francis et al 2001). The winter surveys were reinstated in 2007 and this time included additional 10–30 m strata in an attempt to index elephant fish and red gurnard which were added to the list of target species. Only the 2007, 2012, 2014 and 2016 surveys provide full coverage of the 10–30 m depth range.

The 2014 barracouta biomass estimate was the highest recorded in the east coast South Island winter trawl survey time series core strata (30–400 m). Biomass in the east coast South Island winter trawl survey time series core strata (30–400 m) steadily increased until 2014 when it was more than four-fold larger than the average biomass of the early 1990s, before a 57% decline in 2016 (Table 6, Figure 2). The additional biomass captured in the 10–30 m depth has ranged from 1 to 15% of the biomass in the core plus shallow strata (10–400 m), and in 2016 it accounted for 14%, indicating that shallow strata should continue to be monitored for this species.

A comparison of the pre-recruit and recruited biomass (where recruited fish are over 60 cm long) for the ECSI winter survey, based on the core strata, is shown in Figure 3. During the 1991–93 surveys, the

BARRACOUTA (BAR)

pre-recruit and recruited estimates were similar, but in 1994 and 1996, most of the total biomass was from the recruited fish. For the renewed series, from 2007, the main increase has come from the recruited fish, with significantly higher biomass for recruited fish compared with pre-recruits in the 2009 and 2012 surveys. The 2014 survey indicated an increase in the pre-recruit biomass, although the uncertainty around this estimate is high, and in 2016 both recruited and pre-recruited biomass declined substantially.




Figure 2: Barracouta total biomass and 95% confidence intervals for the all ECSI winter surveys in core strata (30–400 m), and core plus shallow strata (10–400 m) in 2007, 2012, 2014, and 2016.




Figure 3: Barracouta pre-recruit and recruited biomass estimates and associated confidence intervals from the ECSI winter trawl survey core strata (30–400 m). Recruited fish were defined as fish over 60 cm fork length.

Table 6: Relative biomass indices (t) and coefficients of variation (CV) for barracouta for east coast South Island (ECSI) - winter, east coast North Island (ECNI), west coast South Island (WCSI) and Southland survey areas. Biomass estimates for ECSI in 1991 have been adjusted to allow for non-sampled strata (7 & 9 equivalent to current strata 13, 16 and 17). – , not measured; NA, not applicable.

Region	Fishstock	Year	Trip number	Total Biomass estimate	CV (%)	Total Biomass estimate	CV (%)
					30–400 m	10–400 m	
ECSI (winter)	BAR 1	1991	KAH9105	8 361	29	-	-
		1992	KAH9205	11 672	23	-	-
		1993	KAH9306	18 197	22	-	-
		1994	KAH9406	6 965	34	-	-
		1996	KAH9608	16 848	19	-	-
		2007	KAH0705	21 132	17	24 939	19
		2008	KAH0806	25 544	16	-	-
		2009	KAH0905	33 360	16	-	-
		2012	KAH1207	34 325	17	36 526	16
		2014	KAH1402	46 563	19	46 903	19
		2016	KAH1605	19 708	27	23 007	24
ECNI	BAR 1	1993	KAH9304	2 673	15	-	-
		1994	KAH9402	8 433	33	-	-
		1995	KAH9502	2 103	29	-	-
		1996	KAH9602	2 495	23	-	-
WCSI	BAR 7	1992	KAH9203	2 478	14	-	-
		1994	KAH9404	5 298	16	-	-
		1995	KAH9504	4 480	13	-	-
		1997	KAH9701	2 993	19	-	-
		2000	KAH0004	1 787	11	-	-
		2003	KAH0304	4 485	20	-	-
		2005	KAH0503	2 763	13	-	-
Southland	BAR 5	1993	TAN9301	11 587	18	-	-
		1994	TAN9402	6 151	20	-	-
		1995	TAN9502	4 539	17	-	-
		1996	TAN9604	7 693	19	-	-

4.1.3 Length frequency distributions

The length distributions from the east coast South Island winter trawl survey show at least three clear pre-recruit modes at about 20 cm, 35 cm, and 50 cm (combined males, females, and unsexed) consistent with ages of 0+, 1+, and 2+ (Figure 4). Length frequency distributions are consistent among the surveys, showing the presence of the pre-recruited cohorts, with indications that these could be tracked through time (modal progression) (Beentjes et al 2015, 2016). The addition of the 10–30 m depth range does not change the shape of the length distributions (not shown in Figure 4).

BARRACOUTA (BAR)




Figure 4: Scaled length frequency distributions for barracouta in core strata (30–400 m) for the ECSI winter surveys. n, number of fish measured; no., core strata population estimates; c.v., coefficient of variation.




Figure 5 : East coast South Island part of BAR 1 CPUE indices from the standardised lognormal, binomial, and the combined (delta lognormal) models, based on the merged day-level CELR and TCER data for 1989–90 to 2013–14.

4.1.4 CPUE indices

Two sets of standardised CPUE indices were derived for BAR 1: one for the northern waters off the east coast of the North Island (ECNI) and one for the east coast South Island, ECSI (Baird 2016). Each set had three CPUE series defined by form type: a merged CELR/TCER day-level model for 1989–90 to 2013–14; a TCER tow-level model for 2007–08 to 2013–14; and a TCEPR tow-level model for 1989–90 to 2013–14. All ECNI series were rejected by the Working Group because of shifts in targeting through time, high inter-annual variability, and unacceptably low levels of data. Thus, the following sections on CPUE pertain to the ECSI waters only.

Three standardised CPUE series for the east coast South Island part of BAR 1 were prepared, as outlined above, using data from 1989–90 to 2013–14, with each series based on the catch of barracouta in bottom trawl fisheries defined by different target species, including barracouta (Baird 2016). Two CPUE series were rejected by the SINS Working Group: the CPUE index based on the TCEPR data (targeting barracouta, red cod, and arrow squid), primarily because of inter-annual inconsistencies in the underlying catch and effort data; and the short TCER series with only seven years of data.

The SINS Working Group accepted the combined index (delta lognormal model) series based on the daily data from CELR and TCER forms (targeting barracouta, red cod, and tarakihi) as an index of abundance for BAR 1 (Figure 5). After a peak period during 1996–97 and 1997–98, there was a period of relatively lower CPUE from 1998–99 to 2008–09, followed by an increase up to 2012–13, to a level similar to the earlier peak. The most recent index (2013–14) showed a modest drop, but remained above the series mean. The TCER tow-level CPUE series, for which additional explanatory variables were incorporated into the model, was very similar to the CELR/TCER day-level series for the overlapping period (2007–08 to 2013–14). Figure 6 provides a comparison of the ECSI indices with the ECSI winter trawl survey indices. The increase in abundance measured by the trawl survey for 2007 onwards follows a similar trajectory to that for the ECSI CELR/TCER indices.

BARRACOUTA (BAR)




Figure 6: Comparison of the BAR 1 ECSI delta-lognormal indices for 1990–2014 and the recruited biomass (and associated variance) from the ECSI winter trawl survey series. The recruited biomass is based on fish over 60 cm fork length. Each series has been standardised to the mean for concurrent years.

4.2 BAR 5 Southland, Sub-Antarctic

4.2.1 CPUE indices

Marsh (in prep) used unmerged (tow level) data to fit CPUE indices for barracouta to various target fisheries in the BAR 5 region. The WG agreed that the CPUE from the SQU target fishery in Statistical Area 028 was the best series of abundance indices for BAR 5. An alternative CPUE index based on the target BAR and WAR tows was suggested as a sensitivity run. Both series show high catch rates since 2007. The base case CPUE declines from 1990 to 1995, then increases and decreases again until 2007 but after 2007 the index increases and remains high through to 2015 (Figure 7). The alternative series increases from 1995 to 2007 and then oscillates at high catch rates through to 2015 (Figure 8). The current stock status is unknown, due to the lack of a quantitative assessment for this stock.




Figure 7 : Base case BAR 5 CPUE Model: CPUE indices for barracouta from SQU target tows in Statistical Area 028 (1990–2015).




Figure 8 : Alternative BAR 5 CPUE Model: CPUE indices for barracouta from BAR and WAR target tows (1990–2015).

4.3 BAR 7 Challenger, Central (W) Auckland (W)

4.3.1 CPUE indices

McGregor (in prep.) looked at the separate fisheries on the WCNI and WCSI. The three CPUE options for the WCNI all gave similar patterns to the inshore *Kaharoa* WCSI trawl survey. The WG considered that the tow level CPUE was the best data to use to monitor this stock. The CPUE shows an increasing trend from 2000 to 2004 and is then generally flat (Figure 9).




Figure 9: West Coast CPUE for Models 2b (tow level), 3 (JMA target) and 4 (no target) and Trawl Survey abundance index for calendar years 1990–2010. Model 3 (JMA target) is actually based on fishing years, months Nov–May, whereas the other models here are calendar year, Jun–Nov. Trawl survey is based on fishing year.

The WCSI data series shows a similar increase from 2000 and is then generally flat, for the tow level CPUE based on all target from June to October (Figure 10).

BARRACOUTA (BAR)




Figure 10: West Coast South Island current and previous CPUE, West Coast North Island CPUE and trawl survey abundance index for calendar years 1990–2010. Trawl survey is based on fishing year.

4.4 Yield estimates and projections

No estimates of biomass are available for any of the barracouta stocks.

4.5 Other factors

Barracouta are part of the shelf (30–300 m) mixed fishery and are usually the dominant species in these depths around the South Island (except perhaps in good red cod years in the Canterbury Bight). Any increase or decrease in barracouta quotas will have overflow effects onto bycatch species. The economics of targeting on barracouta is probably affected by its availability relative to other more preferred species and this will, in turn, affect fishing patterns.

An analysis of trends in biomass of the Southland fishery suggests that recruitment may have been relatively low in the years after 1989 and that biomass may have declined between surveys by the *Shinkai Maru* (1981 and 1986) and the *Tangaroa* (annually 1993 to 1996). The scale of decline appeared to be greater than could be explained by different catching efficiencies of the two vessels.

5. STATUS OF THE STOCKS

• BAR 1


The current understanding of the BAR 1 stock is that adult barracouta undertake an annual northward migration from the east coast of the South Island to spawn off the east coast of the North Island during July/August–September (see Hurst et al 2012). For the purposes of this analysis barracouta in BAR 1 are assumed to comprise a single stock.

Stock Status	
Year of Most Recent Assessment	2016
Assessment Runs Presented	BAR 1 ECSI CELR/TCER day-level series (target species BAR, RCO, TAR)

Reference Points	Interim Target: B_{MSY} -compatible proxy based on CPUE (average from 1989–90 to 2013–14 of the BAR 1 ECSI CELR/TCER model as defined by Baird (2016)) Soft Limit: 50 % of target Hard Limit: 25 % of target Overfishing threshold: F_{MSY} (assumed)
Status in relation to Target	Very Likely (> 90%) to be at or above the target
Status in relation to Limits	Soft Limit: Very Unlikely (< 10%) to be below Hard Limit: Very Unlikely (< 10%) to be below
Status in relation to Overfishing	Overfishing is Unlikely (< 40%) to be occurring

Historical Stock Status Trajectory and Current Status

CPUE, Catch and TACC Trajectories




Fishery and Stock Trends

Recent trend in Biomass or Proxy

The BAR 1 CPUE series increased steeply from 2002–03 to a peak in 2012–13. The 2013–14 value was lower than the peak, but well above the series mean.

Recent trend in Fishing Mortality or Proxy



BARRACOUTA (BAR)

Other Abundance Indices	The winter ECSI trawl survey series for recruited fish has a trend that is similar to the BAR 1 CPUE index, with a peak in 2014.
Trends in Other Relevant Indicator or Variables	Recent landings (2008–09 to 2013–14) are at a similar level to those recorded during 1994–95 to 1999–2000.


Projections and Prognosis	
Stock Projections or Prognosis	Quantitative stock projections are unavailable.
Probability of Current Catch or TACC causing Biomass to remain below or decline below Limits	Soft Limit: Unlikely (< 40%) as above average pre-recruit abundance was observed in the ECSI trawl survey in 2014. Hard Limit: Unlikely (< 40 %)
Probability of Current Catch or TACC causing Overfishing to continue or to commence	Unknown

Assessment Methodology and Evaluation		
Assessment Type	Level 2: Partial Quantitative Stock Assessment.	
Assessment Method	Standardised CPUE series	
Assessment Dates	Latest assessment: 2016	Next assessment: 2019
Overall assessment quality rank	1 – High Quality	
Main data inputs (rank)	- Catch and effort data - Trawl survey biomass indices and associated length frequencies	1 – High Quality 1 – High Quality
Data not used (rank)	TCEPR CPUE Series (ECSI) Standardised CPUE series (ECNI) Summer ECSI trawl survey data	3 – Low Quality: few vessels and highly variable CPUE 3 – Low Quality: insufficient data and high interannual variability 3 – Low Quality: variable catchability between years
Changes to Model Structure and Assumptions	N/A	
Major Sources of Uncertainty	-	
Qualifying Comments		
-		

Fishery Interactions
Barracouta in the ECSI part of BAR 1 are taken as bycatch by inshore bottom trawl fisheries targeting, amongst others, red cod and tarakihi, and red cod and arrow squid by deepwater vessels. ECSI bycatch also comes from midwater effort targeting jack mackerels. In the ECNI part of BAR 1, most barracouta bycatch is from tarakihi and red gurnard effort; currently, there is little targeting of barracouta in this area. The trawl fishery in the ECSI area is subject to management measures designed to reduce interactions with endemic Hector's dolphins and seabirds. Interactions with other species are currently being characterised.

• **BAR 5**

CPUE analyses were completed for the main fisheries in BAR 5. The relationship between these southern fisheries and the WCSI is uncertain.

Stock Status	
Year of Most Recent Assessment	2017
Assessment Runs Presented	Standardised CPUE Sub-Antarctic (tow level)
Reference Points	Target: 40% B_0 Soft Limit: 20% B_0 Hard Limit: 10% B_0 Overfishing threshold: $F_{40\%B_0}$
Status in relation to Target	Unknown
Status in relation to Limits	B_{2015} is Very Unlikely (< 10%) to be below both the soft and hard limits
Status in relation to Overfishing	Unknown
Historical Stock Status Trajectory and Current Status	
 <p>BAR 5 CPUE Model: CPUE indices for barracouta from SQU target tows in statistical area 028 (1990–2015).</p>	

Fishery and Stock Trends	
Recent Trend in Biomass or Proxy	-
Recent Trend in Fishing Intensity or Proxy	-
Other Abundance Indices	CPUE has remained at a high level since 2008 despite catches at or above the TACC.
Trends in Other Relevant Indicators or Variables	-

Projections and Prognosis	
Stock Projections or Prognosis	-
Probability of Current Catch or TACC causing Biomass to remain below or to decline below Limits	Soft Limit: Very Unlikely (< 10%) Hard Limit: Very Unlikely (< 10%)

BARRACOUTA (BAR)

Probability of Current Catch or TACC causing Overfishing to continue or to commence	Unknown
---	---------

Assessment Methodology and Evaluation		
Assessment Type	Level 2: Partial Quantitative Stock Assessment.	
Assessment Method	Standardised CPUE	
Assessment Dates	Latest assessment: 2016	Next assessment: 2019
Overall assessment quality rank	1 – High Quality	
Main data inputs (rank)	- Commercial CPUE	1 – High Quality
Data not used (rank)	N/A	
Changes to Model Structure and Assumptions	None	
Major sources of Uncertainty	-	

Qualifying Comments
-


Fishery Interactions
Barracouta are taken as a target species in BAR 5 and also as by-catch in the squid and warehou target fisheries. Interactions with other species are currently being characterised.

• BAR 7

CPUE analyses were completed for the main fisheries in BAR 7. The relationship between the WCSI and the fisheries in BAR 5 is uncertain.

Stock Status	
Year of Most Recent Assessment	2016
Assessment Runs Presented	Standardised CPUE (tow level)
Reference Points	Target: 40% B_0 Soft Limit: 20% B_0 Hard Limit: 10% B_0 Overfishing threshold: $F_{40\%B_0}$
Status in relation to Target	Unknown
Status in relation to Limits	B_{2015} is Very Unlikely (< 10%) to be below both the soft and hard limits
Status in relation to Overfishing	Unknown

Historical Stock Status Trajectory and Current Status



West Coast South Island CPUE, West Coast North Island CPUE and trawl survey abundance index for calendar years 1990–2010. Trawl survey is based on fishing year.

Fishery and Stock Trends	
Recent Trend in Biomass or Proxy	-
Recent Trend in Fishing Intensity or Proxy	-
Other Abundance Indices	CPUE has been increasing since 2000.
Trends in Other Relevant Indicators or Variables	-

Projections and Prognosis	
Stock Projections or Prognosis	-
Probability of Current Catch or TACC causing Biomass to remain below or to decline below Limits	Soft Limit: Very Unlikely (< 10%) Hard Limit: Very Unlikely (< 10%)
Probability of Current Catch or TACC causing Overfishing to continue or to commence	Unknown

Assessment Methodology and Evaluation		
Assessment Type	Level 2: Partial Quantitative Stock Assessment.	
Assessment Method	Standardised CPUE	
Assessment Dates	Latest assessment: 2016	Next assessment: 2019
Overall assessment quality rank	1 – High Quality	
Main data inputs (rank)	Commercial CPUE	1 – High Quality
Data not used (rank)	N/A	
Changes to Model Structure and Assumptions	None	
Major sources of Uncertainty	-	

Qualifying Comments
None

Fishery Interactions
Barracouta in BAR 7 are taken as a target on the WCSI and as bycatch in the WCNI jack mackerel and WCSI hoki fisheries. Interactions with other species are currently being characterised.

6. FOR FURTHER INFORMATION


- Bagley, N W; Hurst, R J (1995) Trawl survey of middle depth and inshore bottom species off Southland, February–March 1994 (TAN 9402). *New Zealand Fisheries Data Report* 57. 50 p.
- Bagley, N W; Hurst, R J (1996) Trawl survey of middle depth and inshore bottom species off Southland, February–March 1995 (TAN 9502). *New Zealand Fisheries Data Report* 73. 47 p.
- Bagley, N W; Hurst, R J (1996) Trawl survey of middle depth and inshore bottom species off Southland, February–March 1996 (TAN 9604). *New Zealand Fisheries Data Report* 77. 51 p.
- Baird, S J (2016) Characterisation analyses for barracouta (*Thyrsites atun*) in BAR 1, 1989–90 to 2013–14. *New Zealand Fisheries Assessment Report* 2016/37. 186 p.
- Beentjes, M P; MacGibbon, D J (2013) Review of QMS species for inclusion in the east coast South Island winter trawl survey reports. *New Zealand Fisheries Assessment Report* 2013/35. 102 p.
- Beentjes, M P; MacGibbon, D; Lyon, W S (2015) Inshore trawl survey of Canterbury Bight and Pegasus Bay, April–June 2014 (KAH1402). *New Zealand Fisheries Assessment Report* 2015/14. 136 p.
- Beentjes, M P; MacGibbon, D; Parkinson, D (2016) Inshore trawl survey of Canterbury Bight and Pegasus Bay, April–June 2016 (KAH1605). *New Zealand Fisheries Assessment Report* 2016/61. 135 p.
- Beentjes, M P; Stevenson, M L (2000) Review of the east coast South Island winter trawl survey time series, 1991–96. *NIWA Technical Report* 86. 64 p.
- Beentjes, M P; Stevenson, M L (2001) Review of east coast South Island summer trawl survey time series, 1996–97 to 1999–2000. *NIWA Technical Report* 108. 92 p.
- Boyd, R O; Reilly, J L (2002) 1999/2000 national marine recreational fishing survey: harvest estimates. Draft New Zealand Fisheries Assessment Report. (Unpublished report held by FNZ, Wellington.)
- Bradford, E (1998) Harvest estimates from the 1996 national recreational fishing surveys. New Zealand Fisheries Assessment Research Document 1998/16 27 p. (Unpublished report held by NIWA library, Wellington.)
- Drummond, K L; Stevenson, M L (1995) Inshore trawl survey of the west coast South Island and Tasman and Golden Bays, March–April 1992 (KAH9204). *New Zealand Fisheries Data Report* 63. 58 p.

BARRACOUTA (BAR)

- Drummond, K L; Stevenson, M L (1995) Inshore trawl survey of the west coast South Island and Tasman and Golden Bays, March–April 1994 (KAH9404). *New Zealand Fisheries Data Report* 64. 55 p.
- Drummond, K L; Stevenson, M L (1996) Inshore trawl survey of the west coast South Island and Tasman and Golden Bays, March–April 1994 (KAH9504). *New Zealand Fisheries Data Report* 74. 60 p.
- Francis, R I C; Hurst, R J; Renwick, J A (2001) An evaluation of catchability assumptions in New Zealand stock assessments. *New Zealand Fisheries Assessment Report* 2001/1. 37 p.
- Francis, M P; Paul, L J (2013) New Zealand inshore finfish and shellfish commercial landings, 1931–82. *New Zealand Fisheries Assessment Report* 2013/55. 136 p.
- Grant, C J; Cowper, T R; Reid, D D (1978) Age and growth of snoek, *Leionura atun* (Euphrasen) in South-eastern Australian waters. *Australian Journal of Marine and Freshwater Research* 29: 435–444.
- Harley, S J; Horn, P L; Hurst, R J; Bagley, N W (1999) Analysis of commercial catch and effort data and age determination and catch-at-age of barracouta in BAR 5. New Zealand Fisheries Assessment Research Document 1999/39. 39 p. (Unpublished document held by NIWA library, Wellington.)
- Hatanaka, H; Uozumi, Y; Fukui, J; Aizawa, M; Hurst, R J (1989) Japan New Zealand trawl survey off southern New Zealand, October–November 1983. *New Zealand Fisheries Technical Report* 9. 52 p.
- Horn, P L (2002) Age estimation of barracouta (*Thyrsites atun*) off southern New Zealand. *Marine and Freshwater Research* 53: 1169–1178.
- Hurst, R J (1988a) The barracouta, *Thyrsites atun*, fishery around New Zealand: historical trends to 1984. *New Zealand Fisheries Technical Report* 5. 43 p.
- Hurst, R J (1988b) Barracouta. New Zealand Fisheries Assessment Research Document 1988/8. 29p. (Unpublished document held by NIWA library, Wellington.)
- Hurst, R J; Bagley, N W (1987) Results of a trawl survey of barracouta and associated finfish near the Chatham Islands, New Zealand, December 1984. *New Zealand Fisheries Technical Report* 3. 44 p.
- Hurst, R J; Bagley, N W (1989) Movements and possible stock relationships of the New Zealand barracouta, *Thyrsites atun*, from tag returns. *New Zealand Journal of Marine and Freshwater Research* 23: 105–111.
- Hurst, R J; Bagley, N W (1992) Results of a trawl survey of barracouta and associated finfish near the Chatham Islands, New Zealand, December 1985. *New Zealand Fisheries Technical Report* 30. 36 p.
- Hurst, R J; Bagley, N W (1997) Trends in Southland trawl surveys of inshore and middle depth species 1993–1996. *New Zealand Fisheries Technical Report No.50*. 66p.
- Hurst, R J; Bagley, N W (1994) Trawl survey of middle depth and inshore bottom species off Southland, February–March 1993 (TAN 9301). *New Zealand Fisheries Data Report* 52. 58 p.
- Hurst, R J; Bagley, N W; Uozumi, Y (1990) New Zealand - Japan trawl survey of shelf and upper slope species off southern New Zealand, June 1986. *New Zealand Fisheries Technical Report* 18. 50 p.
- Hurst, R J; Ballara, S L; MacGibbon, D (2012) Fishery characterisation and standardised CPUE analyse for barracouta, *Thyrsites atun* (Euphrasen, 1791) (Gempylidae), 1989–90 to 2007–08. *New Zealand Fisheries Assessment Report* 2012/13. 303 p.
- Hurst, R J; Fenaughty, J M (1985) Report on biomass surveys 1980–84: summaries and additional information. Fisheries Research Division Internal Report 21: 53 p. (Unpublished report held by NIWA library, Wellington.)
- Langley, A D; Bentley, N (2002) An examination of the stock structure of barracouta (*Thyrsites atun*) around the South Island of New Zealand. *New Zealand Fisheries Assessment Report* 2002/30. 57 p.
- Langley, A D; Walker, N A (2002) Characterisation of the barracouta (*Thyrsites atun*) fishery in BAR 1. *New Zealand Fisheries Assessment Report* 2002/44. 37 p.
- Langley, A D; Walker, N (2002) CPUE analysis of the southeast South Island BAR 1 fishery, 1989–90 to 1999–2000. *New Zealand Fisheries Assessment Report* 2002/11. 28 p.
- Marsh, C. (in prep.) Fishery characterisation and standardised CPUE analyses for barracouta *Thyrsites atun*, (Euphrasen, 1791) (Gempylidae), in BAR 5 1989 to 2013–14 (Draft New Zealand Fisheries Assessment Report submitted to FNZ.)
- McGregor, V (in prep.) Fishery characterisation and standardised CPUE analyses for barracouta *Thyrsites atun*, (Euphrasen, 1791) (Gempylidae), 1989 to 2010–11 New Zealand Fisheries Assessment Report (draft)
- Stevenson, M L (1996) Bottom trawl survey of the inshore waters of the east coast North Island, February–March 1995 (KAH9502). *New Zealand Fisheries Data Report* 78. 57 p.
- Stevenson, M L (1998) Inshore trawl survey of the west coast South Island and Tasman and Golden Bays, March–April 1997 (KAH9701). *NIWA Technical Report* 12. 70 p.
- Teimey, L D; Kilner, A R; Millar, R E; Bradford, E; Bell, J D (1997) Estimation of recreational catch from 1991/92 to 1993/94. New Zealand Fisheries Assessment Research Document 1997/15. 43 p. (Unpublished document held by NIWA library, Wellington.)
- Wynne-Jones, J; Gray, A; Hill, L; Heinemann, A (2014) National Panel Survey of Marine Recreational Fishers 2011–12: Harvest Estimates. *New Zealand Fisheries Assessment Report* 2014/67.

BLACK CARDINALFISH (CDL)

(*Epigonus telescopus*)
Akiwa

**1. FISHERY SUMMARY**

Black cardinalfish was introduced into the QMS on 1 October 1998 and quotas were set for QMAs 2–8. Quotas for QMAs 1 and 9 were subsequently set for 1999–00. TACCs were increased from 1 October 2006 in CDL 4 to 66 t and in CDL 5 to 22 t. In these stocks landings were above the TACC for a number of years and the TACCs were increased to the average of the previous eight years plus an additional 10%. From 1 October 2009 the TACC was reduced in CDL 2 to 1620 t, then reduced to 1020 t in 2010–11, and further reduced to 440 t in 2011–12. CDL 1 and CDL 2 have other mortality allocations of 120 t and 20 t respectively. (Table 1).

Table 1: TACs (t), TACCs (t) and allowances (t) for black cardinal fish.

Fishstock	Recreational Allowance	Customary non-commercial Allowance	Other sources of mortality	TACC	TAC
CDL 1	0	0	120	1200	1320
CDL 2	0	0	20	440	460
CDL 3	0	0	-	196	196
CDL 4	0	0	-	66	66
CDL 5	0	-	79	22	22
CDL 6	0	0	-	1	1
CDL 7	0	0	-	39	39
CDL 8	0	0	-	0	0
CDL 9	0	0	-	4	4
CDL 10	0	0	-	0	0
Total	0	0	219	1 968	2 108

1.1 Commercial fisheries

Several species of *Epigonus* are widely distributed in New Zealand waters, but only black cardinalfish (*E. telescopus*) reaches a marketable size and is found in commercial concentrations. It occurs throughout the New Zealand EEZ at depths of 300–1100 m, mostly in very mobile schools up to 150 m off the bottom over hills and rough ground. Black cardinalfish have been caught since 1981 by research and commercial vessels, initially as a bycatch of target trawling for other high value species. The preferred depth range of schools (600–900 m) overlaps the upper end of the depth range of orange roughy and the lower end of alfonsino and bluenose. The exploitation of these species from 1986 resulted in the development of the major cardinalfish fishery in QMA 2.

BLACK CARDINALFISH (CDL)

It is primarily sold domestically due to the short freezer life of fillets. The species has a section of dark flesh under the lateral line that has caused problems with overseas marketing. The fillets can be tainted if this flesh is not removed quickly.

Landings for 1998–99 to 2008–09 are from QMR totals following introduction of the species into the QMS for 1998–99. For the 1982–83 to 1985–86 fishing years, the best estimate of landings was the sum of the FSU Inshore and FSU Deepwater (i.e., FSU Total) catch returns. For 1986–87 to 1988–89 the best estimate was taken as the greater value of either the FSU Total or the LFRR. From the 1989–90 fishing year, the best estimate was taken as the higher of either the LFRR or the sum of the CLR and CELR Landed data.

The best estimate of total landings was split between the nine QMAs and ET (outside the EEZ) based on FSU and QMS data (Table 2). For FSU data (1982–83 to 1987–88 fishing years), catch where area was unknown was pro-rated to QMAs according to the catch level where area was reported. For QMS data (1988–89 to 1994–95 fishing years), catch by area in CELR Landed and CLR reports were scaled to equal the best estimate of the total catch. Commercial landings of black cardinalfish have been made in QMAs 1–9 and outside the EEZ (ET).

In most years since 1982 more than 65% of black cardinalfish landings were from the east coast of the North Island (QMA 2). The large increase in landings from this area in 1986–87 was associated with the development of the orange roughy fishery around the Ritchie Banks and Tuaheni High, and an increase in targeted fishing to establish a catch history when it was anticipated to become a quota species. Landings from the Bay of Plenty (QMA 1) have fluctuated since 1988. The relatively large landings in 1990–91 were a combination of bycatch of the orange roughy fishery and target fishing for black cardinalfish. Between 1991–92 and 2005–06 occasional large catches were taken from outside the EEZ on the northern Challenger Plateau and the Lord Howe Rise.

Figure 1 shows the historical landings and TACC values for the main CDL stocks.

1.2 Recreational fisheries

Recreational fishing for black cardinalfish is negligible.

1.3 Customary non-commercial fisheries

The level of this fishery is believed to be negligible.

1.4 Illegal catch

No information is available about illegal catch.

Table 2: Reported landings (t) of black cardinalfish by QMA and fishing year (1 October to 30 September) from 1982–83 to 2016–17. The data in this table has been updated from that published in the 1998 Plenary Report by using the data through 1996–97 in table 32 on p. 262 of the “Review of Sustainability Measures and Other Management Controls for the 1998–99 Fishing Year - Final Advice Paper” dated 6 August 1998. Data for 1997–98 based on catch and effort returns, since 1998–99 on QMR records.

Year	QMA 1		QMA 2		QMA 3		QMA 4		QMA 5		QMA 6	
	Catch	TACC	Catch	TACC	Catch	TACC	Catch	TACC	Catch	TACC	Catch	TACC
1982–83	-	-	76	-	<1	-	<1	-	-	-	-	-
1983–84	-	-	212	-	7	-	<1	-	-	-	-	-
1984–85	<1	-	189	-	341	-	<1	-	-	-	-	-
1985–86	<1	-	238	-	50	-	3	-	2	-	-	-
1986–87	1	-	1 738	-	72	-	2	-	<1	-	<1	-
1987–88	3	-	1 556	-	28	-	1	-	3	-	-	-
1988–89	305	-	1 434	-	57	-	4	-	-	-	-	-
1989–90	613	-	1 718	-	20	-	18	-	-	-	-	-
1990–91	233	-	3 473	-	598	-	1	-	4	-	-	-
1991–92	7	-	1 652	-	146	-	3	-	<1	-	2	-
1992–93	23	-	1 550	-	519	-	2	-	<1	-	-	-
1993–94	364	-	2 310	-	277	-	10	-	5	-	-	-
1994–95	1 162	-	2 207	-	51	-	7	-	1	-	<1	-
1995–96	1 418	-	2 621	-	57	-	4	-	10	-	-	-

BLACK CARDINALFISH (CDL)

Table 2: [Continued]

Year	QMA 1		QMA 2		QMA 3		QMA 4		QMA 5		QMA 6	
	Catch	TACC	Catch	TACC	Catch	TACC	Catch	TACC	Catch	TACC	Catch	TACC
1996-97	2 001	-	1 910	-	100	-	7	-	-	-	-	-
1997-98	995	-	1 176	-	40	-	351	-	-	-	-	-
1998-99	24	1 200	1 268	2 223	181	196	41	5	-	2	< 1	1
1999-00	980	1 200	2 158	2 223	215	196	36	5	<1	2	< 1	1
2000-01	294	1 200	1 135	2 223	99	196	35	5	74	2	< 1	1
2001-02	455	1 200	1 693	2 223	146	196	29	5	18	2	< 1	1
2002-03	583	1 200	1 845	2 223	172	196	80	5	9	2	< 1	1
2003-04	481	1 200	966	2 223	96	196	148	5	27	2	< 1	1
2004-05	267	1 200	1 102	2 223	43	196	49	5	15	2	< 1	1
2005-06	643	1 200	2 153	2 223	50	196	53	5	< 1	2	< 1	1
2006-07	415	1 200	1 692	2 223	66	196	31	66	10	22	< 1	1
2007-08	202	1 200	861	2 223	7	196	23	66	20	22	<1	1
2008-09	197	1 200	1 135	2 223	52	196	58	66	11	22	< 1	1
2009-10	49	1 200	1 046	1 620	45	196	15	66	3	22	< 1	1
2010-11	84	1 200	736	1 020	17	196	19	66	5	22	< 1	1
2011-12	148	1 200	376	440	79	196	44	66	93	22	< 1	1
2012-13	35	1 200	470	440	40	196	10	66	14	22	1	1
2013-14	160	1 200	282	440	68	196	11	66	19	22	<1	1
2014-15	21	1 200	408	440	209	196	18	66	4	22	<1	1
2015-16	35	1 200	299	440	136	196	30	66	15	22	1	1
2016-17	12	1 200	369	440	101	196	22	66	87	22	2	1
Year	QMA 7		QMA 8		QMA 9		Total (EEZ)		ET		Total	
	Catch	TACC	Catch	TACC	Catch	TACC	Catch	TACC	Catch		Catch	
1982-83	<1	-	-	-	-	-	78	-	-		78	
1983-84	<1	-	-	-	-	-	220	-	-		220	
1984-85	1	-	-	-	-	-	532	-	-		532	
1985-86	<1	-	-	-	45	-	292	-	-		292	
1986-87	<1	-	-	-	-	-	1 814	-	-		1 814	
1987-88	2	-	<1	-	<1	-	1 638	-	-		1 638	
1988-89	2	-	-	-	-	-	1 798	-	2		1 800	
1989-90	15	-	-	-	-	-	2 385	-	<1		2 385	
1990-91	1	-	<1	-	-	-	4 311	-	-		4 311	
1991-92	11	-	-	-	-	-	1 821	-	17		1 838	
1992-93	2	-	-	-	-	-	2 096	-	270		2 366	
1993-94	6	-	-	-	-	-	2 972	-	829		3 801	
1994-95	51	-	-	-	<1	-	3 479	-	231		3 710	
1995-96	26	-	-	-	-	-	4 150	-	340		4 490	
1996-97	27	-	-	-	-	-	4 045	-	522		4 567	
1997-98	76	-	-	-	108	-	2 338	-	405		2 743	
1998-99	16	39	<1	0	< 1	4	1 531	3 670	390		1 921	
1999-00	27	39	0	0	< 1	4	3 415	3 670	962		4 377	
2000-01	2	39	0	0	3	4	1 642	3 670	571		2 213	
2001-02	3	39	0	0	5	4	2 349	3 670	490		2 839	
2002-03	27	39	0	0	5	4	2 721	3 670	275		2 996	
2003-04	2	39	0	0	6	4	1 727	3 670	58		1 785	
2004-05	2	39	0	0	1	4	1 479	3 670	204		1 683	
2005-06	1	39	0	0	2	4	2 901	3 670	44		2 945	
2006-07	1	39	0	0	1	4	2 216	3 751	2		2 218	
2007-08	2	39	<1	0	19	4	1 134	3 751	1		1 135	
2008-09	1	39	0	0	2	4	1 456	3 751	17		1 474	
2009-10	< 1	39	0	0	5	4	1 163	3 148	-		-	
2010-11	< 1	39	0	0	1	4	863	2 548	-		-	
2011-12	< 1	39	0	0	<1	4	742	1 968	-		-	
2012-13	2	39	0	0	4	4	576	1 968	-		-	
2013-14	1	39	0	0	<1	4	542	1 968	-		-	
2014-15	5	39	0	0	1	4	665	1 968	-		-	
2015-16	3	39	0	0	2	4	522	1 968	-		-	
2016-17	5	39	0	0	1	4	599	1 968	-		-	

BLACK CARDINALFISH (CDL)




Figure 1: Reported commercial landings and TACC for the two main CDL stocks. CDL 1 (Auckland East) and CDL 2 (Central East).

1.5 Other sources of mortality

There has been a history of catch overruns (unreported catch) from loss of fish through burst nets, and the discarding at sea of this species while target fishing for higher value species. In the assessment presented here, the total removals were assumed to exceed reported catches by the overrun percentages in Table 3 (Dunn 2009). All yield estimates make an allowance for the current estimated level of overrun of 10%.

Table 3: Catch overruns (%) for CDL 2 by year.

Year	Over-run	Year	Over-run
1982–83	100	1991–92	30
1983–84	100	1992–93	30
1984–85	100	1993–94	30
1985–86	100	1994–95	20
1986–87	50	1995–96	20
1987–88	50	1996–97	20
1988–89	50	1997–98	20
1989–90	50	1998–99 and subsequently	10
1990–01	50		-

2. BIOLOGY

The average size of black cardinalfish landed by the commercial fishery is about 50–60 cm fork length (FL). Length frequency distributions from research surveys are unimodal with a peak at 55–65 cm FL. They reach a maximum length of about 75 cm FL. Otolith readings from 722 fish from QMA 2 have been validated using radiometric and bomb radiocarbon methods, and indicated that this species is relatively slow-growing and long lived (Andrews & Tracey 2007, Neil et al 2008).

Maximum ages of over 100 years were reported, with the bulk of the commercial catch being between 35 and 55 years of age. The validation indicated that fish aged over 60 years tended to be under-aged, by up to 30%. This bias would be likely to have little impact on the estimated growth parameters, but would influence the estimate of natural mortality (M). Life history parameters are given below in Table 4.

Table 4: Life history parameters for black cardinalfish. All estimates are for CDL 2, except the length-weight parameters which are for CDL 2–4.

Fishstock	Estimate	Source
1. Natural mortality (M)	0.034*	(Tracey et al 2000)
Age at recruitment (A_r)	unknown	
Gradual recruitment (A_m)	unknown	
Age at full recruitment	45	(Tracey et al 2000)
Age at maturity (A_s)	35	(Field & Clark 2001)
Gradual maturity (S_m)	13	(Field & Clark 2001)
2. Weight = $a(\text{length})^b$ (weight in g, fork length in cm).		
	Both sexes	
	a	b
	0.113	2.528
		Dunn (2009)
3. Von Bertalanffy growth parameters		
	Both sexes	
	Female	
	Male	
	L_∞	k
	t_0	t_0
	70.8	0.034
	-6.32	-4.62
	70.9	0.038
	67.8	0.034
	-8.39	-8.39

* Because of uncertainties in ageing and M , the Deepwater Fisheries Assessment Working Group used a range of M 's in the assessments.

The reproductive biology of black cardinalfish is not well known (Dunn 2009). Indications from research survey and Observer Programme data are that spawning may occur between November and July. Spawning locations have been identified in CDL 1, CDL 2, CDL 7, CDL 9, and outside the EEZ on the northern Challenger Plateau, Lord Howe Rise, and West Norfolk Ridge. A probit analysis of maturity at length indicated that fish became sexually mature at around 50 cm length, at an age of approximately 35 years (Field & Clark 2001). Maturity was also inferred to be between ages 26 and 44 years (mean 33 years) from changes in $\delta^{13}\text{C}$ in otoliths (Neil et al 2008).

Juveniles are thought to be mesopelagic until they reach a length of about 12 cm (5 years of age), after which they become primarily demersal (Neil et al 2008). Larger juveniles have been caught in bottom trawls at depths of 400–700 m, extending into deeper water as they grow, with adult fish caught primarily at 800–1000 m (Dunn 2009). Prey items from research trawl samples include mesopelagic fish, natant decapod prawns and octopus.

Elevated levels of mercury (Hg) have been recorded in a sample of black cardinalfish from the Bay of Plenty (Tracey 1993).

3. STOCKS AND AREAS

The stock boundaries and number of black cardinalfish stocks in New Zealand are unknown. There are no data on genetics, or known movements of black cardinalfish which indicate possible stock boundaries.

There is evidence that spawning occurs in CDL 1, CDL 2, CDL 7 and CDL 9 and outside the EEZ (e.g., North Challenger, Lord Howe and West Norfolk Ridge). In CDL 2, three geographically close spawning locations have been identified: Tuaheni High, Ritchie Bank, and Rockgarden (Dunn 2009). Juveniles of less than 30 cm have been infrequently identified in CDL 2, and more frequently found on the northern flanks of the Chatham Rise, which is south of the spawning grounds in CDL 2. No spawning grounds have been identified on the Chatham Rise, where adult fish are relatively rare.

4. ENVIRONMENTAL AND ECOSYSTEM CONSIDERATIONS

This section was updated for the 2018 Fishery Assessment Plenary. A more detailed summary from an issue-by-issue perspective is available in the 2017 Aquatic Environment & Biodiversity Annual Review (MPI 2017, <https://www.mpi.govt.nz/dmsdocument/27471-aquatic-environment-and-biodiversity-annual-review-aebar-2017-a-summary-of-environmental-interactions-between-the-seafood-sector-and-the-aquatic-environment>).

4.1 Role in the ecosystem

Black cardinalfish is a part of the mid slope demersal fish assemblage identified by Francis et al (2002). It is widely distributed with a range centred on a depth of about 750 m and latitude about 39.4° S (i.e., central and northern New Zealand). It occupies depths intermediate between the shallower southern community dominated by hoki (about 620 m, 49.5° S) and the deeper southern black oreo (about 930 m, 45.5° S) and smooth oreo (about 1090 m, 44.6° S), and the deeper centrally-located orange roughy (about 1090 m, 41.2° S) (Francis et al 2002). The role in the ecosystem is not well understood; and nor are the effects on the ecosystem of removing about an average of 2300 t of black cardinalfish per year between 1986–87 and 2010–11 from the New Zealand EEZ, mostly from the east coast of the North Island.

4.1.1 Trophic interactions

No detailed feeding studies for black cardinalfish have been documented for New Zealand waters. Prey items observed during research surveys in New Zealand waters include mesopelagic fish, particularly lighthouse fish (*Phosichthys argenteus*), natant decapod prawns, and cephalopods (Tracey 1993). Predators of black cardinalfish are not documented but predation is expected to vary with fish development.

4.1.2 Ecosystem Indicators

Tuck et al (2009) used data from the Sub-Antarctic and Chatham Rise middle-depth trawl surveys to derive indicators of fish diversity, size, and trophic level. However, fishing for cardinalfish occurs mostly deeper than the depth range of these surveys and is only a small component of fishing in the areas considered by Tuck et al (2009).

4.2 Bycatch (fish and invertebrates)

Incidental catch and discards have not been estimated for the black cardinalfish target fishery. Anderson (2009, 2011) summarised the bycatch and discards from the target orange roughy and oreo trawl fisheries from 1999–2000 to 2004–05 and 2005–06 to 2008–09 respectively. The bycatch of these fisheries may be similar to that of the cardinalfish fishery, although both occur somewhat deeper than cardinalfish and oreo fisheries are found further to the south.

4.3 Incidental Capture of Protected Species (seabirds, mammals, and protected fish)

For protected species, capture estimates presented here include all animals recovered to the deck (alive, injured or dead) of fishing vessels but do not include any cryptic mortality (e.g., seabirds struck by a warp but not brought onboard the vessel, Middleton & Abraham 2007).

4.3.1 Seabird interactions

Annual observed seabird capture rates ranged from 0 to 0.9 per 100 tows in orange roughy, oreo, and cardinalfish trawl fisheries between 2002–03 and 2015–16 (Baird 2001, 2004 a, b, 2005, Abraham & Thompson 2009, Abraham et al 2009, Abraham & Thompson 2011, Abraham et al 2016). Capture rates have fluctuated without obvious trend at this low level (Table 5). In the 2015–16 fishing year there were 4 observed captures of seabirds and 2 observed captures of seabirds in 2016–17 in orange roughy, oreo, and cardinalfish trawl fisheries at a rates of 0.3 and 0.2 (respectively) seabirds per 100 observed tows (Table 5, Abraham et al 2016). The average capture rate in deepwater trawl fisheries (including orange roughy, oreo and cardinalfish) for the period from 2002–03 to 2016–17 is about 0.29 birds per 100 tows, a very low rate relative to other New Zealand trawl fisheries, e.g. for scampi (4.43 birds per 100 tows) and squid (13.79 birds per 100 tows) over the same years.

Table 5: Number of tows by fishing year and observed seabird captures in orange roughy, oreo, and cardinalfish trawl fisheries, 2002–03 to 2016–17. No. obs, number of observed tows; % obs, percentage of tows observed; Rate, number of captures per 100 observed tows. Estimates are based on methods described in Abraham et al (2016) and Abraham & Richard (2017, 2018) and available via <https://data.dragonfly.co.nz/psc>. Estimates from 2002–03 to 2015–16 are based on data version 2017v1.

	Fishing effort			Observed captures		Estimated captures	
	Tows	No. obs	% obs	Captures	Rate	Mean	95% c.i.
2002–03	8 873	1 383	15.6	0	0	29	15–48
2003–04	8 007	1 262	15.8	3	0.2	28	16–45
2004–05	8 417	1 619	19.2	7	0.4	48	29–74
2005–06	8 292	1 361	16.4	8	0.6	34	21–51
2006–07	7 371	2 326	31.5	1	0	17	8–28
2007–08	6 731	2 811	41.8	7	0.2	19	12–29
2008–09	6 132	2 374	38.7	7	0.3	20	12–30
2009–10	6 013	2 135	35.6	19	0.9	36	27–49
2010–11	4 179	1 206	28.8	2	0.2	15	6–27
2011–12	3 655	923	25.3	2	0.2	11	5–19
2012–13	3 098	345	11.2	2	0.6	13	6–23
2013–14	3 606	435	12.0	2	0.5	14	6–24
2014–15	3 784	961	25.8	0	0	13	5–24
2015–16	4 085	1 367	33.5	4	0.3	12	6–20
2016–17	3 971	1 226	30.9	2	0.2		

Table 6: Number of observed seabird captures in orange roughy, oreo, and cardinalfish fisheries, 2002–03 to 2016–17, by species and area. The risk category is an estimate of aggregate potential fatalities across trawl and longline fisheries relative to the Population Sustainability Thresholds, PST (from Richard et al 2017, where full details of the risk assessment approach can be found). It is not an estimate of the risk posed by fishing for cardinal fish. These data are available via <https://data.dragonfly.co.nz/psc>, based on data version 2017v1.

Species	Risk Category	Chatham Rise	East Coast South Island	Fiordland	Sub-Antarctic	Stewart Snares Shelf	West Coast South Island	Total
Salvin's albatross	High	13	4	0	3	0	0	20
Southern Buller's albatross	High	3	0	1	0	0	0	4
Chatham Island albatross	High	7	0	0	1	0	0	8
New Zealand white-capped albatross	High	3	0	0	0	0	1	4
Gibson's albatross	High	1	0	0	0	0	0	1
Antipodean albatross	Medium	1	0	0	0	0	0	1
Northern royal albatross	Low	1	0	0	0	0	0	1
Southern royal albatross	Negligible	1	0	0	0	0	0	1
Total albatrosses	-	30	4	1	4	0	1	40
Northern giant petrel	Medium	1	0	0	0	0	0	1
White-chinned petrel	Negligible	2	1	0	0	0	0	3
Grey petrel	Negligible	1	0	0	1	0	0	2
Sooty shearwater	Negligible	0	3	0	0	0	0	3
Common diving petrel	Negligible	2	0	0	0	0	0	2
White-faced storm petrels	Negligible	3	0	0	0	0	0	3
Cape petrel	-	8	1	0	0	0	0	9
Short-tailed shearwater	-	0	0	0	0	1	0	1
Petrels, prions and shearwaters	-	0	0	0	1	0	0	1
Total other birds	-	17	5	0	2	1	0	25

Salvin's albatross was the most frequently captured albatross (50% of observed albatross captures) but eight different species have been observed captured since 2002–03. Cape petrels were the most frequently captured other taxon (36% of observed captures of taxa other than albatross, Table 6). Seabird captures in the orange roughy, oreo, and cardinalfish fisheries have been observed mostly around the Chatham Rise and off the east coast South Island. These numbers should be regarded as only a general guide on the distribution of captures because the observer coverage is not uniform across areas and may not be representative.

BLACK CARDINALFISH (CDL)

The deepwater trawl fisheries (including the cardinal fish target fishery) contributes to the total risk posed by New Zealand commercial fishing to seabirds (see Table 7). The two species to which the fishery poses the most risk are Chatham Island albatross and Salvin's albatross, with this suite of fisheries posing 0.06 and 0.022 of Population Sustainability Threshold (PST) (Table 7). Chatham albatross and Salvin's albatross were assessed as at high risk (Richard et al 2017).

Mitigation methods such as streamer (tori) lines, Brady bird bafflers, warp deflectors, and offal management are used in the orange roughy, oreo, and cardinalfish trawl fisheries. Warp mitigation was voluntarily introduced from about 2004 and made mandatory in April 2006 (Department of Internal Affairs 2006). The 2006 notice mandated that all trawlers over 28 m in length use a seabird scaring device while trawling (being "paired streamer lines", "bird baffle" or "warp deflector" as defined in the notice).

Table 7: Risk ratio of seabirds predicted by the level two risk assessment for the cardinalfish and all fisheries included in the level two risk assessment, 2006–07 to 2015–17, showing seabird species with a risk ratio of at least 0.001 of PST (from Richard et al 2017, where full details of the risk assessment approach can be found). The risk ratio is an estimate of aggregate potential fatalities across trawl and longline fisheries relative to the PST. The DOC threat classifications are shown (Robertson et al 2017 at <http://www.doc.govt.nz/documents/science-and-technical/nztc19entire.pdf>).

Species name	PST (mean)	Risk ratio		Risk category	DOC Threat Classification
		DPW Risk Ratio	TOTAL		
Chatham Island albatross	425.2	0.0602	0.362	High	At Risk: Naturally Uncommon
Salvin's albatross	3 599.5	0.0223	0.78	High	Threatened: Nationally Critical
Northern giant petrel	335.4	0.0052	0.138	Medium	At Risk: Naturally Uncommon
Northern Buller's albatross	1 627.4	0.0024	0.253	Medium	At Risk: Naturally Uncommon
Black petrel	437.1	0.0024	1.153	Very high	Threatened: Nationally Vulnerable
Antipodean albatross	364.3	0.002	0.203	Medium	Threatened: Nationally Critical
Gibson's albatross	496.1	0.0016	0.337	High	Threatened: Nationally Critical
Northern royal albatross	715.1	0.0013	0.043	Low	At Risk: Naturally Uncommon
Flesh-footed shearwater	1 452.8	7.00E-04	0.669	High	Threatened: Nationally Vulnerable
Southern Buller's albatross	1 368.4	6.00E-04	0.392	High	At Risk: Naturally Uncommon
Grey petrel	5 524.1	3.00E-04	0.037	Negligible	At Risk: Naturally Uncommon
Common diving petrel	135 254.8	1.00E-04	0.002	Negligible	At Risk: Relict
New Zealand white-faced storm petrel	331 778.5	1.00E-04	0	Negligible	At Risk: Relict
New Zealand white-capped albatross	10 900.3	1.00E-04	0.353	High	At Risk: Declining
Buller's shearwater	55 991.9	0	0	Negligible	At Risk: Naturally Uncommon
Westland petrel	350.1	0	0.476	High	At Risk: Naturally Uncommon
Sooty shearwater	617 028.2	0	0.002	Negligible	At Risk: Declining
Hutton's shearwater	15 054.3	0	0.001	Negligible	At Risk: Declining
Otago shag	284.0	0	0.144	Medium	Threatened: Nationally Vulnerable
White-headed petrel	34 314.8	0	0.001	Negligible	Not Threatened

4.4 Benthic interactions

Cardinalfish, orange roughy, and oreos are taken using bottom trawls and collectively accounted for about 14% of all tows reported on TCEPR forms to have been fished on or close to the bottom between 1989–90 and 2004–05 (Baird et al 2011). These tows were located in Benthic Optimised Marine Environment Classification (BOMECE, Leathwick et al 2012) classes J, K (mid-slope), M (mid-lower slope), N, and O (lower slope and deeper waters) (Baird & Wood 2012), and 94% were between 700 and 1200 m depth (Baird et al 2011). Deepsea corals in the New Zealand region are abundant and diverse and, because of their fragility, are at risk from anthropogenic activities such as bottom trawling (Clark & O'Driscoll 2003, Clark & Rowden 2009, Williams et al 2010). All deepwater hard corals are protected under Schedule 7A of the Wildlife Act 1953. Rowden et al (2012) mapped the likely coral distributions using predictive models, and concluded that fisheries that pose the most risk to protected corals are these deepwater trawl fisheries.

Trawling for orange roughy, oreo, and cardinalfish, like trawling for other species, is likely to have effects on benthic community structure and function (e.g., Rice 2006) and there may be consequences for benthic productivity (e.g., Jennings et al 2001, Hermsen et al 2003, Hiddink et al 2006, Reiss et al

2009). These consequences are not considered in detail here but are discussed in the Aquatic Environment and Biodiversity Annual Review 2015.

The NZ EEZ contains 17 Benthic Protection Areas (BPAs) that are closed to bottom trawl fishing and include about 52% of all seamounts greater than 1500 m elevation and 88% of identified hydrothermal vents.

4.5 Other considerations

4.5.1 Spawning disruption

Fishing during spawning may disrupt spawning activity or success. Morgan et al (1999) concluded that Atlantic cod (*Gadus morhua*) “exposed to a chronic stressor are able to spawn successfully, but there appears to be a negative impact of this stress on their reproductive output, particularly through the production of abnormal larvae”. Morgan et al. (1997) also reported that “Following passage of the trawl, a 300-m-wide “hole” in the [cod spawning] aggregation spanned the trawl track. Disturbance was detected for 77 min after passage of the trawl.” There is no research on the disruption of spawning black cardinalfish by fishing in New Zealand. Spawning of this species appears to occur between February and July, peaking in April, and catches of black cardinalfish occur throughout the year (Dunn 2005).

4.5.2 Genetic effects

Fishing, environmental changes, including those caused by climate change or pollution, could alter the genetic composition or diversity of a species. There are no known studies of the genetic diversity of cardinalfish from New Zealand. Genetic studies for stock discrimination are reported under “stocks and areas”.

4.5.3 Habitat of particular significance to fisheries management

Habitat of particular significance for fisheries management (HPSFM) does not have a policy definition (Ministry for Primary Industries 2012). O’Driscoll et al (2003) reported spawning black cardinalfish mostly from around the North Island, but higher catch rates of juveniles on the northwest Chatham Rise and Puysegur area (O’Driscoll et al 2003). In both cases, sample sizes were small so these distributions should be treated with caution. It is not known if there are any direct linkages between the congregation of cardinalfish around features and the corals found on those features. Bottom trawling for cardinalfish has the potential to affect features of the habitat that could qualify as habitat of particular significance to fisheries management.

5. STOCK ASSESSMENT

A stock assessment for CDL 2–4 was completed in 2009. No assessments have been made for stocks in other areas. For the purposes of stock assessment, it has been assumed that black cardinalfish on the east coast North Island (CDL 2) are from the same stock as fish on the north Chatham Rise (CDL 3 and CDL 4).

5.1 Assessment inputs

The assessment inputs for CDL 2–4 were catches adjusted by overruns (Table 9), two CPUE indices (Table 8), and length frequency and maturity at length samples (Dunn 2009). The CPUE indices were derived from catch and effort data for fisheries focused on and around specific hill features in CDL 2 (Dunn & Bian 2009) with no overrun included. Whilst the CPUE indices accounted for a substantial proportion of the total catch (65–77%), the spatial extent of the fisheries was small compared with the overall area believed to be occupied by the stock. As a result, the indices may reflect local abundance, but it is less certain that they reflect overall stock biomass. The CPUE was split into two indices, before and after 1 October 1998, because of a change in reported fishing patterns in the late 1990s. This may have been caused, at least in part, by the introduction of the black cardinalfish TACC. The growth parameters used in the assessment are presented in Table 4. Length frequency samples were available for eight years between 1989–90 and 2007–08 from at-sea and market sampling. Maturity was input as the proportions mature at length from samples collected during research trawl surveys of the east coast North Island in 2001 and 2003.

BLACK CARDINALFISH (CDL)

Table 8: Standardised CPUE indices, and their calculated CVs, as used in the stock assessment.

Fishing year	Index a	CV (%)	Index b	CV (%)
1990–91	1.00	46	-	-
1991–92	0.73	43	-	-
1992–93	0.87	42	-	-
1993–94	0.58	46	-	-
1994–95	0.41	45	-	-
1995–96	0.26	39	-	-
1996–97	0.51	42	-	-
1997–98	0.29	47	-	-
1998–99	-	-	1.00	37
1999–00	-	-	0.57	32
2000–01	-	-	0.39	36
2001–02	-	-	0.50	35
2002–03	-	-	0.30	33
2003–04	-	-	0.26	38
2004–05	-	-	0.23	35
2005–06	-	-	0.34	34
2006–07	-	-	0.27	35
2007–08	-	-	0.17	37

Table 9: Estimated catches calculated by summing the CDL 2–4 catches from Table 2 (column 2), and increasing them by the overrun values in Table 3 (column 3), with the combined TACC for CDL 2–4 (column 4).

Year	Reported catch	Catch including overruns	TACC
1982–83	76	152	-
1983–84	219	438	-
1984–85	530	1 060	-
1985–86	291	582	-
1986–87	1 812	2 718	-
1987–88	1 585	2 378	-
1988–89	1 495	2 243	-
1989–90	1 756	2 634	-
1990–91	4 072	6 108	-
1991–92	1 801	2 341	-
1992–93	2 071	2 692	-
1993–94	2 597	3 376	-
1994–95	2 265	2 718	-
1995–96	2 682	3 218	-
1996–97	2 017	2 420	-
1997–98	1 567	1 880	-
1998–99	1 490	1 639	2 424
1999–00	2 409	2 650	2 424
2000–01	1 269	1 396	2 424
2001–02	1 868	2 055	2 424
2002–03	2 097	2 307	2 424
2003–04	1 210	1 331	2 424
2004–05	1 194	1 313	2 424
2005–06	2 256	2 482	2 424
2006–07	1 789	1 968	2 485
2007–08	891	980	2 485

5.2 Model structure and runs

Stock assessments were performed using the stock assessment program, CASAL (Bull et al 2002) to estimate virgin and current biomass (Dunn 2009). Preliminary model runs were completed using all of the observational data. The key assumptions of the final model runs were:

- The biomass information in the data is primarily contained in the CPUE indices. Therefore, a two-step approach was used to produce the final model runs. In the final runs, selectivity and maturity were fixed at estimates from the preliminary runs and the length frequency and maturity data were not fitted. This ensured that any biomass signal from the length frequency data, potentially caused by errors in estimated growth and selectivity, did not dominate the signal from the CPUE trends.

- Runs where maturity and selectivity were estimated separately resulted in selectivity curves displaced to the right of the maturity ogive for $M = 0.04$ and $M = 0.06$, resulting in a proportion of the spawning stock not being available to the fishery (called “cryptic biomass”). The Deepwater Fisheries Assessment Working Group considered that it was unlikely that there existed mature biomass that was not vulnerable to the fishery, and agreed that the age of vulnerability should be fixed to the age at maturity for the base case and for the case with $M = 0.06$. The WG agreed to present a sensitivity model run using $M = 0.04$ and with separately estimated maturity and selectivity to explore the implications of this scenario.
- For runs assuming an M of 0.027, the selectivity and maturity estimates were similar; therefore the two were estimated separately in final runs.
- The base case with M set at 0.04 and vulnerability set equal to the MCMC median of maturity was considered to be the most credible.

Four model runs are therefore presented, two with selectivity assumed to be the same as maturity and M assumed to be either 0.06 or 0.04, and two with selectivity and maturity fitted as separate ogives and M assumed to be 0.04 or 0.027 (Table 10).

Table 10: Four alternative assumptions to the stock assessment.

Model	M	Selectivity
Base	0.04	Equal to MCMC median maturity
Mat&sel	0.04	Estimated separately
M0.027	0.027	Estimated separately
M0.06	0.06	Equal to MCMC median maturity

The model was fitted using Bayesian estimation, and partitioned the population by age (age-groups used were 1–90, with a plus group). The model assumed a single sex, with growth modelled using the von Bertalanffy Growth formula. The stock was considered to reside in a single area, and have a single maturation episode, with maturation modelled by a logistic ogive which was estimated in preliminary model runs. Selectivity of the fishery was assumed to be equal to maturity, or modelled by a logistic ogive estimated in preliminary model runs. The catch equation used was the instantaneous mortality equation from Bull et al (2002), whereby half the natural mortality was applied, followed by the fishing mortality, then the remaining natural mortality. Deterministic recruitment was assumed. A Bayesian estimation procedure was used with a penalty function included to discourage the model from allowing the stock biomass to drop below a level at which the historical catch could not have been taken. Lognormal errors, with known (sampling error) CVs were assumed for the CPUE. In preliminary model runs, an additional process error was estimated and added to the length frequency distributions. Binomial errors were assumed for the proportions mature at length. The final model runs estimated virgin biomass, B_0 , and two catchabilities. Confidence intervals were calculated from a posterior distribution of the model parameters, which was estimated using a Markov Chain Monte Carlo technique.

5.3 Biomass estimates

Biomass estimates depended on the assumed M , with the M0.027 run resulting in a larger and less productive stock, and the M0.06 run in a smaller and more productive stock (Table 11, Figure 2). Estimates of current biomass were lowest in the base case.

The mat&sel run estimated cryptic spawning stock biomass, where vulnerability to the fishery took place after maturity, such that a median of 86% and 62% of the mature biomass was vulnerable to the fishery at virgin and 2009 biomass levels, respectively. It is unclear whether cryptic biomass could occur for black cardinalfish, and it is possible that this result is an artefact generated from the model assumptions. Cryptic biomass was not estimated when maturity and selectivity were estimated separately and M was assumed to be 0.027, and in sensitivity runs the level of cryptic biomass was found to increase as M increased. The wide confidence intervals reflect the uncertainty in the model, which was fitted to only relative biomass indices having relatively high CVs (Table 8).

Table 11: Biomass estimates (medians rounded to the nearest 100 t, with 95% confidence intervals in parentheses) for the four model runs. $B_{current}$ is the mid-year biomass in 2009. $p(B_{2009} < 0.1 B_0)$ is the probability of the mature biomass in 2009 being less than 10% of the virgin mature biomass (B_0). $p(B_{2009} < 0.2 B_0)$ is the probability of the mature biomass in 2009 being less than 20% of the virgin mature biomass (B_0).

Run	B_0 (t)	$B_{current}$ (t)	% B_0	$p(B_{2009} < 0.1 B_0)$	$p(B_{2009} < 0.2 B_0)$
Base	36 800 (32 800–95 400)	4 400 (1 900–60 400)	11.9 (5.9–63.3)	0.41	0.70
Mat&sel	40 800 (35 600–96 700)	7 300 (3 500–61 300)	17.8 (9.9–63.5)	0.13	0.56
$M0.027$	45 100 (39 500–93 500)	6 100 (2 000–53 000)	13.6 (5.0–56.6)	0.32	0.69
$M0.06$	33 800 (25 500–10 700)	8 200 (2 400–82 800)	24.2 (9.6–74.9)	0.16	0.43




Figure 2: Estimated biomass trajectories (solid line) and 95% confidence intervals (shaded area) for the model runs (a) Base, (b) mat&sel, (c) $M0.027$, (d) $M0.06$. The horizontal broken line indicates 20% B_0 .

5.4 Sensitivity analyses

Several sensitivity analyses were conducted (reported in more detail in Dunn 2009). The assessment was found to be relatively insensitive to the assumed catch over-runs. When over-runs were either assumed to be zero, or were doubled for the period before 1998–99 (before the TACC was introduced), the mature stock in 2009 was estimated to be slightly less depleted compared to the Base case, at 13.5% (5.9–67.0%) B_0 , and 12.2% (5.5–58.3%) B_0 , respectively.

5.5 5-year projection results

Forward projections were carried out over a 5 year period using a range of constant catch options. A catch level of 180 t is approximately the level associated with $F = M$, a catch of 890 t is approximately the current (2007–08) catch and a catch of 2490 t is approximately the current (2007–08) TACC. In all projections overrun of 10% was assumed for future catches. For each catch option, three measures of fishery performance were calculated. The first one, % B_0 , is the median biomass in 2009 as a percentage of B_0 . The second one, $P_{0.1}$, is the probability that the biomass at the end of the 5-year period is less than 10% B_0 . The third, $P_{0.2}$, is the probability that the biomass at the end of the 5-year period is less than 20% B_0 . At high future catches the biomass may be reduced to such a low level that the catch is unlikely to be able to be taken (assumed to occur when the exploitation rate exceeds 0.9). This is indicated as P(no catch).

All projections indicate that the biomass would increase for all catch levels near or below the 2008–09 catch (890 t), and would continue to decline at catch levels of 1200 t in all runs except $M = 0.06$, where it would remain about the same (Table 12). In all runs the biomass would decline at catch levels equal to the current TACC (2490 t), and there was a 38–71% probability the biomass would decline to a level where the catch could not be taken.

Table 12: Results from forward projections to 2013 for the model runs. $P_{0.1}$ is the probability of the mature biomass in 2013 being less than 10% of the virgin mature biomass (B_0). $P_{0.2}$ is the probability of the mature biomass in 2013 being less than 20% of the virgin mature biomass (B_0). $P(\text{no catch})$ is the probability that the catch could not be taken, which is assumed to occur if the exploitation rate exceeds 90%. Current (2007–08) values of $\%B_0$ are shown for each run in parenthesis next to the measure. 95% confidence intervals are shown for the $\%B_0$ estimates in 2013. A catch of 180 t is approximately M times the current biomass, 890 t is the current catch and 2490 t is the current TACC.

Run	Measure	Future catch (t)					
		0	180	530	890	1200	2490
Base	$\%B_0$ (11.9)	17.6 (8.5–67.4)	16.5 (7.01–66.0)	14.3 (5.3–63.9)	12.6 (3.6–62.7)	10.2 (2.9–62.6)	5.2 (2.7–56.2)
	$P_{0.1}$	0.11	0.19	0.30	0.40	0.49	0.70
	$P_{0.2}$	0.57	0.60	0.65	0.71	0.74	0.83
	$P(\text{no catch})$	0	0	0	0	0	0.38
mat&sel	$\%B_0$ (17.8)	24.5 (14.0–68.8)	23.6 (12.9–67.8)	20.4 (10.2–65.5)	18.6 (8.0–63.4)	16.2 (6.5–61.7)	9.5 (5.5–57.8)
	$P_{0.1}$	0.00	0.00	0.06	0.14	0.22	0.53
	$P_{0.2}$	0.35	0.38	0.49	0.55	0.61	0.75
	$P(\text{no catch})$	0	0	0	0	0	0.42
M0.027	$\%B_0$ (13.6)	17.9 (7.1–59.4)	16.7 (6.2–59.1)	14.3 (4.5–56.7)	12.0 (2.9–56.5)	10.0 (2.2–55.0)	4.3 (2.0–50.1)
	$P_{0.1}$	0.14	0.19	0.28	0.40	0.49	0.71
	$P_{0.2}$	0.57	0.60	0.67	0.71	0.75	0.84
	$P(\text{no catch})$	0	0	0	0	0	0.41
M0.06	$\%B_0$ (24.2)	33.6 (13.0–80.2)	31.4 (12.5–79.2)	29.8 (10.6–77.5)	26.3 (8.3–77.2)	24.6 (6.7–75.7)	17.4 (4.8–71.2)
	$P_{0.1}$	0.02	0.33	0.07	0.15	0.17	0.35
	$P_{0.2}$	0.27	0.29	0.35	0.40	0.42	0.54
	$P(\text{no catch})$	0	0	0	0	0	0.71

5.6 Updated characterisation and CPUE analyses

A characterisation and CPUE analyses were conducted using catch and effort data to the end of the 2013–14 fishing year (Bentley & MacGibbon, 2016). Catch and effort data were examined in each of nine “zones” which encompassed groups of underwater features where the majority of the cardinalfish catch has been taken: North Colville (NC), Mercury-Colville (MC), White Island (WI), East Cape (EC), Tuaheni High (TH), Richie-Rockgarden (RR), Madden (MD), Wairarapa (WA), and Kaikoura (KK). Within these zones, only tows in the depth range 470–980m (the 2.5th and 97.5th percentiles of the distribution of cardinalfish catch by depth) were considered when characterising effort and performing CPUE analyses.

Catches in each zone have generally declined or remained stable. In CDL 1, most of the catch has come from the Mercury-Colville zone since the early 2000s. In CDL 2, concurrent with a reduction in the TACC, catches have declined in the East Cape, Tuaheni High and Richie-Rockgarden zones since 2010. In these zones, as in CDL 1, most of the cardinalfish is taken in target tows. In contrast, catches in the Wairarapa and Kaikoura zones have remained relatively constant during this period. In these southern two zones a greater proportion of the cardinalfish catch is taken as bycatch from tows that are targeting species other than cardinalfish and orange roughy. There was no evidence of substantial movement of fishing effort between features within zones.

A CPUE analysis was done using data from all nine zones and year effects estimated for each zone. This suggested that the CPUE trends in all zones were generally similar but that the Wairarapa and Kaikoura zones exhibited a flatter trend since 2000. On this basis, a final CPUE standardisation was done with separate year effects estimated for three regions North (zones North Colville, Mercury-Colville and

BLACK CARDINALFISH (CDL)

White Island; i.e. CDL 1), Central (zones East Cape, Tuaheni High, Ritchie-Rockgarden and Madden: i.e. CDL 2 except for Wairarapa) and South (zones Wairarapa and Kaikoura). This standardisation model has the advantage over separate models for each region of using all the available data to estimate vessel coefficients.




Figure 3: CPUE indices by region (see text for definitions of regions). Region/year combinations with less than 30 tows are not shown. Error bars indicate \pm one standard error. Fishing years are indicated by the later calendar year.

6. STATUS OF THE STOCKS

Stock Structure Assumptions

The stock boundaries and number of black cardinalfish stocks in New Zealand is unknown. There are no data on genetics, or known movements of black cardinalfish which indicate possible stock boundaries.


There is evidence that a spawning stock exists in CDL 2, with three geographically close spawning locations identified, on Tuaheni High, Ritchie Bank, and Rockgarden (Dunn 2009). Juveniles of less than 30 cm have been infrequently identified in CDL 2, and more frequently found on the northern flanks of the Chatham Rise, which is south of the spawning grounds in CDL 2. No spawning grounds have been identified on the Chatham Rise, where adult fish are relatively rare.

For the purposes of stock assessment, it has been assumed that black cardinalfish on the east coast North Island (CDL 2) are from the same stock as fish on the north Chatham Rise (CDL 3 and CDL 4).

CDL 2, 3 & 4

Stock Status	
Year of Most Recent Assessment	2009 full assessment 2014 CPUE updated
Assessment Runs Presented	One base case and three sensitivity runs Base case: $M = 0.04$; selectivity equal to maturity Sensitivity runs: various combinations of M and assumptions about the relationship between maturity and selectivity, considered to be less reliable than the base case
Reference Points	Management Target: 40% B_0 Soft Limit: 20% B_0 Hard Limit: 10% B_0 Overfishing threshold: $U_{40\%}$
Status in relation to Target	Very Unlikely (< 10%) to be at or above the target
Status in relation to Limits	<u>Base case:</u> B_{2009} was estimated to be 12% B_0 ; Likely (> 60%) to be below the Soft Limit and About as Likely as Not (40–60%) to be below the Hard Limit. <u>Other model runs:</u> The range of B_{2009} was estimated to be 14–24% B_0 ; About as Likely as Not (40–60%) or Likely (> 60%) to be below the Soft Limit and Unlikely (< 40%) to be below the Hard Limit.
Status in relation to Overfishing	Unknown

Historical Stock Status Trajectory and Current Status



Estimated biomass trajectories (solid line) and 95% confidence intervals (shaded area) for the base case. The horizontal broken line indicates 20% B_0

Fishery and Stock Trends

Recent Trend in Biomass or Proxy	CPUE has been flat since 2008
Recent Trend in Fishing Intensity or Proxy	Unknown
Other Abundance Indices	-
Trends in Other Relevant Indicators or Variables	-

BLACK CARDINALFISH (CDL)

Projections and Prognosis	
Stock Projections or Prognosis	Model projections indicate that the biomass will increase at catch levels near or below the 2007–08 level but will decline sharply at catch levels equal to the TACC.
Probability of Current Catch or TACC causing Biomass to remain below or to decline below Limits	Soft Limit: Likely (> 60%) Hard Limit: About as Likely as Not (40–60%)
Probability of Current Catch or TACC causing Overfishing to continue or to commence	Soft Limit: Likely (> 60%) Hard Limit: Likely (> 60%)

Assessment Methodology and Evaluation		
Assessment Type	2009 Level 1: Full Quantitative Stock Assessment 2014 Level 2: Partial Quantitative Stock Assessment	
Assessment Method	Age-structured CASAL model with Bayesian estimation of posterior distributions	
Assessment Dates	Latest assessment: 2009	Next assessment: Unknown
Overall assessment quality rank	1 – High Quality	
Main data inputs (rank)	- Two commercial catch-per-unit-effort (CPUE) series from the trawl fishery up to 2008 - Estimates of biological parameters	1 – High Quality 1 – High Quality
Data not used (rank)	N/A	
Changes to Model Structure and Assumptions	First accepted assessment for these stocks	
Major sources of Uncertainty	Major sources of uncertainty include the representativeness of the CPUE data, the relationship between CPUE and abundance, the assumption that recruitment has been constant throughout the history of the fishery, estimates of growth and natural mortality and the catch history.	

Qualifying Comments
The TACC was reduced from 2 223 t in 3 stages to the level of 440 t in 2010–11. This level was the maximum annual catch required to rebuild the CDL 2 stock to 30% B_0 within the 24 year period specified in the Harvest Strategy Standard (twice T_{min}). CPUE since 2008 has been flat.

Fishery Interactions
Main associated species are orange roughy, alfonso and, to a lesser extent, hoki. Interactions with other species are currently being characterised.

Other QMAs

There is no information on the status of cardinalfish stocks in other QMAs.

TACCs and reported landings for the 2016–17 fishing year are summarised in Table 13.

Table 13: Summary of TACCs (t) and reported landings (t) for black cardinalfish for the most recent (2016–17) fishing year.


Fishstock	QMA	FMA	2016–17 Actual TACC	2016–17 Reported landings
CDL 1	Auckland (East)	1	1 200	12
CDL 2	Central (East)	2	440	369
CDL 3	South-east (Coast)	3	196	101
CDL 4	South-east (Chatham)	4	66	22
CDL 5	Southland	5	22	87
CDL 6	Sub-Antarctic	6	1	2
CDL 7	Challenger	7	39	5
CDL 8	Central (West)	8	0	0
CDL 9	Auckland (West)	9	4	1
CDL 10	Kermadec	10	0	0
Total			1 968	599

7. FOR FURTHER INFORMATION

- Abraham, E R; Berkenbusch, K; Richard, Y; Thompson, F (2016) Summary of the capture of seabirds, mammals, and turtles in New Zealand commercial fisheries, 2002–03 to 2012–13. *New Zealand Aquatic Environment and Biodiversity Report No.169*. 205 p.
- Abraham, E R; Pierre, J P; Middleton, D A; Cleal, J; Walker, N A; Waugh, S M (2009) Effectiveness of fish waste management strategies in reducing seabird attendance at a trawl vessel. *Fisheries Research*, 95(2), 210–219.
- Abraham, E R; Richard, Y (2017) Summary of the capture of seabirds in New Zealand commercial fisheries, 2002–03 to 2013–14. *New Zealand Aquatic Environment and Biodiversity Report No. 184*. 88 p.
- Abraham, E R; Richard, Y (2018) Estimated capture of seabirds in New Zealand trawl and longline fisheries, 2002–03 to 2014–15. *New Zealand Aquatic Environment and Biodiversity Report No. 197*. 97 p.
- Abraham, E R; Thompson, F N (2011) Estimated Capture of Seabirds in New Zealand Trawl and Longline Fisheries, 2002–03 to 2008–09. *New Zealand Aquatic Environment and Biodiversity Report No. 79*.
- Abraham, E R; Thompson, F N; Berkenbusch, K (2014) Estimated capture of seabirds in New Zealand trawl and longline fisheries, 2002–03 to 2010–11. (Final Research Report for MPI project PRO2010-02 held by FNZ, Wellington.)
- Abraham, E R; Thompson, F N; Oliver, M D (2010) Summary of the capture of seabirds, mammals, and turtles in New Zealand commercial fisheries, 1998–99 to 2007–08. *New Zealand Aquatic Environment and Biodiversity Report No.45*. 149 p.
- Anderson, O F (2009) Fish discards and non-target fish catch in the New Zealand orange roughy trawl fishery: 1999–2000 to 2004–05. *New Zealand Aquatic Environment and Biodiversity Report No.39*. 40 p.
- Anderson, O F (2011) Fish and invertebrate bycatch and discards in orange roughy and oreo fisheries from 1990–91 until 2008–09. *New Zealand Aquatic Environment and Biodiversity Report No. 67*. 61 p.
- Andrews, A H; Tracey, D M (2007) Age validation of orange roughy and black cardinalfish using lead-radium dating. Final Research Report for Ministry of Fisheries Research Project DEE2005-02 Objective 1. 40 p. (Unpublished report held by FNZ, Wellington.)
- Baird, R (2004a) Illegal, unreported and unregulated fishing: an analysis of the legal, economic and historical factors relevant to its development and persistence. *Melbourne Journal of International Law* 5: 299.
- Baird, S J (Ed.). (2001) *Report on the International Fishers' Forum on Solving the Incidental Capture of Seabirds in Longline Fisheries, Auckland, New Zealand, 6-9 November 2000*. Department of Conservation.
- Baird, S J (2004b) Estimation of the incidental capture of seabird and marine mammal species in commercial fisheries in New Zealand waters, 1999–2000. *New Zealand Fisheries Assessment Report 2004/41*. 56 p.
- Baird, S J (2005) Estimation of the incidental capture of seabird species in commercial fisheries in New Zealand waters, 2002–03. *New Zealand Fisheries Assessment Report 2005/2*. 50 p.
- Baird, S J; Wood, B A (2012) Extent of coverage of 15 environmental classes within the New Zealand EEZ by commercial trawling with seafloor contact. *New Zealand Aquatic Environment and Biodiversity Report 89*. 43 p.
- Baird, S J; Wood, B A; Bagley, N W (2011) Nature and extent of commercial fishing effort on or near the seafloor within the New Zealand 200 n. mile Exclusive Economic Zone, 1989–90 to 2004–05. *New Zealand Aquatic Environmental and Biodiversity Report No. 73*. 143 p.
- Baker, C S; Chilvers, B L; Childerhouse, S; Constantine, R; Currey, R; Mattlin, R; van Helden, A; Hitchmough, R; Rolfe, J (2016) Conservation status of New Zealand marine mammals, 2013. *New Zealand Threat Classification Series 14*. Department of Conservation, Wellington. 18 p.
- Baker, C S; Chilvers, L; Constantine, R; DuFresne, S; Mattlin, R H; van Helden, A; Hitchmough, R (2010) Conservation status of New Zealand marine mammals (suborders Cetacea and Pinnipedia), 2009. *New Zealand Journal of Marine and Freshwater Research* 44: 101–115.
- Bentley, N; MacGibbon, D (2016) The fishery for black cardinalfish: characterization and CPUE analyses, 1989–90 to 2013–14. *New Zealand Fisheries Assessment Report 2016/66*.
- Black, J; Wood, R; Berthelsen, T; Tilney, R. (2013) Monitoring New Zealand's trawl footprint for deepwater fisheries: 1989–1990 to 2009–2010. *New Zealand Aquatic Environment and Biodiversity Report No. 110*. 57 p.
- Bull, B.; Francis, R.I.C.C.; Dunn, A.; Gilbert, D.J. (2002). CASAL (C++ algorithmic stock assessment laboratory): CASAL User Manual v1.02.2002/10/21. *NIWA Technical Report 117*. 199 p
- Clark, M R; King, K J (1989) Deepwater fish resources off the North Island, New Zealand: results of a trawl survey, May 1985 to June 1986. *New Zealand Fisheries Technical Report 11*. 55 p.
- Clark, M; O'Driscoll, R (2003) Deepwater fisheries and aspects of their impact on seamount habitat in New Zealand. *Journal of Northwest Atlantic Fishery Science* 31: 441–458.
- Clark, M R; Rowden, A A (2009) Effect of deepwater trawling on the macro-invertebrate assemblages of seamounts on the Chatham Rise, New Zealand. *Deep Sea Research I* 56: 1540–1554.

BLACK CARDINALFISH (CDL)

- Department of Internal Affairs (2006) Seabird Scaring Devices – Circular Issued Under Authority of the Fisheries (Commercial Fishing) Amendment Regulations 2006 (No. F361). *New Zealand Gazette* 6 April 2006: 842–846.
- Dunn, M R (2005) Descriptive analysis of catch and effort data from New Zealand black cardinalfish (*Epigonus telescopus*) fisheries for the fishing years 1979–80 to 2002–03. *New Zealand Fisheries Assessment Report* 2005/32. 47 p.
- Dunn, M R (2007) Analysis of catch and effort data from New Zealand black cardinalfish (*Epigonus telescopus*) fisheries up to the 2004–05 fishing year. *New Zealand Fisheries Assessment Report* 2007/27. 55 p.
- Dunn, M R (2009) Review and stock assessment for black cardinalfish (*Epigonus telescopus*) on the east coast North Island. *New Zealand Fisheries Assessment Report* 2009/39.
- Dunn, M R; Bian, R (2009) Analysis of catch and effort data from New Zealand black cardinalfish (*Epigonus telescopus*) fisheries up to the 2007–08 fishing year. *New Zealand Fisheries Assessment Report* 2009/40. 53 p.
- Field, K D; Clark, M R (2001) Catch-per-unit-effort (CPUE) analysis and stock assessment for black cardinalfish *Epigonus telescopus* in QMA 2. *New Zealand Fisheries Assessment Report* 2001/23. 22 p.
- Field, K D; Tracey, D M; Clark, M R (1997) A summary of information on, and assessment of the fishery for, black cardinalfish, *Epigonus telescopus* (Risso, 1810) (Percoidae: Apogonidae). New Zealand Fisheries Assessment Research Document 97/22. 6 p. (Unpublished report held in NIWA library, Wellington.)
- Francis, M P; Hurst, R J; McArdle, B; Bagley, N W; Anderson, O F (2002) New Zealand demersal fish assemblages. *Environmental Biology of Fishes* 62(2): 215–234.
- Hermesen, J M; Collie, J S; Valentine, P C (2003) Mobile fishing gear reduces benthic megafaunal production on Georges Bank. *Marine Ecology Progress Series* 260: 97–108.
- Hiddink, J G; Jennings, S; Kaiser, M J; Queiros, A M; Duplisea, D E; Piet, G J (2006) Cumulative impacts of seabed trawl disturbance on benthic biomass, production, and species richness in different habitats. *Canadian Journal of Fisheries and Aquatic Sciences* 63: 721–36.
- Jennings, S; Dinmore, T A; Duplisea, D E; Warr, K J; Lancaster, J E (2001) Trawling disturbance can modify benthic production processes. *Journal of Animal Ecology* 70: 459–475.
- Leathwick, J R; Rowden, A; Nodder, S; Gorman, R; Bardsley, S; Pinkerton, M; Baird, S J; Hadfield, M; Currie, K; Goh, A (2012) Benthic-optimised marine environment classification for New Zealand waters. *New Zealand Aquatic Environment and Biodiversity Report* No. 88. 54 p.
- McKenzie, D; Fletcher, D (2006) Characterisation of seabird captures in commercial trawl and longline fisheries in New Zealand 1997/98 to 2003/04. Final Research Report for Ministry of Fisheries project ENV2004/04. 102 p. (Unpublished report held by FNZ, Wellington.)
- Ministry for Primary Industries (2012) Aquatic Environment and Biodiversity Annual Review 2012. Compiled by the Fisheries Management Science Team, Ministry for Primary Industries, Wellington, New Zealand. 390 p.
- Middleton, D A J; Abraham, E R (2007) The efficacy of warp strike mitigation devices: Trials in the 2006 squid fishery. Final Research Report for research project IPA2006/02. (Unpublished report held by FNZ, Wellington.)
- Morgan, M J; Deblois, E M; Rose, G A (1997) An observation on the reaction of Atlantic cod (*Gadus morhua*) in a spawning shoal to bottom trawling. *Canadian Journal of Fisheries and Aquatic Sciences*, 54(S1): 217–223.
- Morgan, M J; Wilson, C E; Crim, L W (1999) The effect of stress on reproduction in Atlantic cod. *Journal of Fish Biology* 54(3): 477–488.
- Neil, H L; McMillan, P J; Tracey, D M; Sparks, R; Marriott, P; Francis, C; Paul, L J (2008) Maximum ages for black oreo (*Alloctytus niger*), smooth oreo (*Pseudocyttus maculatus*) and black cardinalfish (*Epigonus telescopus*) determined by the bomb chronometer method or radiocarbon ageing, and comments on the inferred life history of these species. Final Research Report for Ministry of Fisheries Research Project DEE2005-01 Objectives 1 & 2: 63 p. (Unpublished report held by FNZ, Wellington.)
- O'Driscoll, R L; Booth, J D; Bagley, N W; Anderson, O F; Griggs, L H; Stevenson, M L; Francis, M P (2003) Areas of importance for spawning, pupping or egg-laying, and juveniles of New Zealand deepwater fish, pelagic fish, and invertebrates. *NIWA Technical Report* 119. 377 p.
- Paulin, C; Stewart, A; Roberts, C; McMillan, P (1989) New Zealand Fish: a complete guide. *National Museum of New Zealand Miscellaneous Series* No. 19. 279 p.
- Phillips, N L (2002) Descriptive and catch-per-unit-effort (CPUE) analyses for black cardinalfish (*Epigonus telescopus*) in QMA 1. *New Zealand Fisheries Assessment Report* 2002/55. 54 p.
- Reiss, H; Greenstreet, S P R; Siebe, K; Ehrich, S; Piet, G J; Quirijns, F; Robinson, L; Wolff, W J; Kronke, I (2009) Effects of fishing disturbance on benthic communities and secondary production within an intensively fished area. *Marine Ecology Progress Series* 394: 201–213.
- Rice, J (2006) Impacts of Mobile Bottom Gears on Seafloor Habitats, Species, and Communities: A Review and Synthesis of Selected International Reviews. Canadian Science Advisory Secretariat Research Document 2006/057. 35 p. (available from http://www.dfo-mpo.gc.ca/CSAS/Csas/DocREC/2006/RES2006_057_e.pdf).
- Richard, Y; Abraham, E R (2013) Risk of commercial fisheries to New Zealand seabird populations. *New Zealand Aquatic Environment and Biodiversity Report* No. 109. 58 p.
- Richard, Y.; Abraham, E.R. (2015). Assessment of the risk of commercial fisheries to New Zealand seabirds, 2006–07 to 2012–13. *New Zealand Aquatic Environment and Biodiversity Report* 162. 85 p
- Richard, Y; Abraham, E; Berkenbusch, K (2017) Assessment of the risk of commercial fisheries to New Zealand seabirds, 2006–07 to 2014–15. *New Zealand Aquatic Environment and Biodiversity Report* 191. 104 p.
- Robertson, H A; Baird, K; Dowding, J E; Elliott, G P; Hitchmough, R A; Miskelly, C M; McArthur, N; O'Donnell, C F J; Sagar, P M; Scofield, R P; Taylor, G A (2017) Conservation status of New Zealand birds, 2016. *New Zealand Threat Classification Series* 19.
- Rowden, A A; Berkenbusch, K; Brewin, P E; Dalen, J; Neill, K F; Nelson, W A; Oliver, M D; Probert, P K; Schwarz, A-M.; Sui, P H; Sutherland, D (2012) A review of the marine soft-sediment assemblages of New Zealand. *New Zealand Aquatic Environment and Biodiversity Report* No 96.
- Thompson, F N; Abraham, E R (2009) Six Monthly Summary of the Capture of Protected Species in New Zealand Commercial Fisheries, Summer 2007–08. *New Zealand Aquatic Environment and Biodiversity Report* No. 35.
- Thompson, F N; Abraham, E R; Oliver, M D (2010) Estimation of fur seal bycatch in New Zealand trawl fisheries, 2002–03 to 2007–08. *New Zealand Aquatic Environment and Biodiversity Report* No. 56. 39 p.
- Thompson, F N; Berkenbusch, K; Abraham, E R (2013) Marine mammal bycatch in New Zealand trawl fisheries, 1995–96 to 2010–11. *New Zealand Aquatic Environment and Biodiversity Report* No. 105. 73 p.
- Tracey, D M (1993) Mercury levels in black cardinalfish (*Epigonus telescopus*). *New Zealand Journal of Marine and Freshwater Research* 27: 177–181.
- Tracey, D M; George, K; Gilbert, D J (2000) Estimation of age, growth, and mortality parameters of black cardinalfish (*Epigonus telescopus*) in QMA 2 (east coast North Island). *New Zealand Fisheries Assessment Report* 2000/27. 21 p.
- Tuck, I; Cole, R; Devine, J (2009) Ecosystem indicators for New Zealand fisheries. *New Zealand Aquatic Environment and Biodiversity Report* No. 42. 188 p.
- Williams, A; Schlacher, T A; Rowden, A A; Althaus, F; Clark, M R; Bowden, D A; Stewart, R; Bax, N J; Consalvey, M; Kloser, R J (2010) Seamount megabenthic assemblages fail to recover from trawling impacts. *Marine Ecology* 31 (Suppl. 1): 183–199.

BLADDER KELP ATTACHED (KBB G)*(Macrocystis pyrifera)***1. FISHERY SUMMARY**

Attached bladder kelp (KBB G) was introduced into the Quota Management System (QMS) on 1 October 2010, within FMA 3 and FMA 4 only which have the reporting codes KBB 3G and KBB 4G, respectively. The Total Allowable Catch (TAC), commercial, recreational, customary and other mortality allowances issued to KBB G on entering the QMS, and which remain unchanged, are presented in Table 1.

Bladder kelp, like all other large seaweeds, occurs in one of three states: attached (growing on the substrate); free-floating; and beach-cast. The attached growing state of bladder kelp is the only state managed under the QMS. Fisheries New Zealand will continue to monitor the use of beach-cast and free-floating seaweeds in FMAs 3 and 4, and will reconsider introducing these states into the QMS if sustainability and utilisation risks are identified in the future. Separate codes refer to beach cast bladder kelp in FMA 3 (KBB 3B) and free-floating bladder kelp in FMA 3 and 4 (KBB 3F and KBB 4F). Unless explicitly stated, this section refers to only attached bladder kelp.

Table 1: Total Allowable Catch (TAC, t), Total Allowable Commercial Catches (TACC, t), customary non-commercial (t), recreational, and other mortality allowances for attached bladder kelp on entering the QMS on 1 October 2010.

Fishstock	TAC	TACC	Customary Non-commercial	Recreational	Other Mortality
KBB 3G	1 238	1 237	0.1	0.1	1
KBB 4G	274	273	0.1	0.1	

1.1 Commercial fisheries

Bladder kelp has been used as a dietary supplement, fertilizer, cultivation for bioremediation purposes, as well as abalone and sea urchin feed (Buschmann et al 2006, Gutierrez et al 2006). There is current research evaluating the utilization of bladder kelp as feed for other aquaculture species such as shrimps (Buschmann et al 2006, Cruz-Suárez et al 2009), as well as an evaluation as a possible feedstock for conversion into ethanol for biofuel use (Wargacki et al 2012). Because of the growing demand for bladder kelp, MPI considered that the bladder kelp resource requires active management to ensure its sustainable use, and that management under the QMS was the most appropriate mechanism.

The season for commercial harvest of KBB G has been established between 1 October and 30 September, and catch is measured in greenweight (t).

Restrictions on New Zealand harvests of KBB G have been based on the Californian fishery (where the majority of research into harvesting effects has been conducted) and modified to take into account

BLADDER KELP ATTACHED (KBB G)

differences between California and New Zealand. These differences, compared to the Californian fishery, include reduced nutrients in New Zealand waters, the shallower depth at which KBB G is harvested in New Zealand, and the lack of information on New Zealand stocks.

The single restriction on KBB G harvest, implemented on introduction to the QMS on 1 October 2010, is a maximum cutting depth of 1.2 m.

Harvest of KBB G mainly occurs in QMA 3 and has varied since 2001–02 from 3 to 105 t (Table 2). Landings of KBB G in QMA 4 are minimal, with only 2.47 t reported in the last 15 years (Table 2).

Table 2: Reported landings for KBB G in greenweight (t) by fishing year. Blank cells indicate nil catches. Values above and below the horizontal line represent historic landings prior to QMS introduction and landings post QMS introduction, respectively. * Pre 2010 landings in KBB 3G include a combination of beach cast, free-floating and attached bladder kelp. Pre 2010 landings in KBB 4G may include a combination of free-floating and attached bladder kelp. Post 2010, the reported landings are for attached bladder kelp only.

Fishing Year	KBB 3G	KBB 4G	TACC KBB 3G	TACC KBB 4G
2001–02	104.50*	0.37*		
2002–03	37.00*			
2003–04	7.53*			
2004–05	17.90*			
2005–06	2.82*			
2006–07	8.35*			
2007–08	6.43*	2.10*		
2008–09	63.50*			
2009–10	28.37*			
2010–11	53.34		1 236	273
2011–12	34.25		1 236	273
2012–13	5.00		1 236	273
2013–14	94.00	0.00	1 236	273
2014–15	62.00	0.02	1 236	273
2015–16	30.54	0.00	1 236	273
2016–17	41.77	0.00	1 237	273

1.2 Recreational fisheries

There is no quantitative estimate of recreational harvest of bladder kelp at this time, although it is assumed to be restricted to the collection of beach-cast seaweed for composting. Consequently, recreational harvest of attached bladder kelp is assumed to be negligible.

1.3 Customary non-commercial fisheries

The harvest of bladder kelp by customary Maori is currently unrestricted. There is no quantitative information on the extent of customary harvest of attached bladder kelp (or any other state) in FMAs 3 and 4; however, the customary harvest of attached bladder kelp is likely to be negligible.

1.4 Illegal catch

Since introducing KBB G into the QMS, there is no quantitative or qualitative measure of illegal catch for bladder kelp.

1.5 Other sources of mortality

Hydrographic factors (e.g., tidal surge, nutrient limitation, temperature and salinity stress) and biological processes have been demonstrated to result in significant mortality of bladder kelp in the southern hemisphere (Buschmann et al 2004, 2006). Californian and Chilean studies have shown that grazing by sea urchins can result in the detachment of adult plants and their removal from the population (Dayton 1985a, Tegner et al 1995), and/or the removal of recruits and juvenile plants (Dean & Jacobsen 1984, Dean et al 1988, Vásquez et al 2006). In Chile, infestations of bladder kelp holdfasts by crustaceans (e.g., amphipods and isopods) may increase mortality by decreasing attachment strength (Ojeda & Santelices 1984). Due to their large size and high drag, adult bladder kelp are vulnerable to removal by high water motion (Dayton et al 1984, Seymour et al 1989, Schiel et al 1995, Fyfe & Israel 1996, Graham et al 1997, Fyfe et al 1999), which is considered the primary agent of mortality. In 1994, Fyfe et al (1999) found that winter storms extensively removed floating surface canopies at Pleasant River (north of Dunedin), and

that by February 1995, 50% of surface canopies had reformed. High seasonal and year-to-year variability in wave intensity and plant biomass results in high intra- and inter- annual variability in mortality. In California, uprooted plants may become entangled with attached plants, increasing drag and the likelihood of detachment, which may result in a 'snowball effect' capable of clearing large swaths in the local population (Dayton et al 1984). For example, Seymour et al (1989) observed that mortality of bladder kelp in California due to storm-induced plant detachment and entangled was as great as 94%. Graham et al (1997) observed that bladder kelp holdfast growth in California decreased significantly along a gradient of increasing wave exposure, possibly due to greater disturbance to the bladder kelp surface canopy, which reduces holdfast growth (Barilotti et al 1985, McCleneghan & Houk 1985). Thus, increased water motion and decreased holdfast strength can act in combination to decrease plant survival.

Sedimentation can also increase bladder kelp mortality – movement of bottom sediments can scour or bury bladder kelp spores and recruits, and the resuspension of sediments can reduce the amount of light reaching sub-canopy algae, preventing the attachment and development of spores, and inhibiting the growth of bladder kelp recruits (Dean & Jacobson 1984, Pirker 2002).

Over large spatial scales, elevated temperature also appears to be a major influence on bladder kelp mortality, and is likely to limit the northern distribution of bladder kelp within New Zealand (Hay 1990). For example, Hay (1990) described an apparent retraction of the distribution of bladder kelp within Cook Strait since 1942, presumably due to increasing surface water temperatures. Cavanaugh et al (2011) compared changes in canopy biomass with oceanographic and climatic data in California. They revealed that winter losses of regional kelp canopy biomass were positively correlated with significant wave height, while spring recoveries were negatively correlated with sea surface temperature. On interannual timescales, regional kelp-canopy biomass lagged the variations in wave height and sea surface temperatures by 3 years, indicating that these factors affect cycles of kelp recruitment and mortality. The dynamics of kelp biomass in exposed regions were related to wave disturbance, while kelp dynamics in sheltered regions tracked sea surface temperatures more closely.

Although wave disturbance and sea surface temperature appear to be the predominant sources of bladder kelp mortality, there are no quantitative estimates for these sources of mortality available for New Zealand. Further, the relevance of results from studies conducted outside New Zealand may be limited due to differences in hydrographic environment between New Zealand and other locales.

2. BIOLOGY

Historically, two species of bladder kelp, *Macrocystis pyrifera* (Linnaeus) C.Agardh and *M. integrifolia* Bory, were reported from both Northern and Southern Hemispheres, while *M. angustifolia* Bory and *M. laevis* Hay were reported from the Southern Hemisphere. However, *M. angustifolia*, *M. integrifolia* and *M. laevis* are currently regarded as taxonomic synonyms of *M. pyrifera* (Graham et al 2007, Demes et al 2009). Therefore, for the sake of this document, the four previously recognized species are simply referred to as bladder kelp, *Macrocystis pyrifera*.

Bladder kelp is globally widespread; it is found in the Atlantic Islands (Baardseth 1941, Chamberlain 1965); North America from Alaska to California, Baja and Mexico (e.g., Carr 1994, Graham et al 2007, Cavanaugh et al 2011); Central America (Taylor 1945); South America from Peru to Chile, Argentina and Uruguay (e.g., Vásquez et al 2006, Thiel et al 2007, Macaya & Zuccarello 2010); the Indian Ocean (Silva et al 1996); Tasmania (Cribb 1954, Womersley 1987); the Antarctic and the sub-Antarctic islands (Ricker 1987, John et al 1994) and New Zealand (Hay 1990, Fyfe & Israel 1996, Brown et al 1997, Hepburn et al 2007).

In New Zealand, bladder kelp has a broad latitudinal distribution, occurring in the southern North Island, the South Island, as well as Stewart, Chatham, Bounty, Antipodes, Auckland and Campbell Islands (Chapman & Chapman 1980, Adams 1994, Hurd & Pilditch 2011, Harper et al 2012). Bladder kelp does not persist in New Zealand waters where maximum temperatures exceed 18–19° C for several days (Hay 1990). The northern limit of bladder kelp is between Castle Point and Cape Turnagain on the East coast of the North Island, and Kapiti Island on the west coast of the North Island, and appears to correspond to the Southland current, which brings cool nutrient-rich water north from the south (Hay

BLADDER KELP ATTACHED (KBB G)

1990). The distribution of bladder kelp is generally patchy, and there is both seasonal and interannual variation in abundance (Hay 1990, Pirker et al 2000).

Bladder kelp can grow up to 45 m long in New Zealand, and occurs in water 3–20 m deep. Where the bottom is rocky and affords places for it to anchor, bladder kelp grows in extensive kelp beds with large floating canopies, and frequently forms colonies or large populations in calm bays, harbours or in sheltered offshore waters. It can tolerate a wide range of water motion in New Zealand, including areas where tidal currents reach 5–7 knots (Hay 1990). Smaller plants can be found in shallow pools and channels.




Figure 1: Diagram of the bladder kelp life cycle showing (left side) development of the young diploid sporophyte, increasing frond numbers through production of basal and apical meristematic blades; (right side) growth habit of an adult diploid sporophyte ca two years old, standing in 10 m of water depth, and liberating haploid zoospores; (below center) development of haploid gametophytes from settled zoospores, proceeding to gametogenesis, and fertilization yielding the zygote and, thence, a diploid embryonic sporophyte. From North (1986).

Bladder kelp is a large perennial kelp (individuals persist for up to 5 years in California; North 1994) with a life history progressing from planktonic zoospores (less than 3 days longevity) to microscopic benthic gametophytes (7–30 days longevity) and finally macroscopic benthic sporophytes (the large plants we see along the coast) (Figure 1). Adult sporophytes typically consist of numerous vegetative fronds that arise from longitudinal splits in meristem tissue (undifferentiated plant tissue which gives rise to new cells) located just above the holdfast. Vegetative fronds consist of a stipe (stem) terminating in an apical meristem (the primary point of growth at the tip of a frond) which gives rise to new vegetative blades as the frond develops (Figure 1). Blades are attached to the stipe by a single pneumatocyst (gas bladder), which provides buoyancy to the frond. Continued elongation of the stipe, combined with the production of new blades by the apical meristem, results in elongation of the frond and increases in the number of blades. Fronds continue to grow after reaching the surface, forming canopies (Figure 1). Finally, meristem activity ceases in the apical blade and a terminal blade is formed. In California, frond elongation has been observed occurring at a rate of up to 30 cm per day, making bladder kelp one of the fastest growing organisms on earth. Reproductive blades (called sporophylls) are clustered above the holdfast, forming

from the lowermost two to six blades on each frond (Figure 1). Sporophylls develop reproductive sporangia (spores) that are densely packed in sori (a cluster of sporangia) on the surface of the sporophylls. Californian studies have shown spores within sporangia take about 14 days to mature, with a mean residence time of about 30 days (Tugwell & Branch 1989). Each sporangium releases numerous mature zoospores that develop into gametophytes (North 1986).

A floating surface canopy consisting of numerous vegetative fronds characterizes adult plants. In California, the floating surface canopy comprises 33–50% of total plant biomass, and produces approximately 95% of organic production (Towle & Pearse 1973). Unlike other perennial kelp genera, giant kelp has limited nutrient and photosynphate storage capabilities, which in New Zealand is about 2 weeks (Brown et al 1997); consequently, growth by young fronds, reproductive material, holdfasts and other tissues near the base of the plant is supported by translocation of photosynphates from the canopy, which follows a source-sink relationship (North 1986). Mature canopy tissue exports both upward to the apical meristem at the frond apex, and downward to sporophylls, meristem tissue, holdfasts, and into apical regions of juvenile fronds (Schmitz & Lobban 1976, Lobban 1978, Manley 1984). The ability of bladder kelp to translocate photosynphates allows it to grow in dense aggregations with overlapping canopies that effectively shade out competitors on the bottom, yet supports rapid growth by young fronds, sporophylls, holdfasts and other tissues near the base of the plant.

The reliance on surface fronds for translocated photosynphate, combined with their vulnerability to disturbance, results in considerable spatial and temporal variability in giant kelp productivity and size. For example, Graham et al (1997), observed that bladder kelp holdfast growth in California decreased significantly along a gradient of increasing wave exposure, possibly due to greater disturbance to the bladder kelp surface canopy. Similarly, Miller & Geibel (1973) and McCleneghan & Houk (1985) observed reduced holdfast growth in bladder kelp following the experimental removal of surface canopies in California. Reed (1987) demonstrated that a 75% thinning of vegetative fronds in California led to an approximate 75% decrease in the generation of reproductive blades. Graham (2002) identified shifts in the reproductive condition of Californian bladder kelp from fertile to completely sterile in response to episodic, sub-lethal frond grazing by amphipods. This change in reproductive condition occurred despite relatively constant sporophyll biomass. Finally, in a New Zealand study, Geange (2014) identified an apparent tradeoff between vegetative growth and the generation of reproductive sporophylls. Relative to controls, the removal of surface canopies did not result in decreased frond generation, despite an 86% reduction in the generation of reproductive blades. Geange (2014) also found that 89% of plants became completely sterile 50 days after canopy removal, with effects persisting for up to 83 days.

Growth of bladder kelp in New Zealand appears to be seasonal, with autumn and winter growth rates in 1988 in Otago harbour having been estimated at approximately 1–20 mm per day (Table 3; Brown et al 1997). Brown et al (1997) identified a seasonal pattern of blade relative growth rate (RGR) in Otago Harbour, where blade RGR's during 1986–87 were similar year-round, except for summer when lower rates were recorded. Brown et al (1997) concluded that sufficiently high irradiance levels and seawater nutrient concentrations support relatively constant growth throughout most of the year, but that growth was nutrient-limited during summer months when seawater nitrate levels decline. In a study on Stewart Island, Hepburn et al (2007) found that exposure to waves increased nitrogen uptake, modifying the seasonal pattern of growth by ameliorating the negative effect of low seawater nitrogen concentrations during summer.

BLADDER KELP ATTACHED (KBB G)

Table 3: Growth parameters for KKB G canopy (> 2.25 m) and submerged fronds at Aquarium Point, Otago Harbour during autumn (March/April/May) and winter (June/July/August) 1988. From Brown et al (1997).

Growth parameter	FronD type	
	Canopy	Submerged
<i>FronD-elongation rate</i>		
autumn	1.9 cm d ⁻¹	1.2 cm d ⁻¹
winter	2.0 cm d ⁻¹	1.3 cm d ⁻¹
<i>Relative frond-elongation rate</i>		
autumn	0.0065 d ⁻¹	0.008 d ⁻¹
winter	0.0066 d ⁻¹	0.013 d ⁻¹
<i>Node-initiation rate</i>		
autumn	0.33 nodes d ⁻¹	0.28 nodes d ⁻¹
winter	0.30 nodes d ⁻¹	0.30 nodes d ⁻¹
<i>Relative node-initiation rate</i>		
autumn	0.0047 d ⁻¹	0.0064 d ⁻¹
winter	0.0044 d ⁻¹	0.0089 d ⁻¹
<i>Net blade-elongation rate</i>		
autumn	9.4 cm d ⁻¹	5.4 cm d ⁻¹
winter	12.8 cm d ⁻¹	12.1 cm d ⁻¹
<i>Elongation rate of immature blades</i>		
autumn	0.22 cm d ⁻¹	0.08 cm d ⁻¹
winter	0.21 cm d ⁻¹	0.10 cm d ⁻¹
<i>Relative elongation rate of immature blades</i>		
autumn	0.038 d ⁻¹	0.001 d ⁻¹
winter	0.036 d ⁻¹	0.001 d ⁻¹

3. STOCKS AND AREAS

In New Zealand, patches of bladder kelp are typically small and discrete, usually less than 100 m², although large beds (less than 1 km²) are found along the North Otago coast (Fyfe et al 1999). Although there is currently no data evaluating stock structure for bladder kelp in New Zealand, Alberto et al (2010, 2011) found low but significant genetic differentiation over a 70 km stretch of coast in the Santa Barbara Channel in southern California. In a New Zealand context, where stands of bladder kelp are small and discrete, these results suggest that stocks may display strong spatial structuring; however, these results should be viewed with caution because current regimes in the Santa Barbara Channel are strongly unidirectional.

4. ENVIRONMENTAL AND ECOSYSTEM CONSIDERATIONS

This section was reviewed by the Aquatic Environment Working Group for the May 2013 Fishery Assessment Plenary.

4.1 Role in the ecosystem

Forests of bladder kelp are amongst the most productive marine communities in temperate waters, they act as keystone species, altering the abiotic environment and providing vast amounts of energy and highly structured three-dimensional habitat (Foster & Schiel 1985, Graham 2004, Graham et al 2008). In California, bladder kelp has been identified as altering abiotic and biotic conditions by dampening water motion (Jackson & Winant 1983, Jackson 1998), altering sedimentation (North 1971), shading the sea floor (Reed & Foster 1984, Edwards 1998, Dayton et al 1999, Clark et al 2004), scrubbing nutrients from the water column (Jackson 1977, 1998), stabilising substrata (North 1971), and providing physical habitat for organisms both above and below the benthic boundary layer (Foster & Schiel 1985).

There are three primary components to the provisioning of habitat by attached bladder kelp: the holdfast, the mid-water fronds, and the surface canopy (Foster & Schiel 1985). Studies from California, Canada, Chile, the Sub-Antarctic, and Tasmania have shown that a highly diverse assemblage of organisms colonizes each of these three components. Holdfasts are primarily colonised by algae and invertebrates and encrusted with bryozoans and sponges. The mid-water fronds and surface canopies are host to a variety of sessile and mobile invertebrates (e.g., amphipods, top and turban snails), encrusting bryozoans, and hydroids. Juvenile and adult fishes may also associate with mid-water and canopy fronds, although kelp-fish associations in New Zealand appear to be weaker than those reported in California.

Although the following associations are not exclusive, the major species associated with bladder kelp forests in New Zealand include: (i) understory brown algae, *Ecklonia radiata*, *Carpophyllum flexuosum*, *Marginariella boryana* and *Cystophora platylobium*; (ii) a rich fauna of sessile invertebrates, including *Callana* spp., *Calliostoma granti*, *Cookia sulcata*, *Evechinus chloroticus*, *Haliotis iris*, *Trochus* spp.; and (iii) fishes, including *Notolabrus celidotus*, *N. cinctus*, *Odax pullus* and *Parika scaber* (Pirker et al 2000, Shears & Babcock 2007). Of these species, *Ecklonia radiata*, *Evechinus chloroticus* (kina) and *Haliotis iris* (paua) have significant recreational value.

A significant proportion of annual kelp production becomes free-floating and beach-cast in response to storm events, seasonal mortality, or ageing. Bladder kelp continues to provide habitat resources after detachment from the substratum. Studies in California, Chile, Macquarie Island, South Georgia and Tasmania, have shown that holdfasts, mid-water fronds and canopies can retain epifaunal fishes and mobile and sessile invertebrates when drifting long distances, and play an important role in the dispersal of invertebrates and fishes (Edgar 1987, Vásquez 1993, Helmuth et al 1994, Hobday 2000a,b,c, Smith 2002, Macaya et al 2005, Thiel & Gutow 2005a,b). Mature free-floating individuals may also be important in the connectivity of bladder kelp populations, and may explain low genetic diversity of bladder kelp over large geographic extents in the south eastern Pacific (Thiel et al 2007, Macaya & Zuccarello 2010).

The beach-cast state is either washed back into the sea over subsequent tidal cycles or remains in the beach environment, with New Zealand and Californian studies demonstrating that it is incorporated into physical beach processes, or into the terrestrial or marine food webs through consumption and decomposition (Inglis 1989, Lastra et al 2008). In New Zealand, beach-cast material supports a diverse ecology of organisms through nutrient cycling and decomposition, including various micro- and macro-fauna (Inglis 1989, Marsden 1991), and if washed up high enough on the beach, can aid sand dune formation.

4.2 Incidental catch (fish and invertebrates)

Small scale harvesting experiments carried out in Akaroa Harbour showed that harvesting canopy biomass had no measurable effect on bladder kelp and the dominant understorey species (Pirker et al 2000).

4.3 Incidental catch (marine mammals, seabirds and protected fish)

None known.

4.4. Benthic interactions

None known.

4.5 Other considerations

None known.

5. STOCK ASSESSMENT

Currently there is insufficient information on canopy area and density to allow for a stock assessment for KBB G. Furthermore, due to large temporal and spatial variation in bladder kelp growth, estimates of biomass should be looked at conservatively when applying regional scale management.

Large spatial and temporal fluctuations in biomass within and between individual kelp forests necessitates the need for initial annual stock assessments of targeted beds to determine credible biomass and sustainable yield information to ensure long-term sustainability (Pirker et al 2000). A combination of aerial photography and *in situ* measurements provide an easy method for assessing canopy biomass (Fyfe & Israel 1996, Fyfe et al 1999, Pirker et al 2000).

5.1 Estimates of fishery parameters and abundance

No estimates of fishery parameters or abundance are available at present.

5.2 Biomass Estimates

Maximum biomass occurs in winter (Cummack 1980, Pirker et al 2000). Growth rates and peaks in biomass can vary significantly over very short distances (i.e., kilometres) and temporal scales (i.e.,

BLADDER KELP ATTACHED (KBB G)

seasonally) in response to changes in currents, light, nutrient levels, and other environmental factors. Fyfe et al (1999) found that the wet biomass of closed canopy at Pleasant River in KBB 3 fluctuated from an estimated 10 639 g m⁻² (SE = 1566) in November 1995 to 3761 g m⁻² (SE = 1237) in November 1996. Pirker et al (2000) noted that marked differences exist in the demography of bladder kelp at a spatial scale of only a few kilometres – and that beds decline and regenerate at different times. Because of the apparent rapid spatio-temporal fluctuations in biomass, the status of KBB 3G and KBB 4G biomass is unknown and unable to be reliably estimated using best available information. Therefore, MPI was unable to ascertain whether the current biomass of both attached bladder kelp stocks is stable, increasing or decreasing.

There is some limited information on past harvestable bladder kelp biomass and potential yield at three sites in Akaroa Harbour (Wainui, Ohinepaka, and Mat White Bays: located in KBB 3G) (Pirker et al 2000). Pirker et al (2000) estimated a combined annual harvestable canopy biomass of 377 tonnes for 1999. Further, Pirker et al (2000) concluded that at Akaroa Harbour sites no one forest was capable of supporting the removal of consistent amounts of canopy, although two harvests could be sustained per year – one in late spring/early summer just prior to frond senescence, and then another cut in late autumn/early winter. However, this estimate should be treated with caution – the survey provides only seasonal point estimates of harvestable biomass during the time the survey was conducted, with the 1999 estimate being the highest. Further, the 1999 estimate does not provide an indication of biomass at a QMA level.

There is also some limited information on the location of bladder kelp beds throughout KBB 3, although the biomass of floating surface canopies is unknown. In November 1995, Fyfe et al (1999) used aerial photography to quantify whole plant biomass (surface canopies and subsurface fronds) of bladder kelp forests at Pleasant River. They estimated 42 ha of closed bladder kelp canopy and 43 ha of broken canopy, with a combined biomass of 7900 tonnes (\pm 1300). Shears & Babcock (2007) also provide a per square metre biomass estimates for entire bladder kelp plants from 247 sites within 43 locations across the North and South Islands (Figure 2) between 1999 and 2005. 12.1% of sites surveyed had bladder kelp, with a mean ash free dry weight (AFDW) biomass of 5.43 g m⁻². In KBB 3, biomass of attached bladder kelp ranged between 0.8 g AFDW m⁻² (\pm 0.5, Fiordland) and 374 g AFDW m⁻² (Banks Peninsula, figure 25 Shears & Babcock 2007). Again, estimates from these studies should be treated with caution as they only provide point estimates of biomass, estimates are not of harvestable biomass, and they do not provide estimates of biomass at the QMA level.




Figure 2: Mean biomass (g ash free dry weight m⁻²) of attached bladder kelp at all sites, averaged across 4 depth

5.3 Yield estimates and projections

As absolute biomass has not been estimated, MCY cannot be estimated.

CAY cannot be estimated.

5.4 Other yield estimates and stock assessment results

No information is available.

5.5 Other factors

It is not known whether the biomass of bladder kelp is stable or variable, but the latter is considered more likely.

6. STATUS OF THE STOCKS**KBB 3G****Stock Structure Assumptions**

No information is currently available to determine biological stocks for bladder kelp. Therefore, where quota has been allocated this has been to existing fishery management areas (3 and 4).

Stock Status	
Year of Most Recent Assessment	1995 and 1999
Assessment Runs Presented	Survey biomass from different parts of KBB 3
Reference Points	Interim Target: 40% B_0 Interim Soft Limit: 20% B_0 Interim Hard Limit: 10% B_0 Interim Overfishing threshold: F_{MSY}
Status in relation to Target	Due to the relatively low levels of exploitation it is likely that all stocks are still effectively in a virgin state, therefore they are Very Likely (> 90%) to be at or above the target.
Status in relation to Limits	Very Unlikely (< 10%) to be below the soft and hard limits
Status in relation to Overfishing	Overfishing is Very Unlikely (< 10%) to be occurring

Historical Stock Status Trajectory and Current Status	
	-
Fishery and Stock Trends	
Recent Trend in Biomass or Proxy	Unknown
Recent Trend in Fishing Intensity or Proxy	Fishing is light in KBB 3G averaging 33 t since 2001–02.
Other Abundance Indices	-
Trends in Other Relevant Indicators or Variables	-

Projections and Prognosis	
Stock Projections or Prognosis	Unknown
Probability of Current Catch or TACC causing Biomass to remain below, or to decline below, Limits	Current catches are Very Unlikely (< 10%) to cause declines below soft or hard limits
Probability of Current Catch or TACC causing Overfishing to continue or to commence	Current catches are Very Unlikely (< 10%) to cause overfishing to continue or commence

Assessment Methodology and Evaluation		
Assessment Type	Level 2 Partial quantitative stock assessment	
Assessment Method	Ground-truthed remote sensing biomass surveys	
Assessment Dates	Latest assessment: 1999 and 1995 (in different areas of KBB 3)	Next assessment: Unknown
Overall assessment quality rank	1-High quality: it is very likely that fishing is light and having little impact	

BLADDER KELP ATTACHED (KBB G)

Main data inputs (rank)	Biomass surveys	2 - Medium or mixed quality as surveys only cover part of the range and are dated
Data not used (rank)	-	-
Changes to Model Structure and Assumptions	-	-
Major Sources of Uncertainty	-	-

Qualifying Comments
There are large temporal and spatial fluctuations in biomass within and between beds; therefore, biomass estimates should be utilised conservatively.

Fishery Interactions
Bladder kelp plays an important role in structuring habitats and providing beach-cast material, but harvesting the canopy biomass has no known measurable effect on associated or dependent species.

KBB 4G**Stock Structure Assumptions**

No information is currently available to determine biological stocks for bladder kelp. Therefore where quota has been allocated this has been to existing fishery management areas (3 and 4).

Stock Status	
Year of Most Recent Assessment	None
Assessment Runs Presented	None
Reference Points	Interim Target: 40% B_0 Interim Soft Limit: 20% B_0 Interim Hard Limit: 10% B_0 Interim Overfishing threshold: F_{MSY}
Status in relation to Target	Due to the relatively low levels of exploitation it is likely that all stocks are still effectively in a virgin state, therefore they are Very Likely (> 90%) to be at or above the target
Status in relation to Limits	Very Unlikely (< 10%) to be below the soft and hard limits
Status in relation to Overfishing	Overfishing is Very Unlikely (< 10%) to be occurring

Historical Stock Status Trajectory and Current Status	-
--	---

Fishery and Stock Trends	
Recent Trend in Biomass or Proxy	Unknown
Recent Trend in Fishing Intensity or Proxy	Fishing is very light in KBB 4G with less than 3 t reported since 2001–02.
Other Abundance Indices	-
Trends in Other Relevant Indicators or Variables	-

Projections and Prognosis	
Stock Projections or Prognosis	Unknown
Probability of Current Catch or TACC causing Biomass to remain below, or to decline below, Limits	Current catches are Very Unlikely (< 10%) to cause declines below soft or hard limits
Probability of Current Catch or TACC causing Overfishing to continue or to commence	Current catches are Very Unlikely (< 10%) to cause overfishing to continue or commence

Assessment Methodology and Evaluation		
Assessment Type	-	
Assessment Method	-	
Assessment Dates	-	Next assessment: Unknown
Overall assessment quality rank	-	
Main data inputs (rank)	-	-
Data not used (rank)	-	-
Changes to Model Structure and Assumptions	-	
Major Sources of Uncertainty	-	

Qualifying Comments
There are large temporal and spatial fluctuations in biomass within and between beds; therefore, any biomass estimates in the future should be utilised conservatively.

Fishery Interactions
Bladder kelp plays an important role in structuring habitats and providing beach-cast material, but harvesting the canopy biomass has no known measurable effect on associated or dependent species.

7. RESEARCH NEEDS

Future high priority research areas include: (i) updated (or new in the case of KBB 4G) biomass surveys; (ii) an evaluation of stock structure and inter-stock genetic differentiation; and (iii) quantitative estimates for different sources of mortality.

8. FOR FURTHER INFORMATION

- Adams, N M (1994) *Seaweeds of New Zealand*. Canterbury University Press, 360 p.
- Alberto, F; Raimondi, P; Reed, D C; Coelho, N C; Leblois, R; Whitmer, A; Serrão, E A (2010) Habitat continuity and geographic distance predict population genetic differentiation in giant kelp. *Ecology* 91:49–56.
- Alberto, F; Raimondi, P T; Reed, D C; Watson, J R; Siegel, D A; Mitari, S; Coelho, N; Serrao, E A (2011) Isolation by oceanographic distance explains genetic structure for *Macrocystis pyrifera* in the Santa Barbara Channel. *Molecular Ecology* 20:25432–554.
- Anderson, T W (1994) Role of macroalgal structure in the distribution and abundance of a temperate reef fish. *Marine Ecology Progress Series* 113:279–290.
- Andrews, N L (1945) The kelp beds of the Monterey region. *Ecology* 26:24–37.
- Baardseth, E (1941) The marine algae of Tristan da Cunha. Results of the Norwegian Scientific Expedition to Tristan da Cunha 1937–1938 9:1–173.
- Barilotti, D C; McPeak, R H; Dayton, P K (1985) Experimental studies on the effects of commercial kelp harvesting in central and southern California *Macrocystis pyrifera* kelp beds. *California Fish and Game* 71:4–20.
- Beckley, L E; Branch, G M (1992) A quantitative scuba-diving survey of the sublittoral macrobenthos at subantarctic Marion Island. *Polar Biology* 11:553–563.
- Bernstein, B L; Jung, N (1979) Selective pressures and coevolution in a kelp canopy community in southern California. *Ecological Monographs* 49:335–355.
- Bray, R N; Ebeling, A W (1975) Food, activity, and habitat of three ‘picker-type’ microcarnivorous fishes in the kelp forests off Santa Barbara, California. *Fishery Bulletin* 73:815–829.
- Brown, M T; Nyman, M A; Keogh, J A; Chin, N K M (1997) Seasonal growth of the giant kelp *Macrocystis pyrifera* in New Zealand. *Marine Biology* 129:417–424.
- Buschmann, A H; Moreno, C A; Vásquez, J A; Hernández-González, M C (2006) Population and reproduction strategies of *Macrocystis pyrifera* (Phaeophyta) in southern Chile. *Journal of Applied Phycology* 18:575–582.
- Buschmann, A H; Vásquez, J A; Osorio, P; Reyes, E; Filún, L; Hernández-González, M C; Vega, A (2004) The effect of water movement, temperature and salinity on abundance and reproductive patterns of *Macrocystis* spp. (Phaeophyta) at different latitudes in Chile. *Marine Biology* 145:849–862.
- Carr, M H (1989) Effects of macroalgal assemblages on the recruitment of temperate zone reef fishes. *Journal of Experimental Marine Biology and Ecology* 126:59–76.
- Carr, M H (1991) Habitat selection and recruitment of an assemblage of temperate zone reef fishes. *Journal of Experimental Marine Biology and Ecology* 146:113–137.
- Carr, M H (1994) Effects of macroalgal dynamics on recruitment of a temperate reef fish. *Ecology* 75:1320–1333.
- Cavanaugh, K C; Siegel, D A; Reed, D C; Dennison, P E (2011) Environmental controls of giant kelp biomass in the Santa Barbara Channel. *Marine Ecology Progress Series* 429:1–17.
- Chamberlain, Y M (1965) Marine algae of Gough Island. *Bulletin of the British Museum (Natural History) Botany* 3:176–232.
- Chapman, V J; Chapman, D J (1980) *Seaweeds and their uses*. pp. [i-iv], v-ix, [x], 1–334. London & New York: Chapman & Hall.
- Clark, R P; Edwards, M S; Foster, M S (2004) Effects of shade from multiple kelp canopies on an understory algal assemblage. *Marine Ecology Progress Series* 267:107–119.
- Coyer, J A (1987) The mollusk assemblage associated with fronds of giant kelp (*Macrocystis pyrifera*) off Santa Catalina Island, California. *Bulletin of the Southern California Academy of Sciences* 85:129–138.
- Cribb, A B (1954) *Macrocystis pyrifera* (L.) Ag. in Tasmanian waters. *Australian Journal of Marine and Freshwater Research* 5:1–34.
- Cruz-Suárez, L; Tapia-Salazar, M; Nieto-López, M; Guajardo-Barbosa, C; Ricque-Marie, D (2009) Comparison of *Ulva clathrata* and the kelps *Macrocystis pyrifera* and *Ascophyllum nodosum* as ingredients in shrimp feeds. *Aquaculture Nutrition* 15:421–430.

BLADDER KELP ATTACHED (KBB G)


- Cummack, B.T. (1981) Ecology of *Macrocystis pyrifera* with special reference to growth and development of the sporophyte. MSc Thesis, University of Canterbury, 145 p.
- Dayton, P K (1985a) Ecology of kelp communities. *Annual Review of Ecology and Systematics* 16:215–245.
- Dayton, P K (1985b) The structure and regulation of some South American kelp communities. *Ecological Monographs* 55:447–468.
- Dayton, P K; Currie, V; Gerrodette, T; Keller, B D; Rosenthal, R J; Van Tresca, D (1984) Patch dynamics and stability of some California kelp communities. *Ecological Monographs* 54:253–289.
- Dayton, P K; Tegner, M J; Edwards, P B; Riser, K L (1999) Temporal and spatial scales of kelp demography: the role of oceanographic climate. *Ecological Monographs* 69:219–250.
- Dean, T A; Jacobsen, F R (1984) Growth of juvenile *Macrocystis pyrifera* (Laminariales) in relation to environmental factors. *Marine Biology* 83:301–311.
- Dean, T A; Jacobsen, F R; Thies, K; Lagos, S L (1988) Differential effects of grazing by white sea urchins on recruitment of brown algae. *Marine Ecology Progress Series* 48:99–102.
- DeMartini, E E; Roberts, D A (1990) Effects of giant kelp *Macrocystis* on the density and abundance of fishes in a cobble-bottom kelp forest. *Bulletin of Marine Science* 46:287–300.
- Demes, K W; Graham, M H; Suskiewicz, T S (2009) Phenotypic plasticity reconciles incongruous molecular and morphological taxonomies: the giant kelp, *Macrocystis* (Laminariales, Phaeophyceae), is a monospecific genus (note). *Journal of Phycology* 45:1266–1269.
- Dixon, J; Schroeter, S C; Kastendiek, J (1981) Effects of the encrusting bryozoan, *Membranipora membranacea*, on the loss of blades and fronds by the giant kelp, *Macrocystis pyrifera* (Laminariales). *Journal of Phycology* 17:341–345.
- Ebeling, A W; Laur, D R (1985) The influence of plant cover on surfperch abundance at an offshore temperate reef. *Environmental Biology of Fishes* 12:169–180.
- Edgar, G J (1987) Dispersal of faunal and floral propagules associated with drifting *Macrocystis pyrifera* plants. *Marine Biology* 95:599–610.
- Edwards, M S (1998) Effects of long-term kelp canopy exclusion on the abundance of the annual alga *Desmarestia ligulata* (Light F). *Journal of Experimental Marine Biology and Ecology* 228:309–326.
- Foster, M S; Schiel, D (1985) The ecology of giant kelp forests in California: A community profile. *U.S. Fish and Wildlife Service Biological Report* 85(7).
- Fyfe, J E; Israel, S A (1996) A window on an underwater habitat: Quantifying differences in giant kelp beds using colour aerial photographs and image processing software. 8th International Colloquium of the Spatial Information Research Centre, University of Otago, New Zealand, pp 95–102.
- Fyfe, J; Israel, S A; Chong, A; Ishmail, N; Hurd, C L; Probert, K (1999) Mapping marine habitats in Otago, Southern New Zealand. *Geocarto International* 14:17–26.
- Geange, S (2014) Growth and reproductive consequences of photosynthetic tissue loss in the surface canopies of *Macrocystis pyrifera* (L.) C. Agardh. *Journal of Experimental Marine Biology and Ecology* 453: 70–75.
- Ghelardi, R J (1971) The biology of giant kelp beds (*Macrocystis*) in California: species structure of the holdfast community. *Nova Hedwigia* 32:381–420.
- Graham, M H (1997) Factors determining the upper limit of giant kelp, *Macrocystis pyrifera* Agardh, along the Monterey Peninsula, central California, U.S.A. *Journal of Experimental Marine Biology and Ecology*. 218:127–149.
- Graham, M H (2002) Prolonged reproductive consequences of short-term biomass loss in seaweeds. *Marine Biology* 140:901–911.
- Graham, M H (2004) Effects of local deforestation on the diversity and structure of southern California giant kelp forest food webs. *Ecosystems* 7: 341–357.
- Graham, M; Halpern, B; Carr, M (2008) Diversity and dynamics of California subtidal kelp forests. In: McClanahan, T; Branch, G (editors). *Food webs and the dynamics of marine benthic ecosystems*. Oxford, UK: Oxford University Press. pp 103–134.
- Graham, M H; Harrold, C; Lisin, S; Light, K; Watanabe, J M; Foster, M S (1997) Population dynamics of giant kelp *Macrocystis pyrifera* along a wave exposure gradient. *Marine Ecology Progress Series* 148:269–279.
- Graham, M H; Vasquez, J A; Buschmann, A H (2007) Global ecology of the giant kelp *Macrocystis*: from ecotypes to ecosystems. *Oceanography and Marine Biology: an Annual Review* 45:39–88.
- Gutierrez, A; Correa, T; Muñoz, V; Santibañez, A; Marcos, R; Cáceres, C; Buschmann, A H (2006) Farming of the Giant Kelp *Macrocystis Pyrifera* in Southern Chile for Development of Novel Food Products. *Journal of Applied Phycology* 18:259–267.
- Hallacher, L E; Roberts, D A (1985) Differential utilization of space and food by the inshore rockfishes (Scorpaenidae: *Sebastes*) of Carmel Bay, California [U.S.A.]. *Environmental Biology of Fishes* 12:91–110.
- Harper, M A; Cassie Cooper, V; Chang, F H; Nelson, W A; Broady, P A (2012) Phylum Ochrophyta: brown and golden-brown algae, diatoms, silicoflagellates, and kin. In: *New Zealand inventory of biodiversity. Volume Three. Kingdoms Bacteria, Protozoa, Chromista, Plantae, Fungi*. (Gordon, D.P. Ed.). pp 114–163. Christchurch: Canterbury University Press.
- Hay, C H (1990) The distribution of *Macrocystis* (Phaeophyta: Laminariales) as a biological indicator of cool sea surface temperature, with special reference to New Zealand. *Journal of the Royal Society of New Zealand* 20:313–336.
- Helmuth, B S; Veit, R R; Holberton, R (1994) Long-distance dispersal of subantarctic brooding bivalve (*Gaimardia trapesina*) by kelp rafting. *Marine Biology* 120:421–426.
- Hepburn, C D; Holborow, J D; Wing, S R; Frew, R D; Hurd, C L (2007) Exposure to waves enhances the growth rate and nitrogen status of the giant kelp *Macrocystis pyrifera*. *Marine Ecology Progress Series* 339:99–108.
- Hobday, A J (2000a) Abundance and dispersal of drifting kelp *Macrocystis pyrifera* rafts in the Southern California Bight. *Marine Ecology Progress Series* 195:101–116.
- Hobday, A J (2000b) Age of drifting *Macrocystis pyrifera* (L.) C. Agardh rafts in the Southern California Bight. *Journal of Experimental Marine Biology and Ecology* 253:97–114.
- Hobday, A J (2000c) Persistence and transport of fauna on drifting kelp (*Macrocystis pyrifera* (L.) C. Agardh) rafts in the Southern California Bight. *Journal of Experimental Marine Biology and Ecology* 253:75–96.
- Holbrook, S J; Carr, M H; Schmitt, R J; Coyer, J A (1990) Effect of giant kelp on local abundance of reef fishes: the importance of ontogenetic resource requirements. *Bulletin of Marine Science* 47:104–114.
- Hurd, C L; Durante, K M; Chia, F S; Harrison, P J (1994) Effect of bryozoan colonization on inorganic nitrogen acquisition by the kelps *Agarum fimbriatum* and *Macrocystis integrifolia*. *Marine Biology* 121:167–173.
- Hurd, C L; Pilditch, C A (2011) Flow-induced morphological variations affect diffusion boundary-layer thickness of *Macrocystis pyrifera* (Heterokontophyta, Laminariales). *Journal of Phycology* 47:341–351.
- Inglis, G (1989) The colonisation and degradation of stranded *Macrocystis pyrifera* (L.) C. Ag. by the macrofauna of a New Zealand sandy beach. *Journal of Experimental Marine Biology and Ecology* 125:203–217.
- Jackson, G A (1977) Nutrients and production of giant kelp, *Macrocystis pyrifera*, southern California. *Limnology and Oceanography* 22:979–995.
- Jackson, G A (1998) Currents in the high drag environment of a coastal kelp stand off California. *Continental Shelf Research* 17:1913–1928.
- Jackson, G A; Winant, C D (1983) Effect of a kelp forest on coastal currents. *Continental Shelf Research* 2:75–80.
- John, D M; Pugh, P J A; Tittley, I (1994) Observations on the benthic marine algal flora of South Georgia: a floristic and ecological analysis. *Bulletin of the Natural History Museum, London, Botany* 24(2): 101–114.
- Jones, L G (1971) The biology of giant kelp beds (*Macrocystis*) in California: Studies on selected small herbivorous invertebrates inhabiting *Macrocystis* canopies and holdfasts in southern Californian kelp beds. *Nova Hedwigia* 32:343–367.

- Lastra, M; Page, H M; Dugan, J E; Hubbard, D M; Rodil, I F (2008) Processing of allochthonous macrophyte subsidies by sandy beach consumers: estimates of feeding rates and impacts on food resources. *Marine Biology* 154:163–174.
- Lobban, C S (1978) Translocation of ^{14}C in *Macrocystis pyrifera* (Giant Kelp). *Plant Physiology* 61:585–589.
- Macaya, E C; Boltaña, S; Hinojosa, I A; Macchiavello, J E; Valdivia, N A; Vásquez, N R; Buschmann, A H; Vásquez, J A; Vega, J M A; Thiel, M (2005) Presence of sporophylls in floating kelp rafts of *Macrocystis* spp. (Phaeophyceae) along the Chilean Pacific coast. *Journal of Phycology* 41:913–922.
- Macaya, E C; Zuccarello, G C (2010) Genetic structure of the giant kelp *Macrocystis pyrifera* along the southeastern Pacific. *Marine Ecology Progress Series* 420:103–112.
- Manley, S L (1984) Micronutrient uptake and translocation by *Macrocystis pyrifera* (Phaeophyta). *Journal of Phycology* 20:192–201.
- Marsden, I D (1991) Kelp-sandhopper interactions on a sand beach in New Zealand. I. Drift composition and distribution. *Journal of Experimental Marine Biology and Ecology* 152:61–74.
- McCleneghan, K; Houk, J L (1985) The effects of canopy removal on holdfast growth in *Macrocystis pyrifera* (Phaeophyta; Laminariales). *California Fish and Game* 71:21–27.
- Miller, D J; Geibel, J J (1973) Summary of blue rockfish and lingcod life histories, a reef ecology study, and giant kelp, *Macrocystis pyrifera*, experiments in Monterey Bay, California. State of California, Department of Fish and Game *Fish Bulletin* 158, 135 p.
- Moreno, C A; Jara, H F (1984) Ecological studies of fish fauna associated with *Macrocystis pyrifera* belts in the south of Feuguian Islands, Chile. *Marine Ecology Progress Series* 15:99–107.
- North, W J (1971) The biology of giant kelp beds (*Macrocystis*) in California: introduction and background. *Nova Hedwigia* 32:1–68.
- North, W J (1986) Biology of the *Macrocystis* resource in North America. *Fisheries Technology Papers, Food and Agriculture Organization (FAO), United Nations* 281:265–312.
- North, W J (1994) Review of *Macrocystis* biology. In: Akatsuka I (ed) *Biology of Economic Algae*. Academic Publishing, The Hague, The Netherlands.
- Ojeda, F P; Santelices, B (1984) Ecological dominance of *Lessonia nigrescens* (Phaeophyta) in central Chile. *Marine Ecology Progress Series* 19:83–91.
- Pirker, J J (2002) Demography, biomass production and effects of harvesting giant kelp *Macrocystis pyrifera* (Linnaeus) in southern New Zealand. PhD Thesis. University of Canterbury.
- Pirker, J J; Schiel, D R; Lees, H (2000) Seaweed Products for Barrel Culture Paua Farming. Unpublished Report for Foundation for Research Science and Technology's Technology for Business Growth Development project. 88 p.
- Reed, D C (1987) Factors affecting the production of sporophylls in the giant kelp *Macrocystis pyrifera* (L.) C.Ag. *Journal of Experimental Marine Biology and Ecology*. 113:61–69.
- Reed, D C; Foster, M S (1984) The effects of canopy shading on algal recruitment and growth of a giant kelp (*Macrocystis pyrifera*) forest. *Ecology* 65:937–948.
- Ricker, R W (1987) *Taxonomy and biogeography of Macquarie Island seaweeds*. pp. i–vi, [2], 1–344. London: British Museum (Natural History).
- Schiel, D R; Andrew, N L; Foster M S (1995) The structure of subtidal algal and invertebrate assemblages at the Chatham Islands, New Zealand. *Marine Biology*. 123:355–367.
- Schmitz, K; Lobban, C.S. (1976) A survey of translocation in Laminariales (Phaeophyta). *Marine Biology* 36:207–216.
- Seymour, R J; Tegner, M J; Dayton, P K; Parnell, P E (1989) Storm wave induced mortality of giant kelp *Macrocystis pyrifera* in southern California. *Estuarine Coastal and Shelf Science* 28:277–292.
- Shears, N T; Babcock, R C (2007) Quantitative description of mainland New Zealand's shallow subtidal reef communities. *Science for Conservation* 280. Department of Conservation, Wellington.
- Silva, P C; Basson, P W; Moe, R L (1996) Catalogue of the benthic marine algae of the Indian Ocean. *University of California Publications in Botany* 79: 1–1259.
- Smith, S D A (2002) Kelp rafts in the Southern Ocean. *Global Ecology and Biogeography* 11:67–69.
- Stebbins, T D (1986) Density, distribution, and feeding of the marine snail *Norrisia norrisi* (Mollusca: Gastropoda) on the kelp *Macrocystis pyrifera* (Phaeophyta: Laminariales). *Bulletin of the Southern California Academy of Sciences* 85:69–73.
- Stephens, J S; Morris, P M; Zerba, K; Love, M (1984) Factors affecting fish diversity on a temperature reef: the fish assemblage of Palos Verdes Point [California, U.S.A.], 1974–1981. *Environmental Biology of Fishes* 11:259–275.
- Taylor, W R (1945) Pacific marine algae of the Allan Hancock Expeditions to the Galapagos Islands. *Allan Hancock Pacific Expeditions* 12: i–iv, 1–528.
- Tegner, M J; Dayton, P K; Edwards, P B; Riser, K L (1995) Sea urchin cavitation of giant kelp (*Macrocystis pyrifera*) holdfasts and its effects on kelp mortality. *Journal of Experimental Marine Biology and Ecology* 191:82–99.
- Thiel, M; Gutow, L (2005a) The ecology of rafting in the marine environment. I: The floating substrata. *Oceanography and Marine Biology: An Annual Review* 42:181–264.
- Thiel, M; Gutow, L (2005b) The ecology of rafting in the marine environment. II: The rafting organisms and community. *Oceanography and Marine Biology: An Annual Review* 43:279–418.
- Thiel, M; Macaya, E C; Acuña, E; Arntz, W E; Bastias, H; Brokordt, K; Camus, P A; Castilla, J C; Castro, L R; Cortés, M; Dumont, C P; Escribano, R; Fernandez, M; Gajardo, J A; Gaymer, C F; Gomez, I; González, A E; González, H E; Haye, P A; Illanes, J E; Iriarte, J L; Lancellotti, D A; Luna-Jorquera, G; Luxoro, C; Manriquez, P H; Marín, V; Muñoz, P; Navarrete, S A; Perez, E; Poulin, E; Sellanes, J; Sepúlveda, H H; Stotz, W; Tala, F; Thomas, A; Vargas, C A; Vasquez, J A; Vega, J M (2007) The Humboldt current system of northern-central Chile: oceanographic processes, ecological interactions and socioeconomic feedback. *Oceanography and Marine Biology Annual Review* 45:195–344.
- Thiel, M; Vásquez, J A (2000) Are kelp holdfasts islands on the ocean floor? Indication for temporarily closed aggregations of peracarid crustaceans. *Hydrobiologia* 440:45–54.
- Towle, D W; Pearse, J S (1973) Production of giant kelp, *Macrocystis*, estimated by in situ incorporation of C-14 in polyethylene bags. *Limnology and Oceanography* 18:155–158.
- Tugwell, S; Branch, G M (1989) Differential polyphenolic distribution among tissues in the kelps *Eklonia maxima*, *Laminaria pallida* and *Macrocystis angustifolia* in relation to plant-defence theory. *Journal of Experimental Marine Biology and Ecology* 129:219–230.
- Vásquez, J A (1993) Effects on the animal community of dislodgement of holdfasts of *Macrocystis pyrifera*. *Pacific Science* 47:180–184.
- Vásquez, J A; Vega, J M A; Buschmann, A H (2006) Long term studies on El Niño-La Niña in northern Chile: effects on the structure and organization of subtidal kelp assemblages. *Journal of Applied Phycology*. 18:505–519.
- Wargacki, A J; Leonard, E; Win, M N; Regitsky, D D; Santos, C N S; Kim, P B; Cooper, S R; Raisner, R M; Herman, A; Sivitz, A B; Lakshmanaswamy, A; Kashiyaama, Y; Baker, D; Yoshikuni, Y (2012). An engineered microbial platform for direct biofuel production from brown macroalgae. *Science* 335:308–313.
- Watanabe, J M (1984a) Food preference, food quality and diets of three herbivorous gastropods (Trochidae: *Tegula*) in a temperate kelp forest habitat. *Oecologia* 62:47–52.
- Watanabe, J M (1984b) The influence of recruitment, competition, and benthic predation on spatial distributions of three species of kelp forest gastropods (Trochidae: *Tegula*). *Ecology* 65:920–936.
- Womersley, H B S (1987) *The marine benthic flora of southern Australia*. Part II. South Australian Government Printing Division, Adelaide. 481 p.

BLUE COD (BCO)

(Parapercis colias)

Rawaru



1. FISHERY SUMMARY

Allowances, TACCs and TACs are shown in Table 1.

Table 1: Recreational and Customary non-commercial allowances, other mortality, TACCs and TACs for blue cod by Fishstock.

Fishstock	Recreational Allowance	Customary non-commercial allowance	Other mortality	TACC	TAC
BCO 1	2	2	-	46	46
BCO 2	-	-	-	10	10
BCO 3	-	-	-	163	163
BCO 4	-	-	-	759	759
BCO 5	191	2	20	1 239	1 452
BCO 7	-	-	-	70	20
BCO 8	188	2	2	34	226
BCO 10	-	-	-	10	10

1.1 Commercial fisheries

Blue cod is predominantly an inshore domestic fishery with very little deepwater catch. The major commercial blue cod fisheries in New Zealand are off Southland and the Chatham Islands, with smaller but regionally significant fisheries off Otago, Canterbury, the Marlborough Sounds and Wanganui.

The fishery has had a long history. National landings of up to 3 000 t were reported in the 1930s and catches of 2500 t were sustained for many years in the 1950s and 1960s. Fluctuations in annual landings since the 1930s can be attributed to World War II, the subsequent market for frozen blue cod for a short period of time and then the development of the rock lobster fishery. Annual landings of blue cod also vary with the success of the rock lobster season. Traditionally many blue cod fishers were primarily rock lobster fishers. Therefore, the amount of effort in the blue cod fishery tended to depend on the success of the rock lobster season, with weather conditions in Southland affecting the number of 'fishable' days.

The commercial catch from the BCO 5 fishery is almost exclusively taken by the target cod pot fishery operating within Foveaux Strait and around Stewart Island (Statistical Areas 025, 027, 029 and 030). Similarly, the BCO 3 commercial catch is dominated by the target pot fishery, although blue cod is also taken as a small bycatch of the inshore trawl fisheries operating within BCO 3. Most

BLUE COD (BCO)

of the catch from BCO 3 is taken in the southern area of the Fishstock (Statistical Area 024). Catches from BCO 3 and 5 peak during autumn and winter and the seasonal nature of the fishery is influenced by the operation of the associated rock lobster fishery.

Total landings built up to a peak in 1985, the year before the QMS was implemented. Landings then declined up to 1989, but have since increased, coinciding with a change in the main fishing method from hand-lines to cod pots. Historical landings are shown in Table 2, recent reported landings are shown in Table 3 while Figure 1 shows the historical landings and TACC values for the five main BCO fish stocks.

Since 1994–95, total landings have exceeded 2 000 t annually, peaking at 2 501 t in 2003–04. Historically, the largest catches of blue cod have been taken in BCO 5 (1556 t in fishing year 2003–04). The total catch from this fishery remained relatively stable from 1982 to 1993 and subsequently increased to approach the level of the TACC in 1995–96. Catches have remained stable at this higher level in recent years.

Table 2: Reported landings (t) for the main QMAs from 1931 to 1982.

Year	BCO 1	BCO 2	BCO 3	BCO 4	Year	BCO 5	BCO 7	BCO 8
1931–32	29	0	55	148	1931–32	719	4	4
1932–33	12	0	59	111	1932–33	726	1	5
1933–34	24	5	26	1 055	1933–34	792	3	2
1934–35	17	5	23	1 306	1934–35	1057	0	4
1935–36	18	23	34	1 197	1935–36	284	44	2
1936–37	3	7	27	755	1936–37	113	61	0
1937–38	2	8	31	793	1937–38	172	81	0
1938–39	2	3	19	686	1938–39	94	57	0
1939–40	1	4	33	715	1939–40	135	68	0
1940–41	3	7	39	320	1940–41	177	72	0
1941–42	2	5	30	189	1941–42	128	54	0
1942–43	3	5	20	204	1942–43	139	65	0
1943–44	4	12	31	212	1943–44	221	80	0
1944	3	10	38	216	1944	552	88	0
1945	8	6	45	102	1945	634	109	0
1946	11	9	43	175	1946	715	116	2
1947	8	22	81	278	1947	955	153	1
1948	7	24	74	623	1948	852	88	2
1949	37	6	98	390	1949	929	82	3
1950	5	5	66	485	1950	1005	94	1
1951	4	9	51	494	1951	873	74	2
1952	5	7	53	543	1952	889	95	3
1953	7	20	62	682	1953	414	114	2
1954	5	9	84	603	1954	385	112	2
1955	4	8	83	355	1955	405	79	3
1956	1	7	86	636	1956	656	77	2
1957	2	5	63	1185	1957	581	61	2
1958	2	4	57	892	1958	542	71	2
1959	1	2	51	1158	1959	492	71	1
1960	1	4	48	903	1960	757	65	2
1961	1	2	43	871	1961	590	55	3
1962	1	9	37	550	1962	668	65	3
1963	1	12	46	633	1963	621	60	4
1964	1	107	83	495	1964	462	70	3
1965	1	18	55	742	1965	296	59	2
1966	1	395	35	13	1966	337	79	6
1967	1	437	34	0	1967	518	74	5
1968	1	312	69	0	1968	494	105	2
1969	6	232	92	8	1969	361	60	1
1970	0	402	70	39	1970	432	70	8
1971	1	105	81	36	1971	375	44	2
1972	0	137	60	3	1972	194	63	1
1973	1	127	65	4	1973	571	68	11
1974	0	67	61	1	1974	486	61	16
1975	0	5	42	2	1975	232	58	14
1976	0	103	72	17	1976	254	58	17
1977	2	3	21	46	1977	208	87	19
1978	0	9	49	14	1978	197	104	12
1979	0	17	74	13	1979	217	98	16
1980	1	1	89	1	1980	403	62	18
1981	1	2	69	40	1981	494	79	23
1982	7	0	62	13	1982	356	68	34

BLUE COD (BCO)

Table 3: Reported landings (t) of blue cod by Fishstock from 1983 to 2016–17 and actual TACCs (t) from 1986–87 to 2016–17. QMS data from 1986-present. FSU data 1983–1986.

Fishstock FMA (s)	BCO 1 1 & 9		BCO 2 2		BCO 3 3		BCO 4 4		BCO 5 5 & 6	
	Landings	TACC	Landings	TACC	Landings	TACC	Landings	TACC	Landings	TACC
1983*	23	-	4	-	81	-	192	-	626	-
1984*	39	-	6	-	74	-	273	-	798	-
1985*	21	-	3	-	55	-	274	-	954	-
1986*	19	-	2	-	82	-	337	-	844	-
1986–87	8	30	1	10	84	120	417	600	812	1 190
1987–88	9	40	1	10	148	140	204	647	938	1 355
1988–89	8	42	1	10	136	142	279	647	776	1 447
1989–90	10	45	1	10	121	151	358	749	928	1 491
1990–91	12	45	< 1	10	144	154	409	757	1 096	1 491
1991–92	10	45	1	10	135	154	378	757	873	1 536
1992–93	12	45	4	10	171	156	445	757	1 029	1 536
1993–94	14	45	2	10	142	162	474	757	1 132	1 536
1994–95	13	45	1	10	155	162	565	757	1 218	1 536
1995–96	11	45	2	10	158	162	464	757	1 503	1 536
1996–97	13	45	2	10	156	162	423	757	1 326	1 536
1997–98	16	45	4	10	163	162	575	757	1 364	1 536
1998–99	12	45	2	10	150	162	499	757	1 470	1 536
1999–00	14	45	2	10	168	162	490	757	1 357	1 536
2000–01	15	45	2	10	154	162	627	757	1 470	1 536
2001–02	12	46	2	10	138	163	648	759	1 477	1 548
2002–03	11	46	4	10	169	163	724	759	1 497	1 548
2003–04	9	46	4	10	167	163	710	759	1 556	1 548
2004–05	9	46	5	10	183	163	731	759	1 473	1 548
2005–06	7	46	1	10	183	163	580	759	1 346	1 548
2006–07	6	46	4	10	177	163	747	759	1 382	1 548
2007–08	6	46	3	10	167	163	779	759	1 277	1 548
2008–09	7	46	8	10	158	163	787	759	1 391	1 548
2009–10	8	46	7	10	171	163	691	759	1 210	1 548
2010–11	7	46	8	10	183	163	781	759	1 296	1 548
2011–12	6	46	8	10	166	163	753	759	1 215	1 239
2012–13	9	46	7	10	170	163	739	759	1 207	1 239
2013–14	9	46	8	10	159	163	720	759	1 208	1 239
2014–15	11	46	7	10	175	163	796	759	1 132	1 239
2015–16	9	46	6	10	169	163	758	759	1 099	1 239
2016–17	12	46	10	10	170	163	741	759	1 152	1 239

Fishstock FMA (s)	BCO 7 7		BCO 8 8		BCO 10 10		Total TACC
	Landings	TACC	Landing	TACC	Landings	TACC	
1983*	91	-	53	-	0	-	1 070
1984*	129	-	56	-	0	-	1 375
1985*	169	-	70	-	0	-	1 546
1986*	83	-	42	-	0	-	1 409
1986–87	79	110	22	60	0	10	2 130
1987–88	78	126	44	72	0	10	2 400
1988–89	66	131	32	72	0	10	2 501
1989–90	75	136	34	74	0	10	2 666
1990–91	63	136	28	74	0	10	2 667
1991–92	57	136	25	74	0	10	2 722
1992–93	85	136	32	74	0	10	2 724
1993–94	67	95	21	74	0	10	2 689
1994–95	113	95	24	74	0	10	2 689
1995–96	65	70	31	74	0	10	2 234
1996–97	71	70	38	74	0	10	2 029
1997–98	60	70	15	74	0	10	2 197
1998–99	52	70	35	74	0	10	2 220
1999–00	28	70	30	74	0	10	2 089
2000–01	26	70	22	74	0	10	2 316
2001–02	30	70	17	74	0	10	2 319
2002–03	39	70	13	74	0	10	2 457
2003–04	45	70	10	74	0	10	2 501
2004–05	44	50	7	74	0	10	2 452
2005–06	50	70	20	74	0	10	2 184
2006–07	69	70	34	74	0	10	2 413
2007–08	59	70	22	74	0	10	2 313
2008–09	58	70	18	74	0	10	2 427
2009–10	59	70	16	74	0	10	2 162
2010–11	51	70	16	74	0	10	2 342
2011–12	54	70	10	34	0	10	2 214
2012–13	71	70	12	34	0	10	2 215
2013–14	58	70	12	34	0	10	2 174
2014–15	68	70	8	34	0	10	2 198
2015–16	60	70	4	34	0	10	2 096
2016–17	65	70	5	34	0	10	2 155

BLUE COD (BCO)

Table 4: Reported total New Zealand landings (t) of blue cod for the calendar years 1970 to 1983. Sources MAF and FSU data.

Year	Landings
1970	1 022
1971	644
1972	459
1973	846
1974	696
1975	356
1976	524
1977	383
1978	378
1979	437
1980	536
1981	696
1982	539
1983	1 135




Figure 1: Reported commercial landings and TACC for the five main BCO stocks. From top: BCO 3 (South East Coast), and BCO 4 (South East Chatham Rise) [Continued on next page].




Figure 1 (continued): Reported commercial landings and TACC for the five main BCO stocks. From top: BCO 5 (Southland), BCO 7 (Challenger), and BCO 8 (Central Egmont).

1.2 Recreational fisheries

Blue cod are generally the most important recreational finfish in Marlborough, Otago, Canterbury, Southland and the Chatham Islands. Blue cod are taken predominantly by line fishing, but also by longlining, set netting, potting and spearfishing. The current allowances within the TAC for each Fishstock are shown in Table 1.

BLUE COD (BCO)

1.2.1 Management controls

The main methods used to manage recreational harvests of blue cod are minimum legal size limits (MLS), method restrictions and daily bag limits. Daily bag limits are specified as either blue cod specific (DL) or a combined species limit (CDL). The main management controls have changed over time and vary by Fishstock (Table 5). In addition there have been temporary and seasonal closures in the Marlborough Sounds and several Fiordland Sounds.

Table 5: Changes to minimum legal size (MLS in cm), blue cod specific daily bag limit (DL) and combined species daily bag limit (CDL) by Fishstock from 1986 to present. Slot = slot limit (legal size range). * DS = Doubtful Sounds, TS = Thompson's Sound, BS = Bradshaw Sound. ** C = inner sounds closed. # excluding Challenger East.

Fishstock	BCO 1		BCO 2		BCO 3		BCO3		BCO3		BCO 4	
Area	Auckland		Central (East)		South East (Otago)		North Canterbury		Kaikoura Marine Area		South East (Chatham Is.)	
	MLS	CDL	MLS	CDL	MLS	CDL	MLS	DL	MLS	DL	MLS	CDL
1986	30	30	30	30	30	30	30	30	N/A	N/A	30	30
1993	33	20	33	20	30	30	30	30	N/A	N/A	30	30
1994	33	20	33	20	30	30	30	30	N/A	N/A	30	30
2001	33	20	33	20	30	30	30	10	N/A	N/A	30	30
2008	30	20	33	20	30	30	30	10	N/A	N/A	30	30
2014	30	20	33	20	30	30	30	10	33	6	30	30
2017	30	20	33	20	30	30	30	10	33	6	30	30

Fishstock	BCO5		BCO5		BCO 5		BCO5		BCO 7		BCO7	
Area	Southland & Fiordland (External)		Patersons Inlet		Fiordland internal (excl. DS, TS, BS*)		DS, TS, BS*		Challenger West & South		Challenger East (incl. Marlborough Sounds)	
	MLS	CDL	MLS	DL	MLS	DL	MLS	DL	MLS	DL	MLS	DL
1986	30	30	30	30	33	20	33	20	30	30	30	12
1993	33	30	33	30	33	20	33	20	33	20	33	10
1994	33	30	33	15	33	20	33	20	33	20	28	6
2001	33	30	33	15	33	20	33	20	33	20	28	6
2003	33	30	33	15	33	20	33	20	33	20	30	3
2005	33	30	33	15	33	20	C*	C*	33	20	30	3
2008	33	30	33	15	33	20	C*	C*	33	20	C**	C**
2011	33	30	33	15	33	20	C*	C*	33	20	#SLOT 30–35	2
2014	33	20	33	15	33	20	C*	C*	33	20	#SLOT 30–35	2
2015	33	20	33	15	33	3	33	1	33	20	33	2
2017	33	20	33	15	33	3	33	1	33	20	33	2

Fishstock	BCO8		BCO10	
Area	Central (West)		Kermadec	
	MLS	DL	MLS	CDL
1986	30	30	30	30
1993	33	20	33	20
2014	33	10	33	20
2017	33	10	33	20

During 1992–93, the national minimum legal size (MLS) for blue cod increased from 30 cm to 33 cm for both amateur and commercial fishers, with the exception of BCO 3 and BCO 4 (South East management area). However, this was amended to 30 cm in 2008 for BCO 1, in response to a management review of blue cod in the area. Additionally, the Marlborough Sounds Area (part of BCO 7) had several MLS amendments between 1993 and 2015 (including a closure in the inner sounds followed by a slot limit of 30–35 cm in response to differing management approaches in the Marlborough Sounds. In 2014, the Kaikoura Marine Area in BCO 3 was established and the MLS of blue cod in this area was set at 33 cm.

The recreational daily bag limit (DL) has remained unchanged since 1993 in BCO 1, BCO 2, BCO 3 (South East Otago area), BCO 4, BCO 7 (Challenger West and South area) and BCO 10. In 2001, the recreational daily bag limit (DL) was reduced to 10 in the North Canterbury area (BCO 3). In 2014, the DL was set at 6 in the newly established Kaikoura Marine Area (BCO 3), and the DL was reduced to 20 in Southland and the external waters of the Fiordland marine area (BCO 5). Preceding these changes, the DL in Paterson's Inlet (BCO 5) was reduced from 30 to 15 in 1994. In 2005, new

commercial and recreational rules were introduced to the internal waters of the Fiordland Marine Area and Doubtful Sound, Thompson's Sound and Bradshaw Sound were closed to all blue cod fishing for 10 years. The closure was lifted in 2015 to recreational blue cod fishing and the new DL within Doubtful Sound was set at 1. The DL for the Challenger East area (BCO 7) has reduced five-fold from 10 to 2 since 1993 in response to differing management regimes in the area. In 2014, the DL in BCO 8 was reduced from 20 to 10.

1.2.2 Estimates of recreational harvest

Recreational harvest estimates are given in Table 6. There are two broad approaches to estimating recreational fisheries harvest: the use of onsite or access point methods where fishers are surveyed or counted at the point of fishing or access to their fishing activity; and, offsite methods where some form of post-event interview and/or diary are used to collect data from fishers.

The first estimates of recreational harvest for blue cod were calculated using an offsite approach, the offsite regional telephone and diary survey approach: MAF Fisheries South (1991–92), Central (1992–93) and North (1993–94) regions (Teirney et al 1997). Estimates for 1996 came from a national telephone and diary survey (Bradford 1998). Another national telephone and diary survey was carried out in 2000 (Boyd & Reilly 2002) and a rolling replacement of diarists in 2001 (Boyd et al 2004) allowed estimates for a further year (population scaling ratios and mean weights were not re-estimated in 2001).

The harvest estimates provided by these telephone diary surveys are no longer considered reliable for various reasons. With the early telephone/diary method, fishers were recruited to fill in diaries by way of a telephone survey that also estimates the proportion of the population that is eligible (likely to fish). A “soft refusal” bias in the eligibility proportion arises if interviewees who do not wish to co-operate falsely state that they never fish. The proportion of eligible fishers in the population (and, hence, the harvest) is thereby under-estimated. Pilot studies for the 2000 telephone/diary survey suggested that this effect could occur when recreational fishing was established as the subject of the interview at the outset. Another equally serious cause of bias in telephone/diary surveys was that diarists who did not immediately record their day's harvest after a trip sometimes overstated their harvest or the number of trips made. There is some indirect evidence that this may have occurred in all the telephone/diary surveys (Wright et al 2004).

The recreational harvest estimates provided by the 2000 and 2001 telephone diary surveys are thought to be implausibly high, which led to the development of an alternative maximum count aerial-access onsite method that provides a more direct means of estimating recreational harvests for suitable fisheries. The maximum count aerial-access approach combines data collected concurrently from two sources: a creel survey of recreational fishers returning to a subsample of ramps throughout the day; and an aerial survey count of vessels observed to be fishing at the approximate time of peak fishing effort on the same day. The ratio of the aerial count in a particular area to the number of interviewed parties who claimed to have fished in that area at the time of the overflight was used to scale up harvests observed at surveyed ramps, to estimate harvest taken by all fishers returning to all ramps. The methodology is further described by Hartill et al (2007).

This aerial-access method was first employed, optimised for SNA, in the Hauraki Gulf in 2003–04. It was then extended to survey the wider SNA 1 fishery in 2004–05 and to other areas (SNA 8) and other species, including blue cod in BCO 7 in 2005–06 (Davey et al 2008). The estimates for BCO 7 in 2005–06 are likely to be an underestimate due to less sampling coverage than planned for two key reasons. Fewer flights occurred than planned for the outer Marlborough Sounds due to poor flying conditions (low cloud), and sampling of harvest at boat ramps was not as complete as intended due to the higher than anticipated proportion of fishers who departed and returned to a bach/crib within BCO 7, or Wellington, without being intercepted at a boat ramp within BCO 7. A repeat aerial-access survey was conducted in BCO 7 over the 2015–16 fishing year (Hartill et al 2017) and this was considered by the Marine Amateur Fisheries Working Group to be more reliable than the initial survey. The recreational harvest of from BCO 7 in 2015–16 was about half that in 2005–06 (Table 6), almost all of the decrease being in the Marlborough Sounds.

BLUE COD (BCO)

In response to the cost and scale challenges associated with onsite methods, in particular the difficulties in sampling other than trailer boat fisheries, offsite approaches to estimating recreational fisheries harvest have been revisited. This led to the implementation of a national panel survey during the 2011–12 fishing year. The panel survey used face-to-face interviews of a random sample of 30 390 New Zealand households to recruit a panel of fishers and non-fishers for a full year. The panel members were contacted regularly about their fishing activities and harvest information collected in standardised phone interviews.

Table 6: Recreational harvest estimates for blue cod stocks. The telephone/diary surveys and aerial-access survey ran from December to November but are denoted by the January calendar year. The national panel survey ran through the October to September fishing year but is denoted by the January calendar year. Mean fish weights were obtained from boat ramp surveys (for the telephone/diary and panel survey harvest estimates).

Stock	Year	Method	Number of fish	Total weight (t)	CV
BCO 1	1996	Telephone/diary	34 000	17	0.11
	2000	Telephone/diary	37 000	23	0.31
	2012	Panel survey	17 463	8	0.20
BCO 2	1996	Telephone/diary	145 000	81	0.13
	2000	Telephone/diary	187 000	161	0.25
	2012	Panel survey	53 618	26	0.19
BCO 3	1996	Telephone/diary	217 000	151	11
	2000	Telephone/diary	1 026 000	752	0.29
	2012	Panel survey	212 184	101	0.20
BCO 5	1996	Telephone/diary	171 000	139	0.12
	2000	Telephone/diary	326 000	229	0.28
	2012	Panel survey	72 328	44	0.24
BCO 7	1996	Telephone/diary	356 000	239	0.09
	2000	Telephone/diary	542 000	288	0.20
	2006	Aerial-access	-	149	0.16
	2012	Panel survey	176 152	77	0.17
	2016	Aerial-access	-	75	0.15
BCO 8	1996	Telephone/diary	159 000	79	0.12
	2000	Telephone/diary	232 000	188	0.32
	2012	Panel survey	88 980	48	0.36

1.2.3 Charter vessel harvest

The national marine diary survey of recreational fishing from charter vessels in 1997–98 found blue cod to be the second most frequently landed species nationally and the most frequently landed species in the South Island. Results indicated that recreational harvests from charter vessels (Table 7) follow the same pattern as overall recreational harvest (Table 6). The estimated recreational harvests from charter vessels in BCO 7 exceeded the 1997–98 TACC and the commercial landings in QMA 7.

Table 7: Results of a national marine diary survey of recreational fishers from charter vessels, 1997–98 (November 1997 to October 1998).*

Fishstock	Number caught	CV(%)	Estimated landings (number of fish killed)	Point Estimate (t)
BCO 1	430	18	2 500	2.4
BCO 2	34	50	300	0.2
BCO 3	17 272	29	72 000	58
BCO 5	16 750	36	63 000	51
BCO 7	32 026	13	110 000	76
BCO 8	2	-	-	0

*Estimated number of blue cod harvested by recreational fishers on charter vessels by Fishstock and the corresponding harvest tonnage. The mean weights used to convert numbers to harvest weight were considered the best available at the time (James & Unwin 2000).

1.3 Customary non-commercial fisheries

No quantitative data on historical or current blue cod customary non-commercial catch are available. However, bones found in middens show that blue cod was a significant species in the traditional Maori take of pre-European times.

1.4 Illegal catch

No quantitative data on the levels of illegal blue cod catch are available.

1.5 Other sources of mortality

Blue cod have in the past been used for bait within the rock lobster fishery. Pots are either set specifically to target blue cod or have a bycatch of blue cod that is used for bait. However, these fish are frequently not recorded and the quantity of blue cod used as bait cannot be accurately determined.

Cod pots covered in 38 mm mesh frequently catch undersized blue cod. It has been estimated that in Southland, 65% of blue cod caught in these pots are less than 33 cm. When returned, the mortality of these fish can be high due to predation by mollymawks following commercial boats. It is estimated by the fishing industry that up to 50% of returned fish can be taken. To reduce the problem of predation of returned undersized fish, a minimum 48 mm mesh size was introduced to BCO 5 in 1994. However, no mesh size restrictions exist in any other area.

Recreational line fishing often results in the harvest of undersized blue cod. The survival of these has been shown to be a factor of hook size. A small scale experiment showed that returned undersized fish caught with small hooks (size 1/0) experience 25% mortality, whereas those caught with large hooks (size 6/0) appear to have little or no mortality (Carbines 1999).

2. BIOLOGY

Blue cod is a bottom-dwelling species endemic to New Zealand. Although distributed throughout New Zealand near foul ground to a depth of 150 m, they are more abundant south of Cook Strait and around the Chatham Islands. Growth may be influenced by a range of factors, including sex, habitat quality and fishing pressure relative to location (Carbines 2004a). Size-at-sexual maturity also varies according to location. In Northland, maturity is reached at 10–19 cm total length (TL) at an age of 2 years, whilst in the Marlborough Sounds it is reached at 21–26 cm (TL) at 3–6 years. In Southland, the fish become mature at 26–28 cm (TL), at an age of 4–5 years. Blue cod have also been shown to be protogynous hermaphrodites, with individuals over a large length range changing sex from female to male (Carbines 1998). Validated age estimates using otoliths have shown that blue cod males grow faster and are larger than females (Walsh 2017). The maximum recorded age for this species is about 32 years.

M was estimated using the equation $M = \log_e 100/\text{maximum age}$, where maximum age is the age to which 1% of the population survives in an unfished stock. Using the maximum age of 32 years, (Carbines et al 2007) M was calculated to be 0.14. This estimate seems feasible as in lightly fished areas such as the offshore Banks Peninsula Z is thought to approximate M and was calculated at 0.14 to 0.15 (Beentjes 2012).

Blue cod have an annual reproductive cycle with an extended spawning season during late winter and spring. Spawning has been reported within inshore and mid-shelf waters. It is also likely that spawning occurs in outer-shelf waters. Ripe blue cod are also found in all areas fished commercially by blue cod fishers during the spawning season. Batch fecundity was estimated by Beer et al (2013). Eggs are pelagic for about five days after spawning, and the larvae are pelagic for about five more days before settling onto the seabed. Juveniles (less than about 10 cm TL) are not caught by commercial potting or lining, and therefore blue cod are not vulnerable to the main commercial fishing methods until they are mature. Recreational methods do catch juveniles, but since this species does not have a swim bladder, the survival of these fish is good if they are caught using large hooks (6/0)(which do not result in gut hooking) and returned to the sea quickly (Carbines 1999).

BLUE COD (BCO)

Tagging experiments carried out in the Marlborough Sounds in the 1940s and 1970s suggested that most blue cod remained in the same area for extended periods. A more recent tagging experiment carried out in Foveaux Strait (Carbines 2001) showed that although some blue cod moved as far as 156 km, 60% travelled less than 1 km. A similar pattern was found in Dusky Sound where four fish moved over 20 km but 65% had moved less than 1 km (Carbines & McKenzie 2004). The larger movements observed during this study were generally eastwards into the fiord. The inner half of the fiord was found to drain the outer strata and had 100% residency.

Biological parameters relevant to stock assessment are shown in Table 8.

Table 8: Estimates of biological parameters for blue cod. These estimates are survey specific and reflect varying exploitation histories and environmental conditions. Only von Bertalanffy growth parameters derived from otoliths aged using the Age Determination Protocol for Blue Cod (Walsh 2017) are included in this table.

Fishstock		Estimate			Source			
<u>1. Natural mortality (<i>M</i>)</u>								
All		0.14			Estimated from the maximum age in Carbines et al (2007), using Hoenig's (1983) method.			
<u>2. Von Bertalanffy growth parameters</u>								
		Females			Males			
Survey/year		L_{∞}	K	t_0	L_{∞}	k	t_0	
Dusky Sound (2014)		46.7	0.129	-1.8	50.3	0.222	0.638	Beentjes & Page (2016)
Kaikoura (2015)		40.7	0.174	-1.12	52.3	0.171	-0.27	Beentjes & Page (2017)
Banks Peninsula (2016)		50.2	0.116	-2.07	58.7	0.134	-1.21	Beentjes & Fenwick (2017)
Marlborough Sounds (2017)		32.2	0.52	0.83	39.9	0.37	0.69	Beentjes et al. (in prep)
<u>3. Weight = a(length)^b (Weight in g, length in cm total length).</u>								
Area			a	b		R ²		
Kaikoura	2011	Male	0.011793	3.09246	0.97		Carbines & Haist (2012b)	
	2011	Female	0.007042	3.23949	0.95			
Motunau	2012	Male	0.01490	3.03796	0.98		Carbines & Haist (2012b)	
	2012	Female	0.01384	3.05982	0.97			
Banks Peninsula	2012	Male	0.019138	2.98181	0.98		Carbines & Haist (2012a)	
	2012	Female	0.016939	3.02644	0.96			
North Otago	2013	Male	0.01093	3.10941	0.98		Carbines & Haist (2014b)	
	2013	Female	0.012023	3.09201	0.97			
South Otago	2013	Male	0.008472	3.19011	0.99		Carbines & Haist (2014c)	
	2013	Female	0.008617	3.1863	0.99			
Fiordland (Dusky Sound)	2002	Male	0.007825	3.1727	0.97		Carbines & Beentjes (2003)	
	2002	Female	0.00506	3.2988	0.98			
Stewart Island (Paterson Inlet)	2010	Male	0.00663	3.2469	0.98		Carbines & Haist (2014a)	
	2010	Female	0.00663	3.2469	0.98			

The preliminary results of a mitochondrial DNA analysis (Smith 2012) suggest that the Chatham Island blue cod are likely to be genetically distinct from mainland New Zealand. Over larger distances the mainland New Zealand blue cod appear to show a pattern of Isolation-by-Distance or continuous genetic change among populations.

3. STOCKS AND AREAS

The FMAs are used as a basis for Fishstocks, except FMAs 5 and 6, and FMAs 1 and 9, which have been combined. The choice of these boundaries was based on a general review of the distribution and relative abundance of blue cod within the fishery.

There are no data that would alter the current stock boundaries. However, tagging experiments suggest that blue cod populations may be geographically isolated from each other, and there may be several distinct populations within each management area (particularly those occurring in sounds and inlets).

4. STOCK ASSESSMENT

4.1 Estimates of fishery parameters and abundance

4.1.1 South Island blue cod potting surveys

Marlborough Sounds

In 1995, a fishery independent survey using standardised cod pots at fixed stations provided catch rate estimates for recruited blue cod in Queen Charlotte Sound and outer Pelorus Sound. In 1996 a second potting survey covered all of Pelorus Sound as well as the east coast of D'Urville Island (Blackwell 1997, 1998). A 2001 survey (Blackwell 2002) included Queen Charlotte Sound, Pelorus Sound, and east D'Urville, and a survey in 2004 covered the same areas as 2001 but was expanded to include west D'Urville and Separation Point (Blackwell 2005). In 2007, the surveyed area was the same as 2004 except that Separation Point was dropped. In 2008 a standalone survey of a Cook Strait stratum was carried out and in 2010 the Cook Strait stratum was added to the surveyed area along with those strata used in 2007 (Beentjes & Carbines 2012). A new survey in 2013 used the same strata as 2010 (Beentjes et al 2014). The 2001 to 2008 surveys were reanalysed as part of the 2010 survey so that they were consistent with methods used for recent surveys (Beentjes & Carbines 2012). The 1995 and 1996 surveys, similarly, have been reanalysed as part of the 2013 survey analyses (Beentjes et al 2014). All surveys before 2010 used fixed sites which were selected randomly from a wider list of fixed sites within a given stratum. These fixed locations are available to be used repeatedly on subsequent surveys in that area (Beentjes & Francis 2011). In 2010, experimental random sites were trialled in selected strata. Random sites may have any location (single latitude and longitude) and are generated randomly within each stratum. In 2013 and 2017 (Beentjes et al 2017, Beentjes et al. in prep), full random and full fixed site surveys were conducted concurrently. Of the three random-site surveys only the last two are (2013 and 2017) are comparable.

Throughout the fixed-site surveys, catch rates of total blue cod (all sizes) have tended to be highest around D'Urville Island, lowest in Cook Strait, and similar between Queen Charlotte Sound and Pelorus Sound (Figure 2, Table 9). In Queen Charlotte Sound catch rates progressively declined from 2.1 to 1.1 kg.pot⁻¹ (CVs range 16 to 26%) between 1995 and 2007 before increasing markedly in 2010 to 1.75 kg.pot⁻¹ (Figure 2). From October 2008 to April 2011, the inner Sounds were closed to recreational blue cod fishing and the 2010 potting survey increased abundance in Queen Charlotte Sound is attributed to the closure. In Pelorus Sound, total blue cod catch rates declined from 2.4 to 1.1 kg.pot⁻¹ (CVs range 7 to 19%) over the same period, before increasing again in 2010, to 2.9 kg.pot⁻¹ (Figure 2). Pelorus Sound showed a similar trend in catch rates to Queen Charlotte Sound, dropping markedly from 1996 to 2007 and increasing again in 2010 after two years of closure. In April 2011, a seasonal opening with a "slot" limit (which allowed the take of blue cod between 30 and 35 cm) was introduced for the Marlborough Sounds Management Area, an area that includes inner and outer Queen Charlotte and Pelorus Sounds and east D'Urville. The 2013 survey was carried out two years after the slot limit had been in place, with total blue cod catch rates for both Queen Charlotte and Pelorus Sounds declining compared to 2010, but remaining higher than 2001 to 2007 for Pelorus Sound when the fishery was open, and about the same magnitude as pre-closure for Queen Charlotte Sound (Figure 2). In the D'Urville Island strata, which have been fished

BLUE COD (BCO)

continuously over the same period, catch rates for total blue cod between 2004 and 2013 have been stable, ranging from 3.9 to 4.44 kg.pot⁻¹ (CVs range 8 to 18%) (Figure 2). D'Urville was not closed to fishing in October 2008, but the east side of the island was included in the management area where the "slot limit" has been applicable since April 2011. Cook Strait has three comparable random-site surveys (2010, 2013 and 2017) with the first survey in 2008 being a fixed-site survey which was not comparable. Total blue cod catch rates from the Cook Strait random surveys ranged from 0.7 – 1.1 kg.pot⁻¹ with no trend (Table 9). There were no closures or slot limit management measures for Cook Strait. The proportion of the total biomass within the "slot limit" (30–35 cm) in 2013 was 45%, 49% and 49% for QCH, PEL, and DUR regions respectively, while proportions of biomass above the slot limit were 26%, 25% and 22%, respectively. Sex ratios have been dominated by males in all regions over all surveys (Table 9). The 2017 survey took place 2 years after the slot-limit was removed and in the Marlborough Sounds Area the MLS was increased to 33 cm. In 2017, catch rates from the fixed-site survey in Queen Charlotte Sound were similar to those in 2013, in Pelorus Sound they were similar to 2010, and in D'Urville Island they were about 40% higher than in 2013 (Figure 2, Table 9). The random site surveys in 2013 and 2017 generally have lower catch rates than fixed-site surveys and although the patterns among strata in each region are similar, they do not show the same overall trends as fixed sites by region (Table 9, Figure 2). In Queen Charlotte Sound survey biomass increased markedly, whereas for Pelorus Sound and D'Urville Island there are no significant changes. Cook Strait random-site catch rates show no significant difference from 2010 to 2017. The overall Marlborough Sounds catch rates from 2004 onward (where survey strata are consistent among surveys) indicates that blue cod were more abundant in 2017 than any of the previous years (Figure 2). It is the intention to transition to random-site surveys and conducting both fixed and random site surveys allows comparison of catch rates, length and age composition, and sex ratios between survey designs in the interim. The next survey in the time series will use only a random site design.

A random-site survey of Long Island Marine Reserve in 2017, in which all fish were returned alive (unsexed), had mean catch rates of all blue cod of 8.76 kg.pot⁻¹ (CV of 15%), substantially higher than adjacent fished strata in Queen Charlotte Sound (Table 9). In addition, the mean size was 3.2 cm greater in the marine reserve and length-frequency distributions were bimodal in contrast to the unimodal distributions from adjacent strata in Queen Charlotte Sound.

Growth rates and age compositions were similar for 2013 and 2017. Fixed-site survey Chapman Robson total mortality estimates (Z) for age at recruitment of 6 years were very close at 0.51 in 2013 and 0.53 in 2017 (Table 10). Spawner-per-recruit ratios ($F_{SPR\%}$), however, differed substantially and were 25% in 2013 and 39% in 2017 (The Fisheries New Zealand target is $F_{45\%}$). The difference was primarily a result of having different selectivity ages to the fishery because the MLS increased from 30 cm in 2013 to 33 cm in 2017, and hence these ratios cannot be validly compared. Similarly, random-site survey CR total mortality estimates (Z) for age at recruitment of 6 years were very close at 0.46 in 2013 and 0.52 in 2017 (Table 10). Spawner-per-recruit ratio ($F_{SPR\%}$), also, differed substantially and were 27% in 2013 and 39% in 2017 for the same reasons.




Figure 2: Marlborough Sounds fixed-site and random-site potting survey catch rates of all blue cod by survey year for each region and overall for the Marlborough Sounds. Error bars are 95% confidence intervals. There were no complete fixed-site surveys in QCH in 1996, PEL in 1996, and DUR from 1995 to 2001. For the overall Marlborough Sounds plot, the 2004 and 2007 fixed-site surveys exclude Separation Point, and the random-site surveys exclude Cook Strait, hence the strata are consistent among the surveys for fixed and random site surveys.

BLUE COD (BCO)

Table 9: Summary statistics from standardised blue cod fixed-site and random-site potting surveys in the Marlborough Sounds up to 2017 by region. Mean length and sex ratios are derived from the scaled population length distributions. Results for each region are shown only for surveys where strata have remained the same throughout the time series and results are for all blue cod. For the overall Marlborough Sounds (All MS), the 2004 and 2007 fixed-site surveys exclude Separation Point, and the random-site surveys exclude Cook Strait, hence the strata are consistent among the surveys for fixed and random site surveys. QCH, Queen Charlotte Sound; PEL, Pelorus Sound; DUR, D'Urville; CKST, Cook Strait; LIMR, Long Island Marine Reserve, MS all, all Marlborough Sounds

Region/strat	Year	Site type	Mean length (cm)			Overall	CPUE (kg.pot ⁻¹)	Sex ratio (% male)
			Male	Female	unsexed		range (CV)	
QCH	1995	Fixed	31.0	28.0		2.1	0.74–2.91 (12%)	59%
	1996	–	–	–		–	–	–
	2001	Fixed	28.5	24.3		1.33	0.58–1.69(12%)	61%
	2004	Fixed	27.9	24.2		1.16	0.35–2.01(22%)	51%
	2007	Fixed	29.8	25.7		1.09	0–2.60(15%)	69%
	2010	Fixed	33.2	29.0		2.09	0.60–2.56(18%)	71%
	2013	Fixed	31.7	29.8		1.0	0.32–1.12 (18%)	62%
		Random	32.1	30.3		0.49	0.22–1.07 (27%)	66%
		Fixed	32.2	29.6		0.86	0.18–1.95 (27.3%)	72%
	Random	32.5	30.7		1.04	0.11–1.94 (15%)	73%	
QCH/LIMR	2017	Random	–	–	35.2	8.76	8.76 (14%)	–
PEL	1995	–	–	–		–	–	–
	1996	Fixed	29.8	26.2		2.4	1.0–3.3 (7%)	70%
	2001	Fixed	27.8	22.2		0.67	0.19–1.46(12%)	64%
	2004	Fixed	28.2	23.5		0.96	0.20–2.70(11%)	66%
	2007	Fixed	29.2	24.5		1.07	0.28–3.24(11%)	77%
	2010	Fixed	32.8	28.3		2.9	1.6–3.86(13%)	87%
	2013	Fixed	31.3	27.2		1.95	3.3–4.94(15%)	89%
		Random	33.3	30.1		1.18	0.18–3.96 (12%)	77%
		Fixed	32.0	29.5		3.20	0.11–10.1 (17%)	86%
	Random	32.4	29.8		0.90	0.07–2.77 (23%)	90%	
DUR	1995	–	–	–		–	–	–
	1996	–	–	–		–	–	–
	2001	–	–	–		–	–	–
	2004	Fixed	30.7	27.8		4.23	3.75–4.67(11%)	50%
	2007	Fixed	32.2	29.5		4.15	2.92–5.49(10%)	71%
	2010	Fixed	31.3	28.7		3.82	2.15–5.64(8%)	64%
	2013	Fixed	31.7	29.4		3.88	3.37–4.44(18%)	70%
		Random	32.8	29.9		2.31	1.42–3.28(43%)	57%
		Fixed	32.9	30.6		6.52	4.5–8.7 (15%)	61%
	Random	32.6	30.6		3.59	2.9–4.3 (24%)	65%	
CKST	2008	Fixed	31.9	26.4		1.50	0.30–4.20(15%)	88%
	2010	Random	30.5	25.6		1.06	0.11–1.74(22%)	84%
	2013	Random	31.7	28.4		0.70	0.14–1.62(12%)	83%
	2017	Random	32.3	28.2		1.10	0.08–2.67(28%)	87%
All MS	2004	Fixed	29.1	25.9		1.92	0.37–4.67 (8%)	54
	2007	Fixed	30.7	27.2		1.81	0–5.48 (7%)	72
	2010	Fixed	32.5	28.7		2.83	0.60–5.64 (7%)	75
	2013	Fixed	31.5	29.1		2.68	0.31–4.44 (10%)	76
		Random	32.9	30.0		1.20	0.22–3.96 (21%)	66
		Fixed	32.4	30.2		3.15	0.11–8.73 (10%)	72
		Random	32.5	30.6		1.59	0.06–4.32 (14%)	72

Table 10: Mortality parameters (Z , F and M) and spawner-per-recruit (F_{SPR}) estimates for blue cod from the 2013 and 2017 Marlborough Sounds fixed-site and random-site potting surveys for all regions combined. F , fishing mortality; M , natural mortality; Z , total mortality; Age at recruitment = 6 years equivalent to age at which females reach MLS of 30 cm in 2013, and males and females combined reach MLS of 33 cm in 2017. Otoliths from both surveys were aged using the Age Determination Protocol for blue cod (Walsh 2017). CIs, 95% confidence intervals.

Survey	Region	Site type	M	Z (CIs)	F	F_{SPR}
2013	All regions combined	Fixed	0.14	0.56 (0.40–0.74)	0.42	$F_{25.5\%}$
2017			0.14	0.53 (0.38–0.72)	0.39	$F_{39.0\%}$
2013	All regions combined	Random	0.14	0.53 (0.38–0.70)	0.39	$F_{26.7\%}$
2017			0.14	0.52 (0.37–0.69)	0.38	$F_{39.4\%}$

Banks Peninsula

There have been five fixed site blue cod potting surveys off Banks Peninsula (2002, 2005, 2008, 2012, and 2016), split into geographically separate inshore and offshore areas (Beentjes & Carbines 2003, 2006, 2009; Carbines & Haist 2017; Beentjes & Fenwick 2017). In 2012 and 2016 concurrent random site potting surveys were also carried out and these are intended to replace fixed site surveys as the random surveys provide a more reliable indicator of stock status.

The most recent fixed site inshore survey in 2016 recorded catch rates of 1.26 kg.pot^{-1} (CV 12%), a sex ratio of 67% male, estimated fishing mortality (F) of 1.73 and associated spawner-per-recruit ratio of 4.7% (Table 11). Corresponding values for the 2016 inshore random site survey were 0.53 kg.pot^{-1} (CV 22%), 81% male, $F = 2.1$ and a spawner-per-recruit ratio of 4.3%. For both fixed and random site surveys, the level of exploitation of Banks Peninsula inshore blue cod stocks in 2016 greatly exceeded the Fisheries New Zealand F_{MSY} target reference point of $F_{45\%SPR}$. The very high estimate of total mortality, truncated age composition, strongly skewed sex ratio toward males and extremely low spawner-per-recruit ratio, indicates that the Banks Peninsula inshore blue cod population is heavily overfished. Further, as nearly all females and most males currently caught will be of sub-legal size (less than 30 cm), there is also likely to be significant mortality through catch and return of undersize fish. For the five inshore fixed site surveys there were no trends in survey abundance, length distribution, mean length, or sex ratio. A strong juvenile mode in 2016 can be expected to contribute to increased abundance in about three years when these blue cod recruit to the fishery at 30 cm.

The most recent fixed site offshore survey in 2016 had catch rates of 5.6 kg.pot^{-1} (CV 14%), sex ratio of 65% male, estimated fishing mortality (F) of 0.12 and associated spawner-per-recruit ratio of 40.7% (Table 11). Corresponding values for the 2016 offshore random site survey values were 5.08 kg.pot^{-1} (CV 19%), 57% male, $F=0.05$ and a spawner-per-recruit ratio of 64.3%. For both fixed and random site surveys the level of exploitation (F) of Banks Peninsula offshore blue cod stocks in 2016 is close to or less than the Fisheries New Zealand F_{MSY} target reference point of $F_{45\%SPR}$. The offshore blue cod population, in contrast to inshore, have high catch rates, a wide size range of both males and females, a more balanced sex ratio, and spawner-per-recruit ratio above the target, indicating that they are not overfished. For the five offshore fixed site surveys there were no trends in survey abundance, length distribution, mean length, or sex ratio.

North Canterbury

Kaikoura

There have been four fixed site blue cod potting surveys off Kaikoura (2004, 2007, 2011, and 2015), (Carbines & Beentjes 2006a, 2009; Carbines & Haist 2018; Beentjes & Page 2017). In 2011 and 2015 concurrent random site potting surveys were also carried out and these are intended to replace fixed site surveys in the future as the random surveys provide a more reliable indicator of stock status.

The most recent fixed site survey in 2015 had catch rates of 2.2 kg.pot^{-1} (CV 20%), sex ratio of 66% male, estimated fishing mortality (F) of 0.11 and associated spawner-per-recruit ratio of 53% (Table

BLUE COD (BCO)

11). Corresponding values for the 2015 random site survey were 2.2 kg.pot⁻¹ (CV 19%), 52% male, $F = 0.09$ and a spawner-per-recruit ratio of 58%. For both fixed and random site surveys, the level of exploitation of Kaikoura blue cod stocks in 2015 was less than the Fisheries New Zealand F_{MSY} target reference point of $F_{45\%SPR}$. For the four fixed site surveys, catch rates increased nearly two-fold from 2004 to 2007, and then declined in both 2011 and 2015, with catch rates from the last the lowest of all four surveys (Table 11). Overall blue cod mean length of recruited blue cod (30 cm and over) from fixed sites declined in 2011 and again in 2015. The sex ratio for all blue cod was close to parity for all surveys, with the exception of the 2015 fixed site survey where two-thirds of the blue cod were male. The SB/R ratio calculated for the random survey of 58% is above the target of 45%, suggesting that the population in the survey area is not over-exploited. A strong juvenile mode in 2015 can be expected to contribute to increased abundance in about three to four years when these blue cod recruit to the fishery at 33 cm.

Motunau

There have been four fixed site blue cod potting surveys off Motunau (2005, 2008, 2012, and 2016), (Carbines & Beentjes 2006a, 2009; Carbines & Haist 2018; Beentjes & Sutton 2017). In 2012 and 2016 concurrent random site potting surveys were also carried out and these are intended to replace fixed site surveys in the future as the random surveys provide a more reliable indicator of stock status.

The most recent fixed site survey in 2016 had catch rates of 3.3 kg.pot⁻¹ (CV 13%), sex ratio of 76% male, estimated fishing mortality (F) of 0.62 and associated spawner-per-recruit ratio of 19% (Table 11). Corresponding values for 2016 random site survey were 2.5 kg.pot⁻¹ (CV 27%), 76% male, $F = 0.61$, and a spawner-per-recruit ratio of 19.2%. For both fixed and random site surveys, the level of exploitation of Motunau blue cod stocks in 2016 was greater than the Fisheries New Zealand F_{MSY} target reference point of $F_{45\%SPR}$.

For the four fixed site surveys, catch rates decreased markedly in 2008 and then again in 2016 with a three-fold decline between 2005 and 2016 (Table 11). Overall blue cod mean size steadily declined from 2005 to 2016, with the biggest decreases in 2016. The sex ratio for all blue cod was around 75% male for all fixed site surveys with no trend. A strong juvenile mode in 2015 can be expected to contribute to increased abundance in about three to four years when these blue cod recruit to the fishery at 30 cm. Blue cod abundance and mean size off Motunau has declined and spatial distribution contracted over the eleven years from 2005 to 2016. The very high estimate of total mortality, truncated age composition, strongly skewed sex ratio toward males, and a spawner-per-recruit ratio less than half the target indicates that the blue cod population off Motunau was over-exploited in 2016. Further, as nearly all females and most males currently caught will be of sub-legal size (less than 30 cm), there is also likely to be significant mortality through catch and return of undersize fish.

North Otago

An initial fishery independent fixed site potting survey of blue cod was done in North Otago (also part of BCO 3) in 2005, it produced an overall mean catch rate for all blue cod of 10.14 kg/pot (CV = 5.4%). The catch rate of blue cod 30 cm and over (minimum legal size) was 8.22 kg/pot hour (CV = 5.3%) (Carbines & Beentjes 2006b). In 2009 a second fixed site potting survey (Carbines & Beentjes 2011b) in North Otago produced mean catch rates of blue cod (all sizes) from 6.21 to 19.88 kg per pot per hour. Overall mean catch rate and CV were 11.51 kg per pot per hour and 6.0%, which was consistent with the 2005 survey catch rates. Overall mean catch rate and CV for blue cod 30 cm and over were 8.89 kg per pot per hour and 6.7%, also similar to the 2005 survey results. The overall sex ratio in 2009 was 2.7:1 (male:female), maintaining the bias toward males observed in 2005. Total mortality (Z) for North Otago blue cod populations in 2009 was estimated between 0.25 and 0.36, and were lower than retrospective estimates of Z from the 2005 survey.

In the 2013 North Otago fixed site potting survey (Carbines & Haist 2014b) mean catch rates of blue cod (all sizes) ranged from 2.72 to 8.07 kg per pot per hour. Overall mean catch rate and CV were only 4.96 kg per pot per hour and 12.6%. For blue cod 30 cm and over (minimum legal size), catch rates ranged from 2.02 to 6.42 kg per pot per hour. Overall mean catch rate and CV for blue cod 30

cm and over had dropped to 3.94 kg per pot per hour and 13.7%. The overall sex ratio was 3.3:1 (male:female) and the bias toward males remained consistent for all strata. Z for North Otago blue cod populations in 2013 was estimated between 0.22 and 0.36 and remained consistent with the 2009 survey. The substantial decrease in catch rates in 2013 compared to 2005 and 2009 is of concern. Estimates of Z (0.26, recruitment at 6 years) and percent spawner biomass per recruit ($F_{\%SPR} = 34.11\%$) for the 2013 North Otago fixed site survey are also of some concern.

In the concurrent 2013 North Otago stratified random site potting survey (Carbines & Haist 2014b) mean catch rates of blue cod (all sizes) ranged from 0.94 to 7.46 kg per pot per hour. Overall mean catch rate and CV were 4.16 kg per pot per hour and 13.9%, similar to concurrent fixed sites. For blue cod 30 cm and over, catch rates ranged from 0.46 to 5.28 kg per pot per hour. Overall mean catch rate and CV for blue cod 30 cm and over were 3.01 kg per pot per hour and 14.4%, also similar to fixed sites. The overall sex ratio was 2.13:1 (male:female) and comparatively less bias toward males at random sites.

South Otago

A comparison of fixed and random stratified site potting survey designs was done in three strata off South Otago (also part of BCO 3) in 2009 (Beentjes & Carbines 2011) with similar results. In 2013 a fully stratified random site potting survey of blue cod was done in six strata off South Otago and produced an overall mean catch rate for all blue cod of 6.24 kg/pot (CV = 19.8%) (Carbines & Haist 2014c). The catch rate of blue cod ≥ 30 cm was 5.06 kg/pot hour (CV = 23.03%). The overall sex ratio was 1.22:1 (male:female), with the bias toward males occurring mainly inshore, and some offshore strata having up to 58% females.

Foveaux Strait

A random stratified site potting survey of blue cod was done in Foveaux Strait (also part of BCO 5) in 2010, producing an overall mean catch rate for all blue cod of 4.80 kg/pot (CV = 11.34%). The catch rate of blue cod ≥ 33 cm (minimum legal size) was 2.09 kg/pot hour (CV = 10.87%) (Carbines & Beentjes 2012). In 2014 a second random stratified site potting survey in Foveaux Strait showed a 77% increase in overall mean catch rate of blue cod (all sizes), with an overall mean catch rate and CV of 8.48 kg per pot per hour and 12.85% (Carbines & Haist 2016a). Overall mean catch rate and CV for blue cod ≥ 33 cm had increased 67% to 3.50 kg per pot per hour and 11.26%. The overall sex ratio in 2014 was 0.89:1 (male:female), maintaining the slight bias toward females observed in 2010 (0.86:1).

Paterson Inlet

A fixed site potting survey of blue cod in Paterson Inlet (BCO 5) in 2006 produced an overall mean catch rate for all blue cod of 4.77 kg/pot and CV of 11.9% (set based estimates excluding the marine reserve). The catch rate of blue cod ≥ 33 cm (minimum legal size), was 2.91 kg/pot hour (CV = 12.3%). In 2010 the fixed site survey was repeated along with a concurrent random stratified site survey (Carbines & Haist 2014a). The overall mean catch rate for all blue cod was 4.21 kg/pot and CV of 11.1% from fixed sites, and 0.82 kg/pot and CV of 24.2% from random stratified sites. The overall mean catch rate for ≥ 33 cm blue cod was 3.08 kg/pot and CV of 11.3% from fixed sites, and 0.4 kg/pot and CV of 23.4% from random stratified sites. In 2014 the concurrent fixed site and random stratified site surveys were repeated (Carbines & Haist 2016b). The overall mean catch rate for all blue cod was 4.83 kg/pot and CV of 12.9% from fixed sites, and 1.94 kg/pot and CV of 19.87% from random stratified sites. The overall mean catch rate for ≥ 33 cm blue cod was 2.89 kg/pot and CV of 13.35% from fixed sites, and 1.04 kg/pot and CV of 19.67% from random stratified sites. The fixed site time series from 2006 to 2016 showed extremely stable catch rates in all strata, whereas the random stratified sites overall catch rate had more than doubled from 2010 to 2016. These results suggest that fixed-site catch-rates are hyper stable, and therefore not suited to monitoring blue cod population changes in Paterson Inlet.

Dusky Sound

Three blue cod potting surveys have been carried out in the Dusky Sound. The surveys in 2002 and 2008 were both fixed-site surveys, whereas in 2014, independent fixed-site and random-site surveys were carried out concurrently.

BLUE COD (BCO)

In 2002 the overall mean catch rates for all blue cod from fixed sites were 2.65 kg.pot⁻¹ (CV = 9.2%) and 1.81 kg.pot⁻¹ for recruited blue cod ≥ 33 cm (CV = 8.7%). Catch rates were highest on the open coast (i.e., at the entrance to the Sound; Carbines & Beentjes 2003). The 2008 fixed site survey catch rates were 4.2 kg.pot⁻¹ (CV = 5.8%) for all blue cod and 3.15 kg.pot⁻¹ (CV = 5.9%) for recruited blue cod, considerably higher than in 2002 and again highest catch rates were in the open coast stratum (Carbines & Beentjes 2011a). In 2014 the fixed site catch rates had declined to 3.22 kg.pot⁻¹ (CV = 11.9%) and 2.35 kg.pot⁻¹ (CV = 11.9%), respectively, with highest catch rates on the open coast. The 2014 random site catch rates were less than from fixed sites and were 2.61 kg.pot⁻¹ (CV = 8.6%) for all blue cod and 1.92 kg.pot⁻¹ (CV = 9.6%) for recruited blue cod, also with catch rates highest on the open coast (Beentjes & Page 2017). Overall scaled length and age distributions were similar between the fixed and random site surveys but the sex ratio favoured females in fixed sites (39% male) and was close to parity in random sites (52% male). Fixed site surveys may not be suitable for monitoring the Dusky Sound blue cod population, but at least one more dual fixed and random site survey is required before moving exclusively to random site surveys.

Total mortality (Z) for blue cod from the 2014 random site survey was estimated at 0.25 with Spawner Biomass per Recruit (full recruitment at 8 years for females) estimated at $F_{49\%}$. Mortality estimates from the 2002 and 2008 surveys should not be used due to a recent change in the age determination protocol for blue cod.

Table 11: Summary statistics from standardised blue cod potting surveys of the northeast coast of the South Island (BCO 3). CPUE – catch per unit effort (kg.pot⁻¹); CV – coefficient of variation; Mean length is from population scaled length. . All surveys were reanalysed and reported in Beentjes & Page (2017), Beentjes & Sutton (2017), and Beentjes & Fenwick 2017).

Area/Year	Mean length (cm)		Survey CPUE	CPUE stratum range	Sex ratio (% male)
	Femal	Male			
North Canterbury					
Kaikoura					
2004 (fixed sites)	30.3	32.5	2.62	0.60 – 7.97 (11.1%)	48.7%
2007 (fixed sites)	29.8	32.5	5.0	1.91–20.45 (12.6%)	48.1%
2011 (fixed sites)	27.5	29.1	3.66	2.14 – 11.44 (13.3%)	53.0%
2011 (random sites)	28.5	29.5	2.64	0.61 – 8.22 (16.7%)	46.8%
2015 (fixed sites)	25.7	27.0	2.25	1.58 – 5.07 (20.2%)	66.3%
2015 (random sites)	29.0	30.0	2.21	0.48 – 9.41 (18.9%)	51.7%
Motunau					
2005 (fixed sites)	25.7	29.6	10.2	8.7 – 15.4 (11.4%)	76.6%
2008 (fixed sites)	25.2	29.3	5.5	4.1–8.9 (16.1%)	77.9%
2012 (fixed sites)	24.6	29.1	5.55	4.43–8.70 (11.8%)	71.9%
2012 (random sites)	23.5	28.2	3.01	1.81–6.95 (19.5%)	72.1%
2016 (fixed sites)	22.4	25.8	3.32	2.94–4.66 (12.7%)	75.5%
2016 (random sites)	22.2	26.5	2.48	1.10–7.24 (26.8%)	76.3%
Banks Peninsula					
Inshore					
2002	25.4	28.3	1.12	0.04 – 2.61 (23.2%)	67.9%
2005	27.2	32.7	2.78	1.02 – 4.16 (12.2%)	74.2%
2008	25.5	29.8	1.08	0.07 – 2.3 (17.8%)	70.2%
2012 (fixed sites)	24.7	28.8	1.35	0.60 – 1.88 (12.4%)	67.2%
2012 (random sites)	22.8	27.3	1.23	0.33 – 2.89 (16.6%)	66.1%
2016 (fixed sites)	23.2	26.5	1.26	0.57–2.12 (11.8%)	67.5%
2016 (random sites)	23.8	26.1	0.53	0.09–0.94 (22.2%)	81.3%

Table 11 [Continued]

Offshore					
2002	36.6	37.6	3.39	2.04–4.74 (19.9%)	41.8%
2005	37.4	41.2	6.48	5.68–7.27 (9.4%)	57.2%
2008	35.6	41.8	4.48	3.13 – 5.80 (13.8%)	49.8%
2012 (fixed sites)	33.5	37.4	4.88	3.49 – 6.28 (17.0%)	55.9%
2012 (random sites)	34.1	39.3	3.77	3.69 – 4.09 (36.2%)	59.0%
2016 (fixed sites)	33.6	36.8	5.6	5.09–6.10 (14.1%)	65.2%
2016 (random sites)	36.1	41.3	5.08	5.21–4.54 (19.5%)	57.5%

Table 12: Summary statistics from standardised blue cod potting surveys carried out in the southeast coast of the South Island (BCO 3). CPUE – catch per unit effort (kg.pot⁻¹); CV – coefficient of variation; Mean length, are from population scaled length. North Otago survey - mean length from Beentjes (2012) and Carbines & Haist (2014b), CPUE from Carbines & Beentjes (2006b; 2011) and Carbines & Haist (2014b). South Otago survey - 2009 from Beentjes & Carbines (2011) and 2013 from Carbines & Haist (2014c).

Area/Year	Mean length (cm)		Survey CPUE (kg.pot ⁻¹)	CPUE range (CV) CV is pot-based or set-based*
	Female	Male		
North Otago				
2005 (no stratum 6) (fixed sites)	27.8	32.8	10.1	7.45 - 14.5 (5.4%)
2009 (incl. stratum 6) (fixed sites)	27.4	32.3	11.5	6.21 – 19.88 (*6.8%)
2013 (incl. stratum 6) (fixed sites)	26.9	31.6	5.0	2.72 – 8.07 (*12.6.8%)
2013 (incl. stratum 6) (random sites)	27.6	30.7	4.2	0.94 – 7.46 (*13.9%)
South Otago				
2009** (fixed sites)	29.4	33.6	9.7	3.3–16.9 (*17.1%)
2009 (random sites)	23.7	29.0	4.4	1.2 – 6.0 (*17.8%)
2013 (random sites)	25.5	31.9	6.2	0.8 – 7.4 (*19.9%)

Table 13: Summary statistics from standardised blue cod potting surveys carried out in the south and southwest coast of the South Island (BCO 5). CPUE – catch per unit effort (kg.pot⁻¹); CV – coefficient of variation; Z – Total mortality; $F_{\%SPR}$ estimated for age at full recruitment and $M = 0.14$. Mean length, mean age and Z are from population scaled length and age. Foveaux Strait survey- all results from Carbines & Beentjes 2012, Carbines & Haist 2016a; Paterson Inlet survey -all results from Carbines 2007, Carbines & Haist 2014a, Carbines & Haist 2016b; Dusky Sound - all results from Carbines & Beentjes 2003, 2011a;and Beentjes & Page (2016). Only mean ages, Z estimates and $F_{\%SPR}$ based on otoliths aged with the Age Determination Protocol (Walsh 2017) are included in this table. Results for Paterson Inlet fixed site surveys are not included as they are not reliable.

Area/Year	Mean length		Mean age (years)		CPUE (kg.pot ⁻¹)	CPUE range	Mean Z	$F_{\%SPR}$
	Female	Male	Female	Male				
Foveaux Strait								
2010 (random sites)	27.8	30.5			4.8	1.17 – 14.14		
2014 (random sites)	27.7	30.4			8.5	3.16 – 16.22		
Paterson Inlet								
2006 (fixed sites)	26.9	32.8			4.8	1.47 – 8.42		
2010 (fixed sites)	27.5	32.2			3.2	1.43 – 3.29		
2010 (random sites)	25.9	29.0			0.4	0.22 – 0.53		
Paterson Inlet								
2014 (fixed sites)	26.9	32.3			4.8	1.05 – 7.66		
2014 (random sites)	27.0	29.9			1.94	0.44 – 2.73		

BLUE COD (BCO)

Table 13 [Continued]

Dusky Sound								
2002 (fixed sites)	29.9	34.7			2.65	1.29–8.43		
2008 (fixed sites)	32.2	37.9			4.20	2.49 – 8.13		
2014 (fixed sites)	32.6	35.2	8.1	6.9	3.22	1.87–9.2	0.26 (25%)	48.3%
2014 (random sites)	32.3	33.8	8.2	6.5	2.61	2.04–4.99	0.25 (24%)	49.0%

Trawl survey estimates

Relative abundance indices from trawl surveys are available for BCO 3, BCO 5 and BCO 7, but these have not been used because of the high variance and concerns that this method may not appropriately sample blue cod populations.

4.2 BCO 3




Figure 3: Distribution of landings and number of potlifts for the cod potting method by statistical area and fishing year from trips which landed BCO 3. Circles are proportional within each panel: [catches] largest circle = 95 t in 10/11 for 024; [number potlifts] largest circle = 9641 pots in 05/06 for 024 (Starr & Kendrick in prep).

A standardised CPUE analysis was conducted in 2015 on the target blue cod potting fishery operating in BCO 3. This fishery accounted for two-thirds of the total BCO 3 landings in the 25 years from 1989–90 to 2013–14, predominantly in the two southernmost BCO 3 Statistical Areas: 024 and 026. Together these two areas represented about 90% of the total target blue cod potting fishery over the same 25 years (Figure 3). As found in the previous 2010 analysis, there was misreporting of RCO 3 landings as BCO 3, probably due to data entry errors (Starr & Kendrick 2010). This problem was again resolved before undertaking the CPUE analysis.

The effort data were matched with the landing data at the trip level and the “trip-stratum” stratification inherent in the CELR data was maintained. Two data sets were prepared: one which defined the data set by only selecting trips which fished exclusively in the Areas 018–024 and 026 (designated “statarea”) and the other restricted to trips which exclusively landed BCO 3 (designated “Fishstock”). There was no difference in the CPUE trends estimated by these two data sets. Each analysis was confined to a set of core vessels which had participated consistently in the fishery for a reasonably long period (5 trips in 3 years, resulting in keeping 68 vessels representing 85% of the landings for the “statarea” data set). The explanatory variables offered to the model included fishing year (forced), month, vessel, statistical area, number of pots lifted in a day and number of days fishing in the record. Because there was also an estimated catch of blue cod recorded with nearly every effort record, it was also possible to repeat the standardised analysis based on estimated catch as well as the landed catch. This was done to provide a check on the methods used to groom the

landing data of the spurious RCO 3 landing data. Only a lognormal model based on successful catch records was used as there were too few unsuccessful fishing events to justify pursuing a binomial model.




Figure 4: Comparison of BCO 3 standardised series based on landed greenweight catch data and estimated catch with the three observations from the North Otago potting survey (Starr & Kendrick in prep).

The lognormal standardised model for BCO 3 (Figure 4) showed a declining trend in commercial CPUE from 2002–03 to 2008–09 after a relatively long period of stability, followed by an increasing trend to 2013–14. A model using estimated catches instead of scaled landings showed a similar trend up to 2012–13, when the series based on landed catch increased more rapidly than the estimated catch series. The WG agreed in 2015 that the series based on landed catch was more reliable and consistent with other CPUE analyses done for the Southern Inshore WG.

During the period 2002–03 to 2013–14, commercial catches in all of BCO 3 exceeded the TACC by 5%. As the bulk of the total BCO 3 commercial catch (72%) was taken from Statistical Areas 024 and 026 (along with about 90% of the CPUE data), both the CPUE and catch trends for BCO 3 are strongly influenced by the catches in these areas. Therefore, the Working Group agreed that the CPUE trend presented for the Daily Landed Catch analysis in Figure 4 is representative of the southerly portion of BCO 3 (Areas 024 and 026) and is not applicable to those parts of BCO 3 north of Area 024.

Establishing B_{MSY} compatible reference points

The Working Group accepted mean CPUE from the target BCO cod potting series for the period 1994–95 to 2003–04 as the B_{MSY} -compatible proxy for BCO 3. This period was chosen because catches and CPUE were stable without trend and apparent productivity was good. This period was also used to determine average fishing intensity compatible with the selected B_{MSY} -compatible proxy. The Working Group accepted the default Harvest Strategy Standard definitions for the Soft and Hard Limits at one-half and one-quarter the target, respectively.

4.3 BCO 4

The cod potting fishery in BCO 4 is entirely targeted on blue cod and reported on the daily CELR form. The spatial resolution of the catch effort data is therefore defined by general statistical area, and by day (or part of a day). CPUE was standardised for the cod pot fishery operating in Statistical Areas 049 to 052 (Bentley & Kendrick in prep). The analysis was based on a Weibull model of positive allocated landed catches from a core fleet of vessels. This methodology differs from the previous CPUE standardisation (Kendrick & Bentley 2011) which used a standardisation model with

BLUE COD (BCO)

the assumption of a lognormal error distribution. Detailed examination of model residuals and the distribution of catch per vessel day suggested that the Weibull distribution provided a better fit to the data than the lognormal distribution and other alternative distributions. There appears to have been a change in the underlying frequency distribution of catch categories in the late 1990s, which may be a result of several factors, including changes in the fleet composition, fishing methods, and/or reporting practices. Consequently, the indices for the fishing years up to, and including, 1996/97 are considered to be less reliable, and may not be comparable to, the indices from the latter part of the series.

Overall, the annual indices from the standardisation model have fluctuated without trend since the late 1990s (Figure 5). From 2006–07 to 2012–13 there was a decline in the index, although this was almost fully reversed by a large increase in the index in 2013–14. The indices from the 1990s are lower than those during the latter part of the series and for the aforementioned reasons may not be fully comparable.




Figure 5: Standardised CPUE index for BCO 4 based on records of positive BCO catch by core vessels, 1989–90 to 2013–14 (Bentley & Kendrick in prep.). The indices for the fishing years up to, and including, 1996–97 are considered to be less reliable due to possible changes in fleets, fishing methods and/or reporting practices and may not be comparable to the latter part of the series.

4.4 BCO 5 (Southland)

The first fully quantitative stock assessment for blue cod in BCO 5 was carried out in 2013. A custom-built length-based model, which used Bayesian estimation, was fitted separately to data from Statistical Areas 025, 027 and 030.

4.4.1 Methods

4.4.1.1 Model structure

The stock assessment model is length-based and sex-specific, using growth transition matrices calculated from the von Bertalanffy growth models to transition fish through size bins. This approach is similar to that used for New Zealand rock lobster (Haist et al 2009).

The model is conditioned on the landings for the three modelled fisheries (commercial line, commercial pot, and recreational line), using a Newton-Raphson algorithm to calculate fishing mortality rates for each sex, length bin and fishery. Each fishery is modelled with a selectivity ogive and a retention ogive (Table 14). Catch and catch LFs are a function of the selectivity ogive and landings and landings LFs are a function of the product of selectivity and retention ogives. Separate pre-1993 and post-1992 commercial and recreational fishery retention functions account for the change in minimum legal size (MLS) in 1993. Separate pre-1993 and post-1993 commercial fishery selectivity functions account for change in mesh size regulation at that time, with the assumption that the selectivity change was gradual over 5 years. Discard mortality is assumed for fish that are caught but not landed. Sex change is modelled as a dynamic process, with the proportion of females (at

length) transitioning to males a function of male depletion. S spawning stock biomass (SSB) is measured as the total mature biomass.

A Beverton-Holt stock recruitment relationship is assumed. The standard deviation of recruitment residuals (log-scale) is fixed at 0.6 and the steepness prior is beta distributed (mean= 0.75, std. dev.=0.10). Recruitment residuals are estimated for 1980 to 2010. Fish recruit to the model at age 0+ with 65% of fish recruiting as females.

Natural mortality is modelled assuming a normal prior distribution with a mean of 0.14 and a standard deviation of 0.015. The majority of the prior density is in the range of 0.11 to 0.17, which is the range of uncertainty considered in blue cod potting survey analyses (Beentjes & Francis, 2011).

The populations are initialised at unexploited equilibrium conditions in 1900.

The assumed prior distributions for model parameters are given in Table 15.

Table 14: Model selectivity and retention ogives by fishery, their parametric form, and parameter values if fixed or data fitted in the model to inform their estimation. DHN = double half normal.

Ogives	Type	Parameters if fixed or data to inform
<u>Selectivity</u>		
Commercial line fishery	Logistic	50% selected at 280 mm; 95% selected at 305 mm
Commercial pot fishery <=1992	DHN	Mesh size trial LF
Commercial pot fishery >=1997	Logistic	Logbook & Shed sampling LF
Recreational fishery	DHN	Recreational catch LF
Survey	DHN	Survey LF
<u>Retention</u>		
Commercial line fishery	Knife-edge	MLS (330 mm)
Commercial pot fishery <=1992	Knife-edge	MLS
Commercial pot fishery >=1993	Knife-edge	MLS
Recreational fishery <=1992	Logistic	Recreational landings LF
Recreational fishery >=1993	Logistic	Shifted +3 cm from <=1992 retention curve

Table 15: Assumed prior distributions for model parameters.

Model parameters	Distribution	Parameters
M	Normal	Mean: 0.14 Std. dev: 0.015
S-R steepness	Beta (defined on 0.2 – 1.0)	Mean: 0.75 Std. dev: 0.10
Recruitment variation	Normal-log	Std. dev: 0.60
1995 sex-change d_{max}	Normal-log	Mean: ln(410) Std. dev: 0.05

4.4.1.2 Data

Separate data sets were compiled and analysed for Statistical Areas 025, 027, and 030. The data available for each of these areas differs, and little data are available for the remainder of the BCO 5 Statistical Areas. Combined, Statistical Areas 025, 027 and 030 represent 92% of the recent commercial fishery landings. The general categories of data used in the stock assessment models include: catch and landings; fishery and survey length frequency data (LFs); abundance indices; and biological information on growth, maturation, and sex change.

Historical time series of BCO 5 landings were constructed for three gear types: commercial hand line fishing, commercial pot fishing, and recreational fishing. Additionally, non-reported blue cod catch used as bait in the CRA 8 rock lobster fishery was estimated and included with the commercial landings, and customary catch estimates were included with the recreational harvest.

Commercial landings data are available beginning in 1931 (Warren et al 1997) and these were linearly decreased back to 1900, when the fishery was assumed to begin. The 1989–90 to 2011–12 average proportion of the total BCO 5 catch in each Statistical Area was used to prorate the earlier landings estimates to Statistical Area. A time series of non-reported blue cod used as bait in the rock

BLUE COD (BCO)

lobster fishery was developed based on a 1985 diary study (Warren et al 1997) in conjunction with CRA 8 rock lobster landings.

A time series of recreational blue cod harvest was developed based on the 1991–92 and 1996 diary survey estimates of BCO 5 recreational catch. The average blue cod catch per Southland resident was estimated from the survey data, and assuming a constant per capita catch rate extrapolated to a time series using Southland District population census data.

Commercial fishery LF data were collected through a commercial fishers' logbook project and a shed sampling project from 2009–2011. The shed sampling was sex-specific while the logbook sampling was not. It is unclear whether samples collected for shed sampling were of the entire catch or of landings. Mean size of fish from the shed samples were smaller than those from the logbook programme (for Areas 025 and 027; there were not shed samples from Area 030), which may have resulted because the shed samples were not representative of the entire fishing area. The shed and logbook LF data are each fitted to model predictions of the average commercial catch size distribution for 2009 to 2011.

Recreational fishery LFs were obtained from a 2009–10 study of the Southland recreational blue cod fishery (Davey & Hartill 2011). This study included a boat ramp survey (Bluff, Riverton/Colac, and Halfmoon Bay) and a logbook survey of charter and recreational vessels. Blue cod measured through the boat ramp programme were assumed to represent the landings and fish measured through the logbook programme were assumed to represent the catch.

Length frequency data from a blue cod mesh size selectivity study, conducted by MAF in 1986 at Bluff and Stewart Island, were available. The Length Frequencies from pots fitted with the then-standard 38 mm mesh were assumed to represent the size composition of the BCO 5 commercial pot fishery catch prior to the 1992 and 1994 pot regulation changes. In the model, this data is fitted to the predicted average size distribution of the 1985–1992 potting fishery.

Length Frequency data is also available from random stratified potting surveys conducted in Areas 025 and 030 in 2010. These surveys provide not only length frequency data, but also are one of the few information sources about the population sex structure. These data are fitted in the model assuming domed survey selectivity.

Three sets of data are available that can inform stock abundance estimates: fishery-based standardised CPUE estimates (Table 16), survey-based estimates of total mortality (Z), and a drift underwater video survey (DUV) estimate of absolute stock abundance.

Z estimates were derived from the 2010 Area 025 and Area 030 random-stratified potting survey data using standard methods described in Beentjes & Francis (2011). The distributions of Z estimates are approximately lognormal and are fitted with lognormal priors in the stock assessment model. The mean Z estimate for Area 030 (0.377) is slightly lower than that for Area 025 (0.465).

A DUV survey was conducted in Area 025 in 2010, surveying a number of the random-stratified sites that were sampled during the potting survey. The survey estimate of the mean density of legal-sized blue cod was extrapolated to the total Area 025 area to generate a total abundance estimate. This was fitted to model-predicted 2010 legal-sized blue cod abundance.

The data fitted in the models for each Statistical Area are shown in Table 17 and the assumed error structure of each data series is shown in Table 18.

4.4.1.3 Further assumptions

Sex-specific von Bertalanffy growth parameters are available from Area 025 and Area 030 random-stratified potting surveys. The Area 025 growth models were assumed for Area 027. Both male and female blue cod are assumed to mature at a length of 280 mm (Carbines 2004b).

Sex-change data was available from a 1995 Foveaux Strait study that characterised blue cod by state: male, female, or transitional (Carbines 2004b). The proportions of transitional females by length bin were fitted with a parametric relationship to describe the sex-change process. The maximum proportion transitional was observed at 410 mm.

Assuming that sex-change is a function of the relative abundance of mature males was found to result in fewest model convergence issues. The length at 50% sex change (d_{max}) is modelled as a function of the ratio of mature male biomass in year y (B_y^M) relative to mature male biomass in the virgin state (B_0^M):

$$d_{max} = \lambda \left(\frac{B_y^M}{B_0^M} \right)^\delta$$

where the parameters λ and δ are estimated through the model fitting. In practice, only λ was estimated and δ was fixed. This model results in the form of the sex-change relationship remaining the same except that it is shifted along the length-axis. With this parameterisation it is not possible to fix the 1995 length at 50% sex change (to 410 mm, as observed in the sex transition data set collected in 1995), so a penalty function is used to encourage that value.

Table 16: Standardised CPUE indices for Statistical Areas 025, 027 and 030.

Fishing Year	Area 025	Area 027	Area 030
1990	0.803	0.603	0.925
1991	0.748	0.607	0.860
1992	0.815	0.665	1.026
1993	0.854	0.835	0.846
1994	0.847	0.648	0.689
1995	0.808	0.796	0.669
1996	0.943	1.022	0.657
1997	1.043	1.241	1.011
1998	1.084	1.116	1.141
1999	0.972	1.152	1.224
2000	1.034	1.292	1.185
2001	1.143	1.466	1.098
2002	1.160	1.743	1.453
2003	1.256	1.532	1.422
2004	1.145	1.602	1.359
2005	1.283	1.219	1.262
2006	1.253	1.127	1.172
2007	1.035	0.881	1.093
2008	1.017	0.888	0.924
2009	1.023	0.894	0.939
2010	0.984	0.901	0.961
2011	1.006	0.888	0.839
2012	0.998	0.940	0.819

Table 17: Data series fitted in the stock assessments for Areas 025, 027, and 030.

Data type	Series	Area 025	Area 027	Area 030
LF data:	Shed	✓	✓	-
	Logbook	✓	✓	✓
	Survey	✓	-	✓
	Mesh sel. trials	data common to all areas		
	Rec. landings	data common to all areas		
	Rec. catch	data common to all areas		
Abundance Index:	CPUE	✓	✓	✓
	Survey Z	✓	-	✓
	DUV abundance	✓	-	-

BLUE COD (BCO)

Table 18: Assumed distributions for data fitted in the models.

Data type	Distribution	Parameters
Logbook LF	Multinomial	N: 100
Shed samples LF	Multinomial	N: 100
Mesh size trials LF	Multinomial	N: 100
Recreational catch LF	Multinomial	N: 100
Recreational landings LF	Multinomial	N: 100
Survey LF	Multinomial	N: 100
CPUE	Normal-log	Std. dev: 0.20
Survey Z –Area 025	Normal-log	Mean: -0.782 Std. dev: 0.178
Survey Z –Area 030	Normal-log	Mean: -0.991 Std. dev: 0.173
DUV LegalN	Normal-log	Mean: 15.163 Std. dev: 0.300

4.4.1.4 Calculation of fishing intensity and B_{MSY}

Fishing intensity is measured as the spawning biomass per recruit (SPR). $F_{\%SPR}$ is the ratio of spawning biomass per recruit at a given level of fishing mortality relative to the spawning biomass per recruit in the absence of fishing. This metric was selected to represent fishing intensity because estimates for the entire BCO 5 stock can readily be calculated from the Statistical Area estimates.

MSY statistics are calculated assuming deterministic recruitment and the final years' selectivity and retention ogives. The recreational and customary fisheries are held fixed at the current levels, and only the commercial fishery varied to determine MSY. B_{MSY} is measured as total mature biomass and MSY is presented as the commercial catch at B_{MSY} .

Caution about the interpretation of B_{MSY} estimates

There are several reasons why B_{MSY} , as calculated in this way, is not a suitable target for management of blue cod fisheries. First, it assumes a harvest strategy that is unrealistic in that it involves perfect knowledge (current biomass must be known exactly in order to calculate the target catch) and annual changes in TACC (which are unlikely to happen in New Zealand and not desirable for most stakeholders). Second, it assumes perfect knowledge of the stock-recruit relationship, which is actually very poorly known. Third, it makes no allowance for extended periods of low recruitment. Fourth, it would be very difficult with such a low biomass target to avoid the biomass occasionally falling below 20% B_0 , the default soft limit according to the Harvest Strategy Standard.

4.4.1.5 Biomass estimates

The assessment was conducted in two steps. First, a set of initial exploratory model runs was carried out generating point estimates (MPD runs, which estimate the mode of the posterior distribution). Their purpose was to decide which sets of assumptions should be carried forward to the final runs. The final runs were fully Bayesian, estimating posterior distributions for all quantities of interest.

The modelling assumptions and approaches investigated through the exploratory model runs included: the dynamics of sex-change; what assumptions to make about LF data from the logbook and shed sampling programmes; the magnitude of recruitment variation; the magnitude of error in fits to the CPUE data; the form of the survey and recreational fishery selectivity; and sensitivity to alternative assumptions about recreational catch, bait usage, and discard mortality rates.

Four final runs were chosen by the Working Group: a *base case* and three sensitivities to the *base case*. The sensitivity runs each modify a single assumption of the *base case*. The sex-change power parameter (delta in equation above) is fixed at 0.4 for the *base case*. Two of the sensitivity runs modify this parameter to values of 0.2 and 0.6. The third sensitivity run reduces the recreational catch time series by 50%.

Label	Description
1.1	Base case
1.2	Sex-change power parameter=0.2
1.3	Sex-change power parameter=0.6
1.4	Recreational catch reduced by 50%

Bayesian posterior distributions were estimated for each of these runs using a Markov Chain Monte Carlo (MCMC) approach. For each run a chain of 1 million was completed and the chains thinned to produce a posterior sample of 1000. BCO 5 summary statistics are calculated summing across Areas 025, 027, and 030. B_{MSY} and MSY are calculated assuming these areas account for 92% of the BCO 5 stock.

The model estimates are summarised in Table 19 (estimates of spawning biomass and MSY), Figure 6 (biomass trajectories), Figure 7 (fishing intensity trajectories), and Figure 8 (recruitment trajectories).

The runs with the higher sex-change power parameter (run 1.3) have higher male and lower female spawning abundance in the unfished populations and runs with the lower sex-change power parameter (run 1.2) have lower male and higher female initial abundance. Current biomass and the combined male and female B_0 do not differ much among the runs. Assuming lower recreational catch (run 1.4) results in a slightly lower B_0 estimate and slightly higher current biomass. Area 025 is somewhat more depleted than Areas 027 and 030.




Figure 6: Median estimates of Area 025, Area 027, Area 030, and Areas combined male and female spawning biomass for the base case and sensitivity runs, 1900 – 2012.

BLUE COD (BCO)




Figure 7: Fishing intensity ($F\%_{SPR}$) estimates from the base case runs for Areas 025, 027, 030, and the Areas combined, 1900–2012. The horizontal lines show the median and the vertical lines show the 90% confidence intervals.




Figure 8: Recruitment estimates from the base case runs for Areas 025, 027, 030, and the Areas combined, 1980–2010. The boxes show the interquartile range, the whiskers show the 90% confidence limits, and the bars show the medians.

Fishing intensity has remained below $F_{40\%SPR}$, except in Area 025 for a brief period in the 1990s. Recruitment has been slightly below average in all three Areas over the last decade.

Table 19: Estimates of BCO 5 spawning stock biomass, MSY and B_{MSY} for final runs (medians of marginal posterior distributions, with 90% confidence intervals in parentheses). B_0 and MSY are calculated assuming Areas 025, 027 and 030 represent 92% of the BCO 5 blue cod stock.

Run	B_0 (000 t)	$B_{current}$ (% B_0)	MSY	B_{MSY} (% B_0)
1.1	28(25,31)	39(31,51)	1336(1092,1589)	31(29,35)
1.2	28(26,31)	39(30,50)	1316(1088,1569)	32(29,35)
1.3	27(24,31)	39(30,50)	1345(1114,1607)	31(28,34)
1.4	26(24,29)	40(31,51)	1335(1115,1615)	31(29,35)

4.4.1.6 Yield estimates and projections

Ten-year stock projections were conducted for the three Statistical Areas at constant catch levels, with summary statistics calculated at the end of 5 and 10 years.

Commercial catch levels were based on the current TACC and the average BCO 5 Statistical Area catch split over the past 10 years. Although only 90% of the BCO 5 TACC was caught on average over the past 10 years, with the reduction of the TACC to 1239 t in 2011–12, over 98% of the allowable catch was caught that year. Therefore stock projections based on the full TACC being caught appears reasonable. Alternative catch scenarios were simulated with commercial catch increased and reduced by 20%. Recreational and customary catch was assumed to remain constant at the 2011–12 levels.

Recruitment was simulated by randomly re-sampling (with replacement) from the time series of recruitment deviations, applied to the stock-recruitment relationship. Two alternative recruitment scenarios were simulated: recent recruitments were re-sampled from the 2001–2010 recruitment deviations and long-term recruitments were re-sampled from the 1980–2010 recruitments. Summary statistics were calculated for the BCO 5 FMA by summing B_0 , B_{msy} and projection biomass estimates across the three Statistical Areas.

The projections indicate that under the assumptions of commercial catch at the current TACC and recruitment at recent levels the BCO 5 biomass is unlikely to change much over the next 10 years (Figure 9). Recruitments closer to the long-term average or a reduction in catch from the current TACC results in slight increases in biomass and an increase in catch above the TACC results in a slight decrease in biomass. Although the spawning stock sex ratio is variable among the sensitivity trials, by 2013 and through the projection period the sex ratio remains relatively constant (Table 20).

The probabilities of the projected spawning stock biomass (2018 and 2023) being below the hard limit of 10% B_0 , the soft limit of 20% B_0 , the target of 40% B_0 , and 25%, 50% and 100% of B_{MSY} are presented in Table 21, for the base case model with recent or long-term recruitment and three catch levels and for the sensitivity runs with recent recruitment and commercial catch at the current TACC. With catches at the current TACC, the probability of the stock being less than either the soft or hard limit over the next five years is negligible.

There are no time series of length frequency observations for the BCO 5 stock assessment. So, while the assessment indicates a BCO 5 recruitment pulse in the early 1990s, the information to support this pulse comes solely from the CPUE data, and hence may be spurious.

The sex change predictions also need to be viewed with caution as there are few data to inform the parameters and the form of the equation.

BLUE COD (BCO)

Table 20: Median estimates of the proportion male in the 1900, 2013, 2018 and 2023 BCO 5 spawning stock at alternative recruitment and catch levels for the base case and sensitivity stock projections.

Run	1.1						1.2	1.3	1.4
Recruitment	Recent	Recent	Recent	Long-term	Long-term	Long-term	Recent	Recent	Recent
Catch Level	TACC	1.2·TACC	0.8·TACC	TACC	1.2·TACC	0.8·TACC	TACC	TACC	TACC
1900	0.41	0.41	0.41	0.41	0.41	0.41	0.47	0.39	0.41
2013	0.51	0.51	0.51	0.51	0.51	0.51	0.49	0.51	0.51
2018	0.48	0.51	0.51	0.47	0.51	0.51	0.50	0.48	0.49
2023	0.51	0.52	0.49	0.49	0.51	0.48	0.49	0.52	0.51




Figure 9: Projected BCO 5 spawning biomass (%B₀) assuming recent or long-term recruitment and catch at current TACC or increased/decreased by 20% for the base case run. Median estimates are shown as solid lines and 90% confidence intervals as shaded polygons.

Table 21: Probabilities of SSB being below B₀ and B_{msy} reference levels in 2013, 2018 and 2023 at alternative recruitment and catch levels for the *base case* and sensitivity stock projections.

Run	1.1						1.2	1.3	1.4
Recruitment	Recent	Recent	Recent	Long-term	Long-term	Long-term	Recent	Recent	Recent
Catch Level	TACC	1.2·TACC	0.8·TACC	TACC	1.2·TACC	0.8·TACC	TACC	TACC	TACC
P(B ₂₀₁₃ < 0.1 B ₀)	0	0	0	0	0	0	0	0	0
P(B ₂₀₁₃ < 0.2 B ₀)	0	0	0	0	0	0	0	0	0
P(B ₂₀₁₃ < 0.4 B ₀)	0.538	0.538	0.538	0.538	0.538	0.538	0.576	0.549	0.532
P(B ₂₀₁₃ < 0.25 B _{msy})	0	0	0	0	0	0	0	0	0
P(B ₂₀₁₃ < 0.5 B _{msy})	0	0	0	0	0	0	0	0	0
P(B ₂₀₁₃ < B _{msy})	0.095	0.095	0.095	0.095	0.095	0.095	0.116	0.091	0.078
P(B ₂₀₁₈ < 0.1 B ₀)	0.001	0.002	0	0	0.001	0	0	0	0
P(B ₂₀₁₈ < 0.2 B ₀)	0.010	0.048	0.002	0.003	0.024	0	0.012	0.007	0.015
P(B ₂₀₁₈ < 0.4 B ₀)	0.543	0.694	0.379	0.470	0.622	0.288	0.578	0.578	0.605

Table 21 [Continued]

$P(B_{2018} < 0.25 B_{msy})$	0	0.002	0	0	0	0	0	0	0
$P(B_{2018} < 0.5 B_{msy})$	0.002	0.014	0	0	0.006	0	0.004	0.002	0.005
$P(B_{2018} < B_{msy})$	0.230	0.377	0.114	0.153	0.294	0.069	0.249	0.215	0.262
$P(B_{2023} < 0.1 B_0)$	0.003	0.024	0.002	0	0.005	0	0.007	0.004	0.006
$P(B_{2023} < 0.2 B_0)$	0.053	0.173	0.008	0.019	0.077	0	0.052	0.051	0.074
$P(B_{2023} < 0.4 B_0)$	0.498	0.681	0.271	0.289	0.533	0.110	0.491	0.505	0.553
$P(B_{2023} < 0.25 B_{msy})$	0.001	0.014	0	0	0.002	0	0.004	0.003	0.002
$P(B_{2023} < 0.5 B_{msy})$	0.021	0.107	0.004	0.009	0.037	0	0.025	0.018	0.040
$P(B_{2023} < B_{msy})$	0.256	0.473	0.105	0.113	0.306	0.030	0.272	0.257	0.305

4.5 Other factors

The target blue cod fishery is chiefly a pot fishery and there are few significant bycatch problems. However, in recent years bycatch associated with the inshore fleet of trawlers has increased in BCO 3 and BCO 7. Blue cod is only a very minor bycatch of the offshore fleet.

Before the introduction of the QMS, blue cod landings were affected by factory limits imposed in some parts of Southland, and there were economic constraints to the development of the fishery at the Chatham Islands (BCO 4).

Blue cod fishing patterns have been strongly influenced by the development and subsequent fluctuations in the rock lobster fishery, especially in the Chatham Islands, Southland and Otago. Once a labour intensive handline fishery, blue cod are now taken mostly by cod pots. The fishery had decreased in the past; however, with the advent of cod pots it rapidly redeveloped. Large areas are currently not heavily fished and there are some areas such as the Mernoo Bank, the Puysegur Bank and South Traps which are potentially productive fisheries. Anecdotal information from recreational fishers suggests that there is local depletion in some parts of BCO 3, BCO 5 and BCO 7 where fishing has been concentrated. Blue cod abundance (Carbines & Cole 2009), catch (Cranfield et al 2001) and productivity (Jiang & Carbines 2002, Carbines et al 2004) may also be affected by disturbance of benthic habitat.

5. STATUS OF THE STOCKS

For BCO 1 and 8 recent commercial catch levels are considered sustainable. The status of the remaining fishstocks is summarised below.

- **BCO 3 (Statistical Areas 024 and 026)**

Stock Structure Assumptions


Tagging experiments suggest that blue cod populations may be isolated from each other and there may be several distinct populations within management areas. For the purposes of this summary, BCO 3 is split into two sub-areas along the Statistical Area 022/024 boundary.

Stock Status	
Year of Most Recent Assessment	2015 (CPUE analysis)
Assessment Runs Presented	Standardised CPUE index based on landed catch of BCO target pot fishery
Reference Points	Target: B_{msy} proxy based on mean CPUE for the period 1994–95 to 2003–04 Soft Limit: 50% B_{MSY} Hard Limit: 25% B_{MSY} Overfishing Threshold: F_{MSY} proxy based on mean relative


BLUE COD (BCO)

	exploitation rate for the period 1994–95 to 2003–04
Status in relation to Target	About as Likely as Not (40–60%) to be at or above
Status in relation to Limits	Soft Limit: Unlikely (< 40%) to be below Hard Limit: Very Unlikely (< 10%) to be below
Status in relation to Overfishing	Overfishing is About as Likely as Not (40–60 %) to be occurring

Historical Stock Status Trajectory and Current Status



Cod-potting CPUE index (CP-landed), along with catches and TACC for BCO 3.



Relative Fishing Intensity (catch/CPUE) for BCO 3 (where CPUE=CP(land) and catch=Sum(024 & 026)).

Fishery and Stock Trends

Recent Trend in Biomass or Proxy	Biomass has increased in four of the five years since a nadir reached in 2008–09. It is now near the highest level in the series.
Recent Trend in Fishing Intensity or Proxy	Relative exploitation rate declined since 2011–12, and 2013–14 was below the overfishing threshold.
Other Abundance Indices	The North Otago potting survey has only three index values which do not form a trend and do not match the CP CPUE series very well. The South Otago potting survey has only two index values.

Trends in Other Relevant Indicators or Variables	-
--	---

Projections and Prognosis	
Stock Projections or Prognosis	Stock abundance, as monitored with cod potting CPUE, has fluctuated around a mean level since the early 1990s at levels of commercial catch averaging near 160 t/year. Recreational catch trends are not well known, but there seems to be little cause for concern as long as catches remain near current levels.
Probability of Current Catch or TACC causing decline Biomass to remain below or to decline below Limits	Soft Limit: Unlikely (< 40%) Hard Limit: Very Unlikely (< 10%)
Probability of Current Catch causing Overfishing to continue or to commence	-

Assessment Methodology and Evaluation		
Assessment Type	Level 2: Partial Quantitative Stock Assessment	
Assessment Method	Standardised CPUE analysis of a target cod-potting fishery	
Main data inputs	Catch and effort data derived from the MPI catch reporting data.	
Period of Assessment	Latest assessment: 2015	Next assessment: unknown
Overall Assessment Quality	1 – High Quality	
Main Data Inputs (Rank)	- Catch and effort data	1 – High Quality
Data not used	- North and South Otago potting surveys	3 – Low Quality: insufficient data points to describe trends and inconsistencies with BCO ageing have reduced the quality of age-based mortality estimates
Changes to Model Structure and Assumptions	-	
Major Sources of Uncertainty	-	

Qualifying Comments
As the bulk of the commercial catch (72%) is taken from Statistical Areas 024 and 026, both CPUE and catch trends for BCO 3 are strongly influenced by catches in these areas. A June 2009 change in regulations governing commercial pots (change from 38 mm mesh to 48 mm square grids) will have affected CPUE indices.

Fishery Interactions
Over two thirds of BCO 3 commercial catches are taken in a target cod-potting fishery which has very little interaction with other species. Most of the remaining BCO 3 catch is taken in the inshore bottom trawl fishery operating on the east coast of the South Island, largely directed at flatfish, red cod and tarakihi.

BLUE COD (BCO)


BCO 4

Stock Structure Assumptions


For the purposes of this summary BCO 4 is considered to be a single management unit.

Stock Status	
Year of Most Recent Assessment	2015
Assessment Runs Presented	CPUE index based on landed catch
Reference Points	Interim Target: B_{MSY} proxy based on mean CPUE for the period 2002–03 to 2013–14 (a period with high yield when both catch and CPUE were stable) Soft Limit: 50% B_{MSY} proxy Hard Limit: 25% B_{MSY} proxy Overfishing threshold: F_{MSY} proxy based on mean relative exploitation rate for the period 2002–03 to 2013–14
Status in relation to Target	About as Likely as Not (40–60%) to be at or above the target
Status in relation to Limits	Soft Limit: Very Unlikely (< 10%) to be below Hard Limit: Very Unlikely (< 10%) to be below
Status in relation to Overfishing	About as Likely as Not (40–60 %) to be occurring

Historical Stock Status Trajectory and Current Status



BCO 4 standardised CPUE plotted as two series: 1990–1997 and 1998–2014, representing greater confidence in the latter series. Also plotted are the QMR/MHR landings and the BCO 4 TACC. The orange line represents the interim B_{MSY} proxy of mean CPUE from 2003–2014. The purple line is the interim Soft Limit= $0.5*[B_{MSY} \text{ proxy}]$ and the grey line is the interim Hard Limit= $0.25*[B_{MSY} \text{ proxy}]$.



BCO 4 fishing intensity (=catch/CPUE) plot based on the standardised CPUE series and the QMR/MHR landings. Horizontal orange line represents the mean 2003–2014 fishing intensity associated with the interim B_{MSY} proxy.

Fishery and Stock Trends

Recent Trend in Biomass or Proxy	CPUE has fluctuated without trend since 1997–98
Recent Trend in Fishing Intensity or Proxy	Relative exploitation rate has declined since 2010–11 and in 2013–14 was below the overfishing threshold
Other Abundance Indices	-
Trends in Other Relevant Indicators or Variables	-

Projections and Prognosis

Stock Projections or Prognosis	The current catch and TACC are Unlikely (< 40%) to cause the stock to decline
Probability of Current Catch or TACC causing Biomass to remain below or to decline below Limits	Soft Limit: Very Unlikely (< 10%) Hard Limit: Very Unlikely (< 10%)
Probability of Current Catch or TACC causing overfishing to continue or to commence	-

Assessment Methodology and Evaluation

Assessment Type	Level 2: Partial Quantitative Stock Assessment	
Assessment Method	Fishery characterisation and standardised CPUE analysis	
Assessment Dates	Latest assessment: 2015	Next assessment: 2019
Overall assessment quality rank	1 – High Quality	
Main data inputs (rank)	- Catch and Effort 1997–98 to 2013–14 - Catch and Effort 1989–90 to 1996–97	1 – High Quality 2 – Medium or Mixed Quality: compromised by changes in fleet composition and reporting practices
Data not used (rank)	N/A	
Changes to Model Structure and Assumptions	-	
Major Sources of Uncertainty	-	

BLUE COD (BCO)

Qualifying Comments
-
Fishery Interactions
The catch is almost entirely taken by target cod potting and there is little interaction with other species.


• BCO 5

Stock Structure Assumptions

Tagging experiments suggest that blue cod populations may be isolated from each other and there may be several distinct populations within management areas. For the purposes of this summary, blue cod in Statistical Areas 025, 027 and 030 of BCO 5 are treated as a unit stock. Dusky Sound and Patterson Inlet are assumed to contain discreet populations of BCO, which are monitored with potting surveys.

Stock Status	
Year of Most Recent Assessment	2013
Assessment Runs Presented	One base case model was used to evaluate BCO 5 stock status in this assessment. Three sensitivity runs are also presented.
Reference Points	Interim Management Target: 40% B_0 Soft Limit: 20% B_0 Hard Limit: 10% B_0 Overfishing threshold: F_{MSY}
Status in relation to Target	B_{2013} was estimated to be 39.4% of B_0 ; About as Likely as Not (40–60%) to be at or above the Interim Management Target
Status in relation to Limits	B_{2013} is Very Unlikely (< 10%) to be below the Soft Limit and Exceptionally Unlikely (< 1%) to be below the Hard Limit
Status in relation to Overfishing	Unlikely (< 40%) that overfishing is occurring

Historical Stock Status Trajectory and Current Status



Trajectory of fishing intensity (F_{SPR}) and spawning biomass ($\%B_0$) for BCO 5 from the start of the assessment period in 1990 to 2012. The vertical lines at 10% B_0 , 20% B_0 and 40% B_0 represent the soft limit, the hard limit and the target, respectively, and the shaded area shows the B_{MSY} 90% CI. Estimates are based on MCMC medians and the 2012 90% CI is shown by the crossed lines.

Fishery and Stock Trends	
Recent Trend in Biomass or Proxy	Biomass has been slowly decreasing since 2000.
Recent Trend in Fishing Intensity or	Fishing intensity is estimated to have been relatively constant

BLUE COD (BCO)

Proxy	since 2000.
Other Abundance Indices	-
Trends in Other Relevant Indicators or Variables	Recent recruitment (2002 – 2010) is estimated to be slightly below the long-term average.

Projections and Prognosis	
Stock Projections or Prognosis	BCO 5 biomass is expected to stay steady over the next 5 to 10 years at the 2012 TACC which approximates the 2012 catch.
Probability of Current Catch or TACC causing Biomass to remain below or to decline below Limits	Soft Limit: Very Unlikely (< 10%) Hard Limit: Very Unlikely (< 10%)
Probability of Current Catch or TACC causing Overfishing to continue or to commence	Very Unlikely (< 10%)

Assessment Methodology and Evaluation		
Assessment Type	Level 1 - Full Quantitative Stock Assessment	
Assessment Method	Length-based model with Bayesian estimation of posterior distributions	
Assessment Dates	Latest assessment: 2013	Next assessment: 2019
Overall assessment quality rank	1 – High Quality	
Main data inputs (rank)	<ul style="list-style-type: none"> - CPUE time series - Proportion at length data from surveys and commercial catch - Estimates of biological parameters - DUV survey absolute biomass estimate - Potting survey Z estimates 	1 – High Quality 1 – High Quality 1 – High Quality 1 – High Quality 1 – High Quality
Data not used (rank)	-	-
Changes to Model Structure and Assumptions	New model	
Major Sources of Uncertainty	Degree to which CPUE reflects abundance; the age, size and sex structure of the population; relationship between abundance and sex change dynamics	

Qualifying Comments
-

Fishery Interactions
Historically, significant quantities of blue cod, taken by potting, were used as bait in the commercial rock lobster fishery. Since 1996, reporting of blue cod used for bait is mandatory and included as part of the commercial catch reporting. Some blue cod are landed as bycatch in rock lobster pots and oyster dredges.

Research needs
Research into the sex change dynamics of blue cod would assist in improving the information that goes into the BCO 5 stock assessment. Histological analysis of gonads from the randomly stratified surveys would be a useful approach to assess sex change dynamics. Catch sampling should be undertaken in BCO 5 and needs to be scheduled as part of the medium term research plan.

BLUE COD (BCO)

BCO 7 - Marlborough Sounds only

Stock Structure Assumptions

For the purposes of this summary BCO - Marlborough Sounds is considered to be a single management unit.

Stock Status	
Year of Most Recent Assessment	2018
Assessment Runs Presented	Catch rates from the fixed and random site Marlborough Sounds potting surveys
Reference Points	Target1: B_{MSY} -compatible proxy based on the Marlborough Sounds potting survey (to be determined) Target 2: Target: $F_{SB45\%}$ ($F_{SB45\%} = 0.26$). Soft Limit: 20% B_0 Hard Limit: 10% B_0 Overfishing threshold: $F_{SB45\%}$
Status in relation to Target	F is Unlikely ($< 40\%$) to be at or below the target
Status in relation to Limits	Unknown
Status in relation to Overfishing	Overfishing is Likely ($> 60\%$) to be occurring
Historical Stock Status Trajectory and Current Status	
<div><div><div><div><div>Marlborough Sounds surveys</div><div><div><div>Queen Charlotte Sound</div><div><div><div><div><div></div><div></div></div><div><div><div>Fixed</div><div>Random</div></div></div><div><div><div>1995</div><div>1996</div><div>2001</div><div>2004</div><div>2007</div><div>2010</div><div>2013</div><div>2017</div></div><div><div>2.1</div><div>1.3</div><div>1.2</div><div>1.1</div><div>2.1</div><div>1.0</div><div>0.5</div><div>1.0</div></div></div></div><div><div>Cook Strait</div><div><div><div><div></div><div></div></div><div><div><div>Fixed</div><div>Random</div></div></div><div><div><div>2008</div><div>2010</div><div>2013</div><div>2017</div></div><div><div>1.6</div><div>1.1</div><div>0.7</div><div>1.1</div></div></div></div></div></div><div><div><div><div><div>Pelorus Sound</div><div><div><div><div></div><div></div></div><div><div><div>Fixed</div><div>Random</div></div></div><div><div><div>1995</div><div>1996</div><div>2001</div><div>2004</div><div>2007</div><div>2010</div><div>2013</div><div>2017</div></div><div><div>2.4</div><div>0.7</div><div>1.2</div><div>1.1</div><div>2.9</div><div>2.0</div><div>1.2</div><div>3.2</div></div></div></div><div><div>All Marlborough Sounds excluding Cook Strait</div><div><div><div><div></div><div></div></div><div><div><div>Fixed</div><div>Random</div></div></div><div><div><div>2004</div><div>2007</div><div>2010</div><div>2013</div><div>2017</div></div><div><div>2.0</div><div>1.8</div><div>2.9</div><div>2.1</div><div>3.2</div></div><div><div>1.2</div><div>1.6</div></div></div></div></div></div><div><div><div><div><div>D'Urville Island</div><div><div><div><div></div><div></div></div><div><div><div>Fixed</div><div>Random</div></div></div><div><div><div>1995</div><div>1996</div><div>2001</div><div>2004</div><div>2007</div><div>2010</div><div>2013</div><div>2017</div></div><div><div>4.4</div><div>4.2</div><div>3.9</div><div>4.0</div><div>6.5</div><div>4.0</div><div>2.3</div><div>3.7</div></div></div></div><div><div></div><div>Survey year</div></div></div></div></div><div><p>Marlborough Sounds fixed-site and random-site potting survey catch rates of all blue cod by survey year for each region and overall for the Marlborough Sounds. Error bars are 95% confidence intervals. There were no complete fixed-site surveys in QCH in 1996, PEL in 1996, and DUR from 1995 to 2001. For the overall Marlborough Sounds plot, the 2004 and 2007 fixed-site surveys exclude Separation Point, and the random-site surveys exclude Cook Strait, hence the strata are consistent among the surveys for fixed and random site surveys.</p></div></div></div></div></div></div></div></div></div></div></div></div></div></div></div></div>	
Fishery and Stock Trends	

BLUE COD (BCO)

Recent Trend in Biomass or Proxy	The Marlborough Sounds fixed site potting survey indices of abundance increased markedly in 2010 in the Queen Charlotte Sound and Pelorus regions following the closure of the fishery in the inner sounds in 2008 (QCH, PEL). The survey indices were stable in the D’Urville region where the fishery remained open (DUR). The QCH and PEL fisheries were reopened to a limited size range of blue cod (slot limit) in April 2011 and the estimated 2013 survey abundance in those regions declined, but no change was observed in DUR. In 2017, abundance in QCH was not different to 2013, whereas for PEL and DUR abundance was the highest of any of the surveys. The overall Marlborough Sounds catch rates from 2004 onward (where survey strata are consistent among surveys) indicates that blue cod were more abundant in 2017 than any of the previous surveys. Cook Strait random-site surveys show no trend in abundance from 2010 to 2017. There are only two random site surveys for the other regions (2013 and 2017), not enough to comment on trends.	
Recent Trend in Fishing Mortality or Proxy	Regulatory changes to the recreational fishery (e.g. fishery closures, changes to MLS and daily bag limits) are likely to have resulted in a reduction in fishing mortality up to April 2011, after which mortality increased with the re-opening of the fishery. Fishing mortality was at least twice natural mortality for the random and fixed site surveys in 2017, and the spawning biomass per recruit ratio was 39% (i.e. lower than the target of 45%).	
Other Abundance Indices	The mean length of catches taken during the 2010 blue cod potting survey (following the closure) tended to be larger than those observed in previous surveys and this has generally been maintained in 2013 and 2017	
Trends in Other Relevant Indicators or Variables	Sex ratio is strongly skewed in favour of males. For Marlborough Sounds overall, the percent male from random sites surveys in 2013 was 66%, and in 2017 it was 72% (Table 9). Blue cod catch rates were about 10-fold higher, and length about 5 cm larger overall in the Long Island Marine Reserve (head of QCH) compared to adjacent fished strata in Queen Charlotte Sound. This is a strong indication that fishing pressure has reduced size and abundance of blue cod in the Marlborough Sounds.	
Projections and Prognosis		
Stock Projections or Prognosis	Biomass is expected to increase under current management controls.	
Probability of Current Catch or TACC causing Biomass to remain below or to decline below Limits	Soft Limit: Unknown Hard Limit: Unknown	
Probability of Current Catch or TACC causing overfishing to continue or to commence	Current catches are Likely (> 60%) to cause overfishing to continue	
Assessment Methodology and Evaluation		
Assessment Type	2 – Partial Quantitative Stock Assessment	
Assessment Method	Fishery-independent potting survey. Fixed and random sites in QCH, PEL, DUR, and random sites in CKST.	
Assessment Dates	Latest assessment: 2018	Next assessment: 2022
Overall assessment quality rank	2 – Medium Quality: mortality estimates compromised by regulation changes	
Main data inputs (rank)	- Potting survey catch rates from fixed and random site surveys.	1 – High Quality

BLUE COD (BCO)

	- Length and age composition of catches from random and fixed site potting surveys in 2017	1 – High Quality
Data not used (rank)	N/A	
Changes to Model Structure and Assumptions	-	
Major Sources of Uncertainty	<ul style="list-style-type: none">- Uncertainty in the estimate of M- Frequent regulatory changes for this fishery are likely to have resulted in inconsistent fishing mortality over the lifetime of recent cohorts.- The predominance of males suggests fishing mortality may be higher than estimated.- Trends for random and fixed site surveys between 2013 and 2017 were contradictory in some areas. Random site surveys are believed to be better indicators of population abundance.	
Qualifying Comments		
The survey has been transitioning from a fixed-site to a random-site stratified potting survey. The 2010 survey comprised a full fixed-site survey along with a partial random-site survey in selected strata, whereas 2013 and 2017 included full fixed and full random site surveys carried out simultaneously. The next survey will be based on random sites only.		

Fishery Interactions

Most of the BCO catch is taken by recreational fishers using line methods. There is a reasonably high catch of associated species in this fishery, such as spotted and other wrasses as well as other targeted species such as tarakihi. Most of the commercial catch is taken by potting and has little bycatch.

Table 22: Summary of yields (t), TACCs (t), and reported landings (t) for blue cod from the most recent fishing year.

Fishstocks	QMA	FMA	2016–17 Actual TACC	2016–17 Reported landings
BCO 1	Auckland	1 & 9	46	12
BCO 2	Central (East)	2	10	10
BCO 3	South-East (Coast)	3	163	170
BCO 4	South-East (Chatham Rise)	4	759	741
BCO 5	Southland and Sub-Antarctic	5 & 6	1 239	1 152
BCO 7	Challenger	7	70	60
BCO 8	Central (Egmont)	8	34	5
BCO 10	Kermadecs	10	10	0
Total			2 332	2 150

6. FOR FURTHER INFORMATION

- Beentjes, M P (2012) Correction of catch at age, Z estimates, and SPR estimates for blue cod potting surveys. 46 p. Final Research Report for Ministry of Fisheries project SEA201109. (Unpublished report held by Fisheries New Zealand, Wellington.)
- Beentjes, M P; Carbines, G D (2003) Abundance of blue cod off Banks Peninsula in 2002. *New Zealand Fisheries Assessment Report 2003/16*. 25 p.
- Beentjes, M P; Carbines, G D (2005) Population structure and relative abundance of blue cod (*Parapercis colias*) off Banks Peninsula and in Dusky Sound, New Zealand. *New Zealand Journal of Marine and Freshwater Research* 39: 77–90.
- Beentjes, M P; Carbines, G D (2006) Abundance of blue cod in Banks Peninsula in 2005. *New Zealand Fisheries Assessment Report 2006/01*. 24 p.
- Beentjes, M P; Carbines, G D (2009) Abundance of blue cod in Banks Peninsula in 2008. *New Zealand Fisheries Assessment Report 2009/28*.
- Beentjes, M P; Carbines, G D (2011) Relative abundance, size and age structure, and stock status of blue cod off south Otago in 2010. *New Zealand Fisheries Assessment Report 2011/42*. 60 p.
- Beentjes, M P; Carbines, G D (2012) Relative abundance, size and age structure, and stock status of blue cod from the 2010 survey in Marlborough Sounds, and review of historical surveys. *New Zealand Fisheries Assessment Report 2012/43*. 137 p.
- Beentjes, M.P.; Fenwick, M. (2017). Relative abundance, size and age structure, and stock status of blue cod off Banks Peninsula in 2016. *New Zealand Fisheries Assessment Report 2017/30*. 81 p.
- Beentjes, M P; Francis, R I C C (2011) Blue cod potting surveys: standards and specifications. Version 1. *New Zealand Fisheries Assessment Report 2011/29*.
- Beentjes, M. P; Michael, K; Pallentin, A; Parker, S; Hart, A (2014) Relative abundance, size and age structure, and stock status of blue cod from the 2013 survey in Marlborough Sounds. *New Zealand Fisheries Assessment Report 2014/14*. 85 p.
- Beentjes, M P; Page, M (2016) Relative abundance, size and age structure, and stock status of blue cod in Dusky Sound, Fiordland in 2014. *New Zealand Fisheries Assessment Report 2016/42*. 51 p.


- Beentjes, M P; Page, M (2017) Relative abundance, size and age structure, and stock status of blue cod off Kaikoura in 2015. *New Zealand Fisheries Assessment Report 2017/16*.
- Beentjes, M P; Michael, K; Parker, S; Pallentin, A; Hart, A (2017) Blue cod relative abundance, size and age structure, and habitat surveys of Marlborough Sounds in 2013. *New Zealand Fisheries Assessment Report 2017/61*. 110 p.
- Beentjes, M P; Page, M; Sutton, C; Olsen, L; Walsh, C; Buckthought, D; Sprong, K (in prep). Relative abundance, size and age structure, and stock status of blue cod in the Marlborough Sounds in 2017. Draft New Zealand Fisheries Assessment Report 201X/XX. XX p.
- Beentjes, M.P.; Sutton, C. (2017). Relative abundance, size and age structure, and stock status of blue cod off Motunau in 2016. *New Zealand Fisheries Assessment Report 2017/17*.
- Beer, N A; Wing, S R; Carbines, G (2013) First estimates of batch fecundity for *Paraperis colias*, a commercially important temperate reef fish. *New Zealand Journal of Marine and Freshwater Research* 47: 587–594.
- Bell, J D; Bell, S M; Teirney, L D (1993) Results of the 1991–92 Marine Recreational Fishing Catch and Effort Survey, MAF Fisheries South Region. *New Zealand Fisheries Data Report*: 39.
- Bentley, N; Kendrick, T H (in prep.) Fishery characterisation and Catch-Per-Unit-Effort indices for blue cod in BCO 4; 1989–90 to 2013–14. Draft New Zealand Fisheries Assessment Report.
- Blackwell, R G (1997) Abundance, size composition, and sex ratio of blue cod in the Marlborough Sounds, September 1995. *New Zealand Fisheries Data Report* 88. 17 p.
- Blackwell, R G (1998) Abundance, size and age composition, and yield-per-recruit of blue cod in the Marlborough Sounds, September 1996. *NIWA Technical Report* 30. 16 p.
- Blackwell, R G (2002) Abundance and size composition of recruited blue cod in the Marlborough Sounds, September 2001. Final Research report for Ministry of Fisheries Research Project BCO2001-01. (Unpublished report held by Fisheries New Zealand).
- Blackwell, R G (2005) Abundance and size composition of recruited blue cod in the Marlborough Sounds, September 2005. Final Research report for Ministry of Fisheries Research Project BCO2003-01. (Unpublished report held by Fisheries New Zealand).
- Blackwell, R G (2009) Abundance and size composition of blue cod in the Marlborough Sounds, and Tasman Bay September–October 2007. Final Research report for Ministry of Fisheries Research Project BCO2006-01. (Unpublished report held by Fisheries New Zealand).
- Boyd, R O; Gowing, L; Reilly, J L (2004) 2000–2001 national marine recreational fishing survey: diary results and harvest estimates. Final Research Report for Ministry of Fisheries project REC2000/03. 81 p. (Unpublished report held by Fisheries New Zealand, Wellington.)
- Boyd, R O; Reilly, J L (2002) 1999–00 national marine recreational fishing survey: harvest estimates. Draft New Zealand Fisheries Assessment Report held by Fisheries New Zealand Wellington.
- Bradford E. (1998) Harvest estimates from the 1996 national recreational fishing surveys. New Zealand Fisheries Assessment Research Document 1998/16. 27 p. (Unpublished document held in NIWA library, Wellington.)
- Carbines, G D (1998) Blue cod age validation, tagging feasibility and sex-inversion. Final report to the Ministry of Fisheries for Project SOBC04. 77 p. (Unpublished document held by Fisheries New Zealand Wellington.)
- Carbines, G D (1999) Large hooks reduce catch-and-release mortality of blue cod *Paraperis colias* in the Marlborough Sounds of New Zealand. *North American Journal of Fisheries Management* 19(4): 992–998.
- Carbines, G D (2000) Comparisons of age and growth of blue cod within the Marlborough Sounds (BCO7). Final report to the Ministry of Fisheries for Project BCO9801. (Unpublished document held by Fisheries New Zealand Wellington.)
- Carbines, G D (2001) Movement patterns and stock mixing of blue cod in Southland. Final report to the Ministry of Fisheries for Project BCO9702. (Unpublished document held by Fisheries New Zealand Wellington.)
- Carbines, G (2004a) Age, growth, movement and reproductive biology of blue cod (*Paraperis colias*-Pinguipedidae): Implications for fisheries management in the South Island of New Zealand. Unpublished Ph.D. thesis, University of Otago, Dunedin, New Zealand. 211 p.
- Carbines, G (2004b) Age determination, validation, and growth of blue cod *Paraperis colias*, in Foveaux Strait, New Zealand. *New Zealand Journal of Marine and Freshwater Research* 38: 201–214.
- Carbines, G D (2007) Relative abundance, size, and age structure of blue cod in Paterson Inlet (BCO 5), November 2006. *New Zealand Fisheries Assessment Report 2007/37*. 31 p.
- Carbines G D; Beentjes, M P (2003) Relative abundance of blue cod in Dusky Sound in 2002. *New Zealand Fisheries Assessment Report 2003/37*. 25 p.
- Carbines, G; Beentjes, M P (2006a) Relative abundance of blue cod in North Canterbury in 2004 and 2005. *New Zealand Fisheries Assessment Report 2006/30*. 26 p.
- Carbines, G; Beentjes, M P (2006b) Relative abundance of blue cod in north Otago in 2005. *New Zealand Fisheries Assessment Report 2006/29*. 20 p.
- Carbines, G D; Beentjes, M P (2009) Relative abundance, size and age structure, and mortality of blue cod in north Canterbury (BCO 3) in 2007/08. *New Zealand Fisheries Assessment Report 2009/37*.
- Carbines, G D; Beentjes, M P (2011a) Relative abundance, size and age structure, and stock status of blue cod in Dusky Sound, Fiordland, in 2008. *New Zealand Fisheries Assessment Report 2011/35*. 56 p.
- Carbines, G D; Beentjes, M P (2011b) Relative abundance, size and age structure, and stock status of blue cod off north Otago in 2009. *New Zealand Fisheries Assessment Report 2011/36*. 57 p.
- Carbines, G D; Beentjes, M P (2012) Relative abundance, size and age structure, and stock status of blue cod in Foveaux Strait in 2010. *New Zealand Fisheries Assessment Report 2012/39*. 66 p.
- Carbines, G; Cole, R G (2009) Using a remote drift underwater video (DUV) to examine dredge impacts on demersal fishes and benthic habitat complexity in Foveaux Strait, Southern New Zealand. *Fisheries Research* 96: 230–237.
- Carbines, G; Dunn, A; Walsh, C (2007) Age composition and estimates of mortality of blue cod from seven relative abundance South Island potting surveys. Inshore Stock Assessment Working Group Meeting paper, INS WG 2007/24. (Unpublished working group paper held by Fisheries New Zealand.)
- Carbines, G; Dunn, A; Walsh, C (2008) Age composition and derived estimates of total mortality for blue cod taken in South Island potting surveys, 2002–2005. *New Zealand Fisheries Assessment Report 2008/68*. .
- Carbines, G D; Haist, V (2012a) Relative abundance, size and age structure, and stock status of blue cod off Banks Peninsula in 2012. SINS-WG-2012-23. (Unpublished working group paper held by Fisheries New Zealand.)
- Carbines, G D; Haist, V (2012b) Relative abundance, size and age structure, and stock status of blue cod off North Canterbury (Kaikoura & Motunau) in 2011/12. SINS-WG-2012-24. (Unpublished working group paper held by Fisheries New Zealand.)
- Carbines, G D; Haist, V (2014a) Relative abundance, size and age structure, and stock status of blue cod in Paterson Inlet of BCO 5 in 2010. *New Zealand Fisheries Assessment Research Report 2014/14*. 84 p.
- Carbines, G D; Haist, V (2014b) Relative abundance, size structure, and stock status of blue cod off North Otago in 2013. SINS-WG-2014-30. (Unpublished working group paper held by Fisheries New Zealand.)

BLUE COD (BCO)

- Carbines, G D; Haist, V (2014c) Relative abundance, size structure, and stock status of blue cod off South Otago in 2013. SINS-WG-2014-31. (Unpublished working group paper held by Fisheries New Zealand.)
- Carbines, G D; Haist, V (2016a) Relative abundance of blue cod in the Foveaux Strait. SINS-WG-2016-23. (Unpublished working group paper held by Fisheries New Zealand.)
- Carbines, G D; Haist, V (2016b) Relative abundance of blue cod in Paterson Inlet. SINS-WG-2016-24. (Unpublished working group paper held by Fisheries New Zealand.)
- Carbines, G; Haist, V (2017) Relative abundance, size and age structure, and stock status of blue cod off Banks Peninsula in 2012. *New Zealand Fisheries Assessment Report 2017/37*. 126 p.
- Carbines, G.D.; Haist, V. (2018). Relative abundance, size and age structure, and stock status of blue cod off Kaikoura and north Canterbury in 2011–12. Comparisons of potting survey designs and estimates of pot catchability and size selectivity. *New Zealand Fisheries Assessment Report 2018/06*. 97 p.
- Carbines, G; Jiang, W; Beentjes, M P (2004) The impact of oyster dredging on the growth of blue cod, *Parapercis colias*, in Foveaux Strait, New Zealand. *Aquatic Conservation* 14: 491–504.
- Carbines, G; McKenzie, J (2004) Movement patterns and stock mixing of blue cod in Dusky South in 2002. *New Zealand Fisheries Assessment Report 2004/36*. 28 p.
- Cole, R (1999) A comparison of abundance, population size structure, and sex ratio of blue cod *Parapercis colias* sampled by pot and diver count methods in the Marlborough Sounds. Final report to the Ministry of Fisheries for Project BCO9701. (Unpublished report held by Fisheries New Zealand, Wellington.)
- Cranfield, H J; Carbines, G; Michael, K P; Dunn, A; Stotter, D R; Smith, D L (2001) Promising signs of regeneration of blue cod and oyster habitat changed by dredging in Foveaux Strait, southern New Zealand. *New Zealand Journal of Marine and Freshwater Research* 35: 897–908.
- Davey, N K; Hartill, B (2011) Survey of the Southland recreational blue cod fishery during the 2009–2010 fishing year. *New Zealand Fisheries Assessment Report 2011/57*. 44 p.
- Davey, N K; Hartill, B; Caimey, D G; Cole, R G (2008) Characterisation of the Marlborough Sounds recreational fishery and associated blue cod and snapper harvest estimates. *New Zealand Fisheries Assessment Report 2008/31*. 63 p.
- Francis, M P; Paul, L J (2013) New Zealand inshore finfish and shellfish commercial landings, 1931–82. *New Zealand Fisheries Assessment Report 2013/55*. 136 p.
- Haist, V; Breen, P A; Starr, P J (2009) A new multi-stock length-based assessment model for New Zealand rock lobsters (*Jasus edwardsii*). *New Zealand Journal of Marine and Freshwater Research* 43(1): 355–371.
- Hartill, B; Bian, R; Armiger, H; Vaughan, M; Rush, N (2007) Recreational marine harvest estimates of snapper, kahawai, and kingfish in QMA 1 in 2004–05. *New Zealand Fisheries Assessment Report 2007/26*. 44 p.
- Hartill B.; Davey, N.; Bradley, A.; Carter, M.; Olsen, L.; Bian, R. (2017). Aerial-access recreational harvest estimates for snapper and blue cod in FMA 7 in 2015–16. *New Zealand Fisheries Assessment Report 2017/34*. 28 p.
- Hoenig, J M (1983) Empirical use of longevity data to estimate mortality rates. *Fishery Bulletin* 81: 898–903.
- James, G D; Unwin, M J (2000) National marine diary survey of recreational fishing from charter vessels, 1997–98. *NIWA Technical Report* 70. 51 p.
- Jiang, W; Carbines, G D (2002) Diet of blue cod, *Parapercis colias*, living on undisturbed biogenic reefs and on seabed modified by oyster dredging in Foveaux Strait, New Zealand. *Aquatic Conservation* 12: 257–272.
- Kendrick, T H; Bentley, N (2011) Fishery characterisation and Catch-Per-Unit-Effort indices for blue cod in BCO 4; 1989–90 to 2008–09. Progress Report for Ministry of Fisheries project BCO2009-04. (Unpublished report held by Fisheries New Zealand, Wellington.)
- Langley, A D (2005) Summary of catch and effort data from the BCO 3 and BCO 5 fisheries, 1989–90 to 1999–2000. *New Zealand Fisheries Assessment Report*. 2005/30. 28 p.
- Leach, B F; Boocock, A S (1993) Prehistoric fish catches in New Zealand. *Tempus Reparatum. BAR International Series* 584: 38 p.
- Mace, J T; Johnston, A D (1983) Tagging experiments on blue cod (*Parapercis colias*) in the Marlborough Sounds, New Zealand. *New Zealand Journal of Marine and Freshwater Research* 17: 207–211.
- McGregor, G A (1988) Blue cod. New Zealand Fisheries Assessment Research Document 1988/41. 11 p. (Unpublished document held in NIWA library, Wellington.)
- Mutch, P G (1983) Factors influencing the density and distribution of the blue cod (*Parapercis colias*). (Unpublished M.Sc. thesis held in University of Auckland library, Auckland.)
- Rapson, A M (1956) Biology of the blue cod (*Parapercis colias* Foster) of New Zealand (Unpublished Ph.D. thesis held in Victoria University library, Wellington.)
- SeaFIC (2005) Report to the Inshore Fishery Assessment Working Group: BCO 5 characterisation and CPUE analysis. 35 p. (Unpublished report held by Seafood New Zealand, Wellington.)
- Smith, H M (2012) Characterisation of the Mitochondrial Genome and the Phylogeographic Structure of Blue Cod (*Parapercis colias*). (Unpublished M.Sc. thesis held by Victoria University of Wellington.)
- Starr, P J; Kendrick, T H (2009) Report to Southeast Finfish Management LTD: Review of the BCO 5 fishery. 51 p. (Unpublished report held by Seafood New Zealand, Wellington.)
- Starr, P J; Kendrick, T H (2011) Report to Southeast Finfish Management Ltd: Review of the BCO 5 Fishery. 67 p. (Unpublished report held by Seafood New Zealand, Wellington.)
- Teirney, L; Bell, S; Bell, J (1992) MAF Fisheries South Region Survey of Marine Recreational Fishers — Summary of Findings. New Zealand Fisheries Management: Regional Series: 1, 23p. (Available at NIWA library).
- Teirney, L D; Kilner, A R; Millar, R E; Bradford, E; Bell, J D (1997) Estimation of recreational catch from 1991/92 to 1993/94. New Zealand Fisheries Assessment Research Document 1997/15. 43 p. (Unpublished document held by NIWA library, Wellington.)
- Walsh, C. (2017). Age determination protocol for blue cod (*Parapercis colias*). *New Zealand Fisheries Assessment Report 2017/15*. 34 p.
- Warren, E J (1994) The blue cod fishery in the Marlborough Sounds. 30 p. MAF Fisheries Central Internal Report available at NIWA library.
- Warren, E J; Grindley, R M; Carbines, G D; Teirney, L (1997) Characterisation of the Southland blue cod fishery (1991–1996). (MFish South report available at Invercargill City Library.)
- Wright, P; Gowing, L; McClary, D; Boyd, R O (2004) 2000–2001 National Marine recreational fishing survey: direct questioning of fishers compared with reported diary data. Final Research report of Ministry of Fisheries Research Project REC2000-01: Objective 2. 28 p. (Unpublished report held by Fisheries New Zealand, Wellington.)

BLUE MACKEREL (EMA)*(Scomber australasicus)*

Tawatawa

**1. FISHERY SUMMARY**

Blue mackerel were introduced into the QMS on 1 October 2002. Since then allowances, TACCs and TACs (Table 1) have not changed.

Table 1: Recreational and Customary non-commercial allowances, TACCs and TACs for blue mackerel by Fishstock.

Fishstock	Recreational Allowance	Customary Non-Commercial Allowance	TACC	TAC
EMA 1	40	20	7 630	7 690
EMA 2	5	2	180	187
EMA 3	1	1	390	392
EMA 7	1	1	3 350	3 352
EMA 10	0	0	0	0
Total	47	24	11 550	11 621

1.1 Commercial fisheries

Blue mackerel are taken by a variety of methods but for most of these methods the catches are very low. The largest and most consistent catches have been from the target purse seine fishery in EMA 1, 2 and 7, and as non-target catch in the jack mackerel mid-water trawl fishery in EMA 7. Most catch is taken north of latitude 43° S (Kaikoura). Historical estimated and recent reported blue mackerel landings and TACCs are shown in Tables 2 and 3, while Figure 1 shows the historical landings and TACC values for these three main stocks. Since 1983–84 the catch of blue mackerel in New Zealand waters has grown substantially (Table 3), primarily in the purse seine fishery in EMA 1, and have averaged about 10 000 t annually since 1990–91.

Most blue mackerel purse seine catch comes from the Bay of Plenty (BoP) and East Northland, where it is primarily taken between July and December. Purse seine fishing effort on blue mackerel has been strongly influenced by the availability and market value of other pelagic species, particularly skipjack tuna and kahawai, with effort increasing as limits have been placed on the purse seine catch of kahawai. The purse seine fishery has accounted for more than 97% of annual EMA 1 landings since at least 1990, and about 90% of this was targeted (Ballara 2016).

Total blue mackerel catches peaked in 1991–92 at more than 15 000 t, of which 60–70% was taken by purse seine. More recently, commercial landings of over 12 500 t were taken in 1998–99 (13 500 t), 2000–01 (13 100 t) and 2004–05 (12 750 t), with the highest landings recorded in EMA 1 and EMA 7. EMA 1 landings exceeded the TACC in 2004–05, 2006–07, 2009–10, 2011–12 and 2014–15. The 2004–05, 2005–06, and 2008–09 EMA 7 landings also exceeded the TACC. The EMA 7 landings in 2016–17, however, were the lowest since the mid-1980s. Landings from EMA 2 and EMA 3 have been well

BLUE MACKEREL (EMA)

below the TACCs since the early to mid-1990s; they are primarily a bycatch of purse seine (EMA 2) and trawl (EMA 3) fisheries.

The blue mackerel catch from EMA 7 is now principally non-target catch from the jack mackerel mid-water trawl fishery. Highest catches are taken during June, July and October in areas 034 and 035 on the WCSI and areas 041 and 801 further north (WCNI). Fishing has shifted from south to north in the last decade. Since the late 1990s, a fleet of Ukrainian vessels has taken most of the catch in the JMA 7 target fishery and these vessels have taken the EMA as bycatch. Since 2004, 0–11% of the EMA 7 catch has been taken annually by purse seine, down from an average of about 25% between 1991 and 2003.

A number of factors have been identified that can influence landing volumes in the blue mackerel fisheries. In the purse seine fishery, blue mackerel has become the second most preferred species because of decreased TACCs on kahawai. Skipjack tuna is the preferred species and blue mackerel will not be targeted once the skipjack season has begun in late-spring, early summer. Thus, early arrival of skipjack can result in reduced volumes of blue mackerel being landed.

Management of company quota is complicated by the relative timing of the fishing season and the fishing year and this, along with the timing of the main market, may influence whether the blue mackerel TACC can all be taken in a particular year. The fishing season usually begins in about July–August, runs through to the end-beginning of subsequent fishing years, and finishes in about November. The main market for purse seined blue mackerel takes up to 80% of the catch and requires premium fish to be available from early spring. To meet the demands of this market and to minimise the costs of storing fish from the previous season, fishing companies must carry over some proportion of their quota for a given year until fish become available the following season. If availability is delayed until after October 1, only 10% of the total quota can then be carried over into the new fishing year.

Because blue mackerel is taken principally as bycatch in the jack mackerel TCEPR target fishery in JMA 7, factors influencing the targeting of jack mackerel also affect blue mackerel landings. Other bycatch species taken in this fishery include barracouta, gurnard, John dory, kingfish, and snapper, and, although non-availability of ACE is unlikely to be constraining in the first three of these, the same is not true of kingfish and snapper. Fishing company spokespersons have stated that known hotspots of snapper are avoided. Other factors in this fishery include strategies to avoid the catch of marine mammals, and a code of practice operates in which gear is not deployed between 2 a.m. and 4 a.m. It is unknown whether this affects total landing volumes.

Table 2: Reported landings (t) for the main QMAs from 1931 to 1982.

Year	EMA 1	EMA 2	EMA 3	EMA 7	Year	EMA 1	EMA 2	EMA 3	EMA 7
1931–32	0	0	0	0	1957	0	0	0	0
1932–33	0	0	0	0	1958	0	0	0	0
1933–34	0	0	0	0	1959	0	0	0	0
1934–35	0	0	0	0	1960	0	0	0	0
1935–36	0	0	0	0	1961	0	0	0	0
1936–37	0	0	0	0	1962	0	0	0	0
1937–38	0	0	0	0	1963	0	0	0	0
1938–39	0	0	0	0	1964	0	0	0	0
1939–40	0	0	0	0	1965	0	0	0	0
1940–41	0	0	0	0	1966	0	0	0	0
1941–42	0	0	0	0	1967	0	0	0	0
1942–43	0	0	0	0	1968	0	0	0	0
1943–44	0	0	0	0	1969	0	0	0	0
1944	0	0	0	0	1970	0	0	0	0
1945	0	0	0	0	1971	0	0	0	0
1946	0	0	0	0	1972	0	0	0	0
1947	0	0	0	0	1973	0	0	0	0
1948	0	0	0	0	1974	38	8	0	6
1949	0	0	0	0	1975	10	0	0	2
1950	0	0	0	0	1976	50	49	0	0
1951	0	0	0	0	1977	34	135	0	0
1952	0	0	0	0	1978	14	55	0	128
1953	0	0	0	0	1979	185	31	0	317
1954	0	0	0	0	1980	752	32	0	407
1955	0	0	0	0	1981	459	49	0	1363
1956	0	0	0	0	1982	305	0	0	791

Notes

1. The 1931–1943 years are April–March but from 1944 onwards are calendar years.


2. Data up to 1985 are from fishing returns: Data from 1986 to 1990 are from Quota Management Reports.

3. Data for the period 1931 to 1982 are based on reported landings by harbour and are likely to be underestimated as a result of under-reporting and discarding practices. Data includes both foreign and domestic landings.

Table 3: Reported landings (t) of blue mackerel by QMA, and where area was unspecified (Unsp.), from 1983–84 to 2016–17. CELR data from 1986–87 to 2000–01. MHR data from 2001–02 to present.

Fishing year	QMA					Unsp	Total
	1	2	3	7	10#		
1983–84*	480	259	44	245	0	1	1 028
1984–85*	565	222	18	865	0	73	1 743
1985–86*	618	30	190	408	0	51	1 296
1986–87	1 431	7	424	489	0	49	2 399
1987–88	2 641	168	864	1 896	0	58	5 625
1988–89	1 580	< 1	1 141	1 021	0	469	4 211
1989–90	2 158	76	518	1 492	0	< 1	4 245
1990–91	5 783	94	478	3 004	0	0	9 358
1991–92	10 926	530	65	3 607	0	0	15 128
1992–93	10 684	309	133	1 880	0	0	13 006
1993–94	4 178	218	223	1 402	5	0	6 025
1994–95	6 734	94	154	1 804	10	149	8 944
1995–96	4 170	119	173	1 218	0	1	5 680
1996–97	6 754	78	340	2 537	0	< 1	9 708
1997–98	4 595	122	78	2 310	0	< 1	7 104
1998–99	4 505	186	62	8 756	0	4	13 519
1999–00	3 602	73	3	3 169	0	0	6 847
2000–01	9 738	113	6	3 278	0	< 1	13 134
2001–02	6 368	177	49	5 101	0	0	11 694
2002–03	7 609	115	88	3 563	0	0	11 375
2003–04	6 523	149	1	2 701	0	0	9 373
2004–05	7 920	9	< 1	4 817	0	0	12 746
2005–06	6 713	13	133	3 784	0	0	10 643
2006–07	7 815	133	42	2 698	0	0	10 688
2007–08	5 926	6	122	2 929	0	0	8 982
2008–09	3 147	2	88	3 503	0	0	6 740
2009–10	8 539	3	14	3 260	0	0	11 816
2010–11	6 630	2	9	1 996	0	0	8 638
2011–12	8 080	2	28	2 707	0	0	10 817
2012–13	7 213	3	100	2 401	0	0	9 716
2013–14	6 860	4	29	1 200	0	0	8 092
2014–15	8 134	16	87	892	0	0	9 129
2015–16	7 226	18	27	761	0	0	8 033
2016–17	7 551	83	126	625	0	0	8 385

* FSU data, # Landings reported from QMA 10 are probably attributable to Statistical Area 010 in the Bay of Plenty (i.e., QMA 1).

**Figure 1: Reported commercial landings and TACC for the three main EMA stocks. EMA 1 (Auckland East) (continued on next page).**

BLUE MACKEREL (EMA)




Figure 1: Reported commercial landings and TACC for the three main EMA stocks. From top: EMA 2 (Central East), and EMA 7 (Challenger to Auckland West).

1.2 Recreational fisheries

Blue mackerel does not rate highly as a recreational target species although it is popular as bait. There is some uncertainty with all recreational harvest estimates for blue mackerel and there is some confusion between blue and jack mackerels in the recreational data.

Recreational catch in the northern region (EMA 1) was estimated at 114 000 fish by a diary survey in 1993–94 (Bradford 1996), 47 000 fish in a national recreational survey in 1996 (Bradford 1998), 84 000 fish (CV 42%) in the 2000 survey (Boyd & Reilly 2005) and 58 000 fish (CV 27%) in the 2001 survey (Boyd et al 2004). The surveys suggest a harvest of 35–90 t per year for EMA 1, insignificant in the context of the commercial catch. Estimates from other areas are very low (between 500 and 3000 fish) and are likely to be insignificant in the context of the commercial catch.

The harvest estimates provided by telephone-diary surveys between 1993 and 2001 are no longer considered reliable for various reasons. A Recreational Technical Working Group concluded that these harvest estimates should be used only with the following qualifications: a) they may be very inaccurate; b) the 1996 and earlier surveys contain a methodological error; and c) the 2000 and 2001 estimates are implausibly high for many important fisheries. In response to these problems and the cost and scale challenges associated with onsite methods, a National Panel Survey was conducted for the first time throughout the 2011–12 fishing year. The panel survey used face-to-face interviews of a random sample of 30 390 New Zealand households to recruit a panel of fishers and non-fishers for a full year. The panel members were contacted regularly about their fishing activities and harvest information collected in

standardised phone interviews. Harvest estimates for blue mackerel are given in Table 4 (from Wynne-Jones et al 2014 and Hartill & Davey 2015).

Table 4: Recreational harvest estimates for blue mackerel stocks (Wynne-Jones et al 2014). Mean fish weights were obtained from boat ramp surveys; for blue mackerel the value used was 1.039 kg (Hartill & Davey 2015).

Stock	Year	Method	Number of fish	Total weight (t)	CV
EMA 1	2011/12	Panel survey	18 438	19.2	0.36
EMA 2	2011/12	Panel survey	3 346	3.5	-
EMA 7	2011/12	Panel survey	11 194	11.6	0.42
EMA total	2011/12	Panel survey	32 976	34.3	0.25

A repeat of the National Panel Survey is being conducted over the 2017–18 October fishing year. Results are expected in early 2019.

1.3 Customary non-commercial fisheries

Quantitative information on the current level of customary non-commercial catch is not available.

1.4 Illegal catch

There is no known illegal catch of blue mackerel.

1.5 Other sources of mortality

There is no information on other sources of mortality.

2. BIOLOGY

The geographical distribution and habitat of blue mackerel vary with life history stage. Juvenile and immature blue mackerel are northerly in their distribution, having been recorded from commercial and research catches around the North Island and into Golden and Tasman Bay at the top of the South Island.

By contrast, adults have been recorded around both the North and South Islands to Stewart Island and across the Chatham Rise almost to the Chatham Islands. Sporadic catches of small numbers of yearling blue mackerel have been made by bottom trawl in shallow waters.

The distribution of blue mackerel at the surface is seasonal and differs from its known geographical range. During summer, surface schools are found in Northland, BoP, South Taranaki Bight, and Kaikoura, but they disappear during winter, when only occasional individuals are found in Northland and the BoP. A possible corollary to this winter disappearance comes from the peak in bycatch of blue mackerel in the winter jack mackerel mid-water trawl fishery in EMA 7. This suggests an increased partitioning of the population in deeper water at this time of the year, reflecting an observed behavioural characteristic of the related Atlantic species, *Scomber scombrus*. Summaries from aerial sightings data show that blue mackerel can be found in mixed schools with jack mackerel (*Trachurus* spp.), kahawai (*Arripis trutta*), skipjack tuna (*Katsuwonus pelamis*) and trevally (*Pseudocaranx dentex*), and that its appearance in mixed schools varies seasonally.

Blue mackerel are serial spawners, releasing eggs in batches over several months. Based on gonad condition, sexual maturity for both sexes of blue mackerel taken in the Great Australian Bight between January 1979 and December 1980 was estimated to be about 28 cm FL, which translates to an age of about 2 years. Eggs are pelagic and development rate is dependent on temperature. In plankton surveys, blue mackerel eggs have been found from North Cape to East Cape, with highest concentrations from Northland, the Hauraki Gulf, and the Western BoP. Eggs have been described throughout the Hauraki Gulf from November to the end of January, at surface temperatures in the range 15–23°C. Individuals in spent or spawning condition have been taken in a few tows off Tasman Bay and Taranaki, in EMA 7 and in the BoP in EMA 1.

Age and growth studies suggest a difference in the age structures of catches taken in the BoP (New Zealand, EMA 1) and New South Wales (Australia). For fish from the New South Wales study, a peak was found at 1 year that accounted for more than 55% of the fish sampled, with a maximum age of 7

BLUE MACKEREL (EMA)

yr. The BoP results show a much broader distribution, with a maximum age of 24 yr, and a mode in the data around 8 to 10 yr. Growth parameters estimated in the BoP study are given in Table 5. Following a quantitative test of competing growth models in the BoP study, no evidence was found of statistically significant differences in growth between the sexes in BoP blue mackerel.

Australian studies may underestimate the ages of larger, older blue mackerel in their catch. The Australian method for estimating blue mackerel ages is based on reading otoliths whole in oil, whereas the New Zealand method is based on otolith thin-sections (Marriott & Manning 2011). Results from the New South Wales study referred to above, suggest that blue mackerel 25–40 cm fork length may be 3–7 years old. Using the New Zealand method, fish in this length range could be as old as 16 years. Australian scientists, reading whole otoliths, may be missing opaque zones near the margin, which are visible in sectioned otoliths.

Table 5: Von Bertalanffy growth parameters for Bay of Plenty (EMA 1) blue mackerel (Manning et al 2006).

	Males	Females	Both sexes
L_{∞}	52.49	53.10	52.79
K	0.15	0.15	0.15
t_0	-3.29	-3.18	-3.19
Age range	1.8–21.9	1.8–21.9	1.8–21.9
N	240	269	509

Although Australian scientists have validated the timing of the first opaque zone in blue mackerel otoliths, their results do not cover the complete life history defined using either the Australian or New Zealand method. A study attempting to validate the New Zealand age estimation method using lead-radium dating indicated that blue mackerel in New Zealand are a relatively long lived small pelagic species, living to at least 17 to 49 years, with the real age most likely nearer the lower value (Marriott et al 2010). While this range of age estimates is less than desirable for the validation of the growth zone counting method for this species, the findings are consistent with the New Zealand method where otolith ageing studies from commercial catches have blue mackerel living to at least 24 years.

Instantaneous natural mortality (M) for male and female fish was estimated using Hoenig's method (Morrison et al 2001a). Based on age estimates from otoliths collected during the mid 1980s when fishing pressure was presumably light, natural mortality estimates of 0.22 for males and 0.20 for females were derived.

In New Zealand, the diet of blue mackerel has been described as zooplankton, which consists mainly of copepods, but also includes larval crustaceans and molluscs, fish eggs and fish larvae. Feeding involves both filtering of the water and active pursuit of prey, with blue mackerel able to take much smaller animals than, for example, kahawai can.

3. STOCKS AND AREAS

Sampling of eggs, larvae, and spawning blue mackerel indicate at least three spawning centres for this species: Northland-Hauraki Gulf; Western BoP; and South Taranaki Bight. Nothing is known of migratory patterns or the fidelity of fish to a particular spawning area. Examination of mitochondrial DNA shows no geographical structuring between New Zealand and Australian fish. Meristic characters show significant regional differentiation within New Zealand fisheries waters and, combined with parasite marker information, Smith et al (2005) sub-divided blue mackerel into at least three stocks in New Zealand fisheries waters: EMA 1, EMA 2, and EMA 7. No information is currently available on the stock affinity of fish in EMA 3.

4. STOCK ASSESSMENT

4.1 EMA 1

4.1.1 Estimates of fishery parameters and abundance

Analysis of aerial sightings data for east Northland (part of EMA 1) from 1985–86 to 2002–03 found no apparent trends in abundance, apart from a peak off east Northland in 1991–92 for both the number of schools and the estimated tonnage, and a further strong signal for the number of schools and the estimated tonnage from 2000–01 to 2002–03.

Using market and catch sampling data collected from 2002 to 2005, estimated numbers-at-length and numbers-at-age were calculated based on all available groomed length and length-at-age data (Manning et al 2007). These were done separately by sex and scaled to estimates of the total catch from the purse seine fishery. Results showed that the EMA 1 purse seine fishery was composed of fish between 2–21 years of age, although most were between 5 and 15 years.

4.2. EMA 7

4.2.1 Estimates of fishery parameters and abundance

A standardised CPUE analysis for EMA 7 was carried out using TCEPR tow by tow data from the mid-water trawl jack mackerel target fishery up to 2013–14 in which blue mackerel form a significant and important bycatch (Ballara 2016). The initial dataset comprised tows that targeted jack mackerel with blue mackerel caught as bycatch. Tows that targeted blue mackerel were not considered as they constituted a small amount of catch and effort (about 30 tows each year for the last 10 years by all vessels) and they were confined to a few areas in the fishery and were directed at large sub-surface schools of blue mackerel. Tows that targeted jack mackerel but did not report any blue mackerel catch were also excluded. The data used for the CPUE analyses consisted of catch and effort by core vessels that targeted jack mackerel; core vessels were those participating in the fishery for five or more years, and reporting at least 20 tows per vessel-year. Estimates of relative year effects were obtained using a forward stepwise multiple regression method, where the data were fitted using binomial-lognormal model structure.

Separate standardisations were carried out to two subgroups of core vessels corresponding to an early and late period of the data series respectively. CPUE indices were developed for the early time series from 1989–90 to 1997–98 using catch and effort by 12 core vessels and the late time series from 1996–97 to 2013–14 using catch and effort by 7 core vessels (Table 6). The residual deviance explained was 33% for the early time series and 35% for the late time series. For both data series, the main terms selected by the models are statistical area, vessel, and month.

The early time series increased from 1990 to 1992, and was then relatively constant to 1998. The late time series declined steadily from 1997 to about 2005, and has been relatively constant since then (although the three most recent years produced the lowest indices from this series). Similar trends were also apparent for the later series analysed separately by WCSI and WCNI areas (Figure 2). The series from 2000 onwards shows a decline of more than 50%.

The WG concluded that standardised CPUE series based on the blue mackerel bycatch in the WCNI and WCSI jack mackerel trawl fishery appears to provide reliable indices of abundance.

Using market and catch sampling data collected from 2002 to 2005, estimated numbers-at-length and numbers-at-age were calculated based on all available groomed length and length-at-age data (Manning et al 2007). These were done separately by sex and scaled to estimates of the total catch from the purse seine and the trawl fisheries. Results showed that the EMA 7 purse seine fishery was composed of fish between 2–24 years of age, although most were between 5 and 15 years. Catch-at-age in the EMA 7 mid-water trawl TCEPR bycatch (jack mackerel target) fishery also showed a wide range, with fish between 2–24 years represented, and small peaks evident between 10 and 11 years in both sexes. These results were generally consistent with those from previous years, although relatively low numbers of small fish in the sampled fisheries were noted.

BLUE MACKEREL (EMA)

Table 6: Standardised lognormal CPUE catch/hr indices for the core West coast TCEPR tow-by-tow target JMA data indices for fishing years 1990–2014. The Standardised CPUE indices for the early series is from 1990 to 1998 (from Fu & Taylor 2011) and for the late series from 1997 to 2014 (Ballara 2016).

Year	Indices	CV	Indices	CV
1990	0.67	0.20	–	–
1991	0.87	0.10	–	–
1992	1.24	0.11	–	–
1993	1.01	0.13	–	–
1994	0.99	0.09	–	–
1995	1.05	0.07	–	–
1996	0.87	0.11	–	–
1997	1.34	0.08	2.27	0.09
1998	1.13	0.08	1.99	0.07
1999	–	–	2.22	0.05
2000	–	–	2.03	0.05
2001	–	–	1.66	0.05
2002	–	–	1.73	0.04
2003	–	–	1.17	0.05
2004	–	–	0.80	0.04
2005	–	–	0.70	0.04
2006	–	–	0.86	0.04
2007	–	–	0.60	0.04
2008	–	–	0.69	0.04
2009	–	–	0.84	0.04
2010	–	–	0.71	0.04
2011	–	–	0.75	0.04
2012	–	–	0.55	0.05
2013	–	–	0.57	0.05
2014	–	–	0.52	0.06




Figure 2: Blue mackerel CPUE for 1997–2014 for West coast (WC); WCSI, and WCNI. Indices have been standardised to have a mean of one.

4.3 Biomass estimates

No estimates of biomass are available for any blue mackerel stocks.

4.4 Other factors

Catch sampling in the period from 2002 to 2005 indicated that catch-at-length and catch-at-age is relatively stable between years in EMA 1. Although total mortality in EMA 1 is poorly understood, the relatively stable age-length composition between years and the number of year-classes that compose the catch-at-age within fishing years, suggested that blue mackerel may be capable of sustaining the catch levels at that time in EMA 1.

5. STATUS OF THE STOCKS


Based on studies of stock structure within New Zealand waters blue mackerel may be sub-divided into at least three stocks: EMA 1, EMA 2, and EMA 7. No information is currently available on the stock affinity of fish in EMA 3.

Little is known about the status of blue mackerel stocks and no estimates of current and reference biomass, or yield, are available for any blue mackerel area.

• EMA 1

For EMA 1, the stability of the age composition data and the large number of age classes that comprise the catches suggests that blue mackerel may be capable of sustaining current commercial fishing mortality, at least in the short-term.

• EMA 7



Fishery and Stock Trends	
Recent Trend in Biomass or Proxy	CPUE has continued to decline since 2009
Recent Trend in Fishing Intensity or Proxy	
Other Abundance Indices	-

BLUE MACKEREL (EMA)

Trends in Other Relevant Indicators or Variables	Broad age structure of the trawl catch (2004–05) did not support a large decrease in biomass from 1999 to 2005 as suggested by the CPUE series
--	--

Projections and Prognosis	
Stock Projections or Prognosis	Unknown
Probability of Current Catch or TACC causing Biomass to remain below or to decline below Limits	Unknown
Probability of Current Catch or TACC causing Overfishing to continue or to commence	Unknown

Assessment Methodology and Evaluation		
Assessment Type	Level 2 - Partial Quantitative Stock Assessment	
Assessment Method	Standardised CPUE from the jack mackerel target fishery WCSI and WCNI	
Assessment Dates	Latest assessment: 2016	Next assessment: 2019
Overall assessment quality rank	1 – High Quality	
Main data inputs (rank)	- Standardised CPUE - Proportions at age data from the commercial trawl fishery	1 – High Quality 1 – High Quality
Data not used (rank)		
Changes to Model Structure and Assumptions	-	
Major sources of Uncertainty	-	

Qualifying Comments
The decline in CPUE from 1999 to 2005 was not consistent with the broad range of ages in the trawl catch. However, no recent age data are available.

Fishery Interactions
There is a small target fishery for blue mackerel on the WCNI but the bulk of the catch is taken as bycatch in the jack mackerel mid-water trawl fishery on the WCSI and WCNI, which has issues with bycatch of kingfish and snapper. Incidental interactions and associated mortality of common dolphins occur in the jack mackerel fishery. Interactions with other species are currently being characterised.

Table 7: Summary of reported landings (*t*) and TACCs by QMA for the most recent fishing year.


Fishstock	FMA	2016–17 TACC	2016–17 Reported Landings
EMA 1	1	7 630	7 551
EMA 2	2	180	83
EMA 3	3–6	390	126
EMA 7	7–9	3 350	625
EMA 10	10	0	0
TOTAL		11 550	8 385

6. FOR FURTHER INFORMATION

- Ballara, S (2016). Characterisation analyses for blue mackerel (*Scomber australasicus*) in EMA 1, 2, 3, and 7, 1989–90 to 2013–14. *New Zealand Fisheries Assessment Report 2016/04*. 108 p.
- Boyd, R O; Gowing, L; Reilly, J L (2004) 2000–2001 national marine recreational fishing survey: diary results and harvest estimates. Draft New Zealand Fisheries Assessment Report (Unpublished report held by Fisheries New Zealand, Wellington.). 93 p.
- Boyd, R O; Reilly, J L (2005) 1999/2000 National marine recreational fishing survey: harvest estimates. Final Research Report for Ministry of Fisheries Research Project REC9803. (Unpublished report held by Fisheries New Zealand, Wellington.).
- Bradford, E (1996) A comparison of the 1993–94 diary and boatramp surveys of recreational fishing in the Ministry of Fisheries North region. New Zealand Fisheries Assessment Research Document 1996/5. 21 p. (Unpublished report held by NIWA library, Wellington.)
- Bradford, E (1998) Harvest estimates from the 1996 national marine recreational fishing surveys. New Zealand Fisheries Assessment Research Document 1998/16. 27 p. (Unpublished report held by NIWA library, Wellington.)
- Bradford, E; Taylor, P R (1995) Trends in pelagic fish abundance from aerial sightings data. New Zealand Fisheries Assessment Research Document 1995/8. 60 p. (Unpublished report held by NIWA library, Wellington.)
- Fu, D (2013). Characterisation analyses for blue mackerel (*Scomber australasicus*) in EMA 1, 2, 3, and 7, 1989–90 to 2009–10. *New Zealand Fisheries Assessment Report 2013/16*. 54 p.
- Fu, D; Taylor, P R (2007) Standardised CPUE analysis for blue mackerel (*Scomber australasicus*) in EMA 7, 1989–90 to 2004–05. *New Zealand Fisheries Assessment Report 2007/33*. 33 p.
- Fu, D; Taylor, P R (2011) Characterisation and standardised CPUE analyses for blue mackerel (*Scomber australasicus*) in EMA 7, 1989–90 to 2008–09. *New Zealand Fisheries Assessment Report 2011/56*.
- Hartill, B; Davey, N (2015) Mean weight estimates for recreational fisheries in 2011–12. *New Zealand Fisheries Assessment Report 2015/25*.
- Langley, A; Anderson, F (1998) Sampling the length and age composition of EMA 1 blue mackerel catches from the 1997–98 target purse seine fishery. Unpublished research report, held by Sanfords Tauranga Ltd.
- Manning, M J; Devine, J A; Marriott, P M; Taylor, P R (2007) The length and age composition of the commercial catch of blue mackerel (*Scomber australasicus*) in EMA 1 and EMA 7 during the 2004–05 fishing year. *New Zealand Fisheries Assessment Report 2007/35*. 36 p.
- Manning, M J; Marriot, P M; Taylor, P (2006) The length and age composition of the commercial catch of blue mackerel (*Scomber australasicus*) in EMA 1 during the 2002–03 fishing year, including a comparison with data collected during the 1997–98 fishing year, and some remarks on estimating blue mackerel ages from otoliths. *New Zealand Fisheries Assessment Report 2006/42*. 42 p.
- Manning, M J; Marriot, P M; Taylor, P R (2007) Length and age composition of the commercial catch of blue mackerel (*Scomber australis*) in EMA 1 and 7 during the 2003–04 fishing year. *New Zealand Fisheries Assessment Report 2007/13*. 41 p.
- Marriott, P M; Manning, M J (2011) Reviewing and refining the method for estimating blue mackerel (*Scomber australasicus*) ages. *New Zealand Fisheries Assessment Report 2011/11*.
- Marriott, P M; Manning, M J; Andrews, A H (2010) Investigating blue mackerel (*Scomber australasicus*) age-estimation error. Final Research Report for Ministry of Fisheries Research Project EMA2005-02, Objectives 2–3. 42 p. (Unpublished document held by Fisheries New Zealand, Wellington.)
- Morrison, M; Taylor, P; Marriott, P; Sutton, C (2001) An assessment of information on blue mackerel (*Scomber australasicus*) stocks. *New Zealand Fisheries Assessment Report 2001/44*. 26 p.
- Parrish, R H; MacCall, A D (1978) Climatic variation and exploitation in the Pacific mackerel fishery. *California Department of Fish and Game Bulletin* 167. 110 p.
- Robertson, D A (1978) Blue Mackerel, pilchard, anchovy, sprat, saury, and lantern fish. *New Zealand Fisheries Research Division Occasional Publication* 15. 85–89.
- Rohde, K (1987) Different populations of *Scomber australasicus* in New Zealand and south-eastern Australia, demonstrated by a simple method using monogenean sclerites. *Journal of Fish Biology* 30(6): 651–657.
- Sette, O E (1950) Biology of the Atlantic mackerel (*Scomber scombrus*) of North America. Part II. Migrations and habits. *Fishery Bulletin of the United States Fish and Wildlife Service* 51(49): 251–358.
- Shaun-ror, W (1970) Age and growth of Taiwan spotted mackerel, *S. australasicus*. MSc Thesis, Department of Zoology, National Taiwan University. 38 p. [In Chinese].
- Smith, P J; Diggles, B; Kim, S (2005) Stock structure of blue mackerel, *Scomber australasicus*. *New Zealand Fisheries Assessment Report 2005/43*. 38 p.
- Stevens, J D; Hausfeld, H F; Davenport, S R (1984) Observations on the biology, distribution and abundance of *Trachurus declivis*, *Sardinops neopilchardus* and *Scomber australasicus* in the Great Australian Bight. *Report (CSIRO Marine Laboratories) No. 164*. 29 p.
- Stewart, J; Ferrell, D J; Andrew, N L (1998) Ageing yellowtail (*Trachurus novaezelandiae*) and blue mackerel (*Scomber australasicus*) in New South Wales. *New South Wales Fisheries Final Report Series, No. 3*. 59 p.
- Stewart, J; Ferrell, D J; Andrew, N L (1999) Validation of the formation and appearance of annual marks in the otoliths of yellowtail (*Trachurus novaezelandiae*) and blue mackerel (*Scomber australasicus*) in New South Wales. *Australian Journal of Marine and Freshwater Research* 50: 389–395.
- Taylor, P R (2002) A summary of information on blue mackerel, (*Scomber australasicus*), characterisation of its fishery in QMAs 7, 8, and 9, and recommendations on appropriate methods to monitor the status of its stock. *New Zealand Fisheries Assessment Report 2002/50*. 68 p.
- Taylor, P R (2008) Factors affecting fish size and landed volumes in the purse-seine and TCEPR charter-boat fisheries in 2004–05 and 2005–06. *New Zealand Fisheries Assessment Report 2008/32*. 17 p.
- Wynne-Jones, J; Gray, A; Hill, L; Heinemann, A (2014) National Panel Survey of Marine Recreational Fishers 2011–12: Harvest Estimates. *New Zealand Fisheries Assessment Report 2014/67*.

BLUE MOKI (MOK)*(Latridopsis ciliaris)*

Moki

**1. FISHERY SUMMARY****1.1 Commercial fisheries**

Most blue moki landings are taken by setnet or trawl on the east coast between the Bay of Plenty (BoP) and Kaikoura, although small quantities are taken in most New Zealand coastal waters. While the proportions of the total commercial landings taken by setnet and trawl have varied over time, setnetting has been the predominant method, accounting for 50–60% of the annual catch during 1989–90 to 2011–12. The proportion of the catch taken by set net declined in the more recent years (to 2015–16) and catches by the two methods were at about parity during this period.

Reported landings and TACCs are given in Tables 1 and 2, while an historical record of landings and TACC values for the two main MOK stocks are depicted in Figure 1. Landings of blue moki peaked in 1970 and 1979 at about 960 t. Blue moki stocks appeared to have been seriously depleted by fishing prior to 1975 and this resulted in the sum of allocated ITQs being markedly less than the sum of the catch histories.

Table 1: Total reported landings (t) of blue moki from 1979 to 1985–86.

Year	1979*	1980*	1981*	1982*	1983†	1983–84†	1984–85†	1985–86†
Landings	957	919	812	502	602	766	642	636

*MAF data.

†FSU data.

Total annual landings of blue moki were substantially constrained when it was introduced into the QMS. In MOK 1, landings increased as the TACC was progressively increased. Since the TACC was set at 400 t (1995–96) landings have fluctuated around the TACC, which was subsequently increased to 403 t in 2001–02.

Landings from MOK 3 increased from the mid 2000s and exceeded the TACC of 127 t from 2010–11. The TACC was increased to 160 t in 2014–15.

The combined MOK 1 and 3 catch fluctuated around 500 t per annum during 1994–95 to 2009–10. Since then annual catches have been about 550 t.

BLUE MOKI (MOK)

Table 2: Reported landings (t) and actual TACCs (t) of blue moki by Fishstock from 1986–87 to 2016–17. Source – QMS data. MOK 10 is not tabulated; no landings have ever been reported from MOK 10.

Fishstock FMA (s)	MOK 1		MOK 3		MOK 4		MOK 5		Total	
	1,2,7,8,9		3		4		5 & 6			
	Landings	TACC	Landings	TACC	Landings	TACC	Landings	TACC	Landings	TACC
1986–87	109	130	52	60	0	20	3	40	164	260
1987–88	183	142	95	62	0	20	2	40	280	274
1988–89	134	151	121	64	0	20	3	40	258	285
1989–90	202	156	89	65	11	25	1	43	303	299
1990–91	264	157	93	71	1	25	2	43	360	306
1991–92	285	157	66	71	2	25	2	43	355	306
1992–93	289	157	94	122	1	25	4	43	388	358
1993–94	374	200	102	126	4	25	5	43	485	404
1994–95	418	200	90	126	< 1	25	3	43	511	404
1995–96	435	400	91	126	1	25	3	43	530	604
1996–97	408	400	66	126	2	25	3	43	479	604
1997–98	416	400	78	126	3	25	2	43	500	604
1998–99	468	400	78	126	< 1	25	4	43	551	604
1999–00	381	400	56	126	1	25	5	43	443	604
2000–01	420	400	67	126	5	25	6	43	499	604
2001–02	365	403	77	127	8	25	2	44	451	608
2002–03	380	403	87	127	2	25	6	44	475	608
2003–04	372	403	60	127	2	25	6	44	440	608
2004–05	418	403	70	127	3	25	11	44	502	608
2005–06	408	403	69	127	1	25	5	44	483	608
2006–07	402	403	90	127	< 1	25	11	44	504	608
2007–08	401	403	125	127	< 1	25	8	44	533	608
2008–09	413	403	103	127	1	25	8	44	525	608
2009–10	386	403	129	127	< 1	25	6	44	521	608
2010–11	421	403	144	127	< 1	25	10	44	574	608
2011–12	427	403	137	127	< 1	25	6	44	571	608
2012–13	385	403	159	127	< 1	25	5	44	549	608
2013–14	393	403	134	127	<1	25	7	44	535	608
2014–15	376	403	146	160	<1	25	6	44	529	631
2015–16	395	403	183	160	<1	25	8	44	587	631
2016–17	387	403	162	160	<1	25	7	44	556	631




Figure 1: Reported commercial landings and TACC for the two main MOK stocks: MOK 1 (Auckland, Central, and Challenger). Note: these figures do not show data prior to entry into the QMS.




Figure 1 [Continued]: Reported commercial landings and TACC for the two main MOK stocks: MOK 3 (South East Coast). Note: these figures do not show data prior to entry into the QMS.

1.2 Recreational fisheries

Popular with recreational fishers, blue moki are taken by beach anglers, setnetting and spearfishing. Annual estimates of recreational harvest were obtained from diary surveys in 1991–94, 1996 and 1999–2000 (Tables 3a and 3b).

Table 3a: Estimated number and weight of blue moki harvested by recreational fishers by Fishstock and survey. Surveys were carried out in different years in the MAF Fisheries regions: South in 1991–92, Central in 1992–93 and North in 1993–94 (Teirney et al 1997).

Fishstock	Survey	Number	CV(%)	Survey harvest (t)
MOK 1	North	6 000	-	5–15
MOK 1	Central	38 000	28	40–80
MOK 1	South	2 000	-	0–5
MOK 3	South	31 000	33	40–70
MOK 5	South	7000	33	5–15

Table 3b: Estimates of annual number and weight of blue moki harvested by recreational fishers from national diary surveys in 1996 (Bradford 1998) and Dec1999–Nov 2000 (Boyd & Reilly 2005). The mean weights used to convert numbers to catch weight are considered the best available estimates. Estimated harvest is also presented as a range to reflect the uncertainty in the point estimates.

Fishstock	Number caught	CV	Estimated harvest range (t)	Point estimate (t)
				1996
MOK 1	63 000	14	80–110	93
MOK 3	16 000	18	20–30	24
MOK 5	9000	-	-	-
				1999–2000
MOK 1	81 000	37	82–180	131
MOK 3	36 000	32	36–70	53
MOK 5	38 000	89	7–115	61

The harvest estimates provided by telephone-diary surveys between 1993 and 2001 are no longer considered reliable for various reasons. A Recreational Technical Working Group concluded that these harvest estimates should be used only with the following qualifications: a) they may be very inaccurate; b) the 1996 and earlier surveys contain a methodological error; and c) the 2000 and 2001 estimates are implausibly high for many important fisheries. In response to these problems and the cost and scale challenges associated with onsite methods, a National Panel Survey was conducted for the first time throughout the 2011–12 fishing year. The panel survey used face-to-face interviews of a random sample of 30 390 New Zealand households to recruit a panel of fishers and non-fishers for a full year. The panel members were contacted regularly about their fishing activities and harvest information collected in standardised phone interviews. Harvest estimates for blue moki are given in Table 4 (from Wynne-Jones et al 2014 and Hartill & Davey 2015).

BLUE MOKI (MOK)

Table 4: Recreational harvest estimates for blue moki stocks (Wynne-Jones et al 2014). Mean fish weights were obtained from boat ramp surveys; for blue moki the value used was 2.026 kg (Hartill & Davey 2015).

Stock	Year	Method	Number of fish	Total weight (t)	CV
MOK 1	2011/12	Panel survey	21 945	44.5	0.31
MOK 3	2011/12	Panel survey	5 739	11.6	0.53
MOK 5	2011/12	Panel survey	243	0.5	1.02
MOK total	2011/12	Panel survey	27 927	56.6	0.28

A repeat of the National Panel Survey is being conducted over the 2017–18 October fishing year. Results are expected in early 2019.

1.3 Customary non-commercial fisheries

A traditional Maori fishery exists in some areas, particularly the eastern BoP and East Cape regions. No quantitative information is available on the level of customary non-commercial catch.

Iwi in the Cape Runaway area have a strong view that blue moki are of special significance in the history and life of the community. They believe that blue moki come to spawn in the waters around Cape Runaway and there are traditional fishing grounds, where in earlier years fishing took place in accordance with customary practices. In addition, these local Iwi consider the taking of blue moki by nets in this area to be culturally offensive.

Since September 1996, fishing by the methods of trawling, Danish seining and setnetting has been prohibited at all times within a two nautical-mile wide coastal band beginning at the high water mark and extending from Cape Runaway to a stream tributary at Oruiti Beach. Note this is not a legal description, for full details please refer to the Fisheries Act (Auckland and Kermadec Areas Commercial Fishing Regulations 1986, Amendment No. 13).

1.4 Illegal catch

No quantitative estimates are available.

1.5 Other sources of mortality

Some blue moki caught for use as rock lobster bait have not been reported. While little information is available, this practice appears to have been most common in Stewart Island and the Chatham Islands, and may have accounted for about 45 t and 60 t in Stewart and Chatham respectively in the past. The use of blue moki as bait has not been considered in the determination of *MCY*.

2. BIOLOGY

Blue moki grow rapidly at first, attaining sexual maturity at 40 cm fork length (FL) at 5–6 years of age. Growth then slows, and fish of 60 cm FL are 10–20 years old. Fish over 80 cm FL and 43 years old have been recorded (Manning et al 2009).

Many adults take part in an annual migration between Kaikoura and East Cape. The migration begins off Kaikoura in late April/May as fish move northwards. Spawning takes place in August/September in the Mahia Peninsula to East Cape region (the only known spawning ground), with the fish then returning south towards Kaikoura. The larval phase for blue moki lasts about 6 months.

Juvenile blue moki are found inshore, usually around rocky reefs, while most adults school offshore over mainly open bottom. Some adults do not join the adult schools but remain around reefs.

Biological parameters relevant to the stock assessment are shown in Table 5.

Table 5: Estimates of biological parameters for blue moki.

Fishstock	Estimate		Source
1. Natural mortality (M)			
All areas	0.14		Francis (1981b)
For maximum observed age of 33 yr.			
MOK 1	0.10		Manning et al (2009)
For maximum observed age of 44 yr.			
2. Weight = $a(\text{length})^b$ (Weight in g, length in cm fork length).			
	Both sexes		
	a	b	
All areas	0.055	2.713	Francis (1979)
3. von Bertalanffy growth parameters			
	Both sexes		
	L_∞	k	t_0
All areas	66.95	0.208	-0.029
			Francis (<i>pers. comm.</i>)

The estimate of natural mortality, given a maximum age of 43 years and using the equation $M = \log_e 100/\text{maximum age}$, is 0.1. Note that the maximum age for this calculation is meant to be the maximum age that 1% of the unfished population will reach, however, as this is not known, the maximum observed age was used here.

3. STOCKS AND AREAS

There are no new data which would alter the stock boundaries given in previous assessment documents.

Blue moki forms one stock around the North Island and the South Island north of Banks Peninsula. No information is available to indicate stock affiliations of blue moki in other areas (southern South Island and Chatham Rise) so these fish are currently divided into three Fishstocks.

4. STOCK ASSESSMENT

There are no new data which would alter the yield estimates given in the 1996 Plenary Report. The yield estimates are based on commercial landings data only and have not changed since the 1992 Plenary Report.

4.1 Estimates of fishing mortality

Estimates of total mortality (Z) for MOK 1 were obtained from catch curve analysis of catch sampling data collected during 2004–05 and 2005–06. Samples were taken from both the target setnet fishery and from bycatch from the TAR 2 trawl fishery. When data were pooled across the two years, sexes and fishing methods, Z estimates ranged from 0.11 to 0.14, depending on assumed age-at-full recruitment (ages 4–12 years were tested). Assuming a value of natural mortality of 0.10 (based on a maximum age of 44 years), this suggests that recent fishing mortality is likely to be in the range of about 0.01 to 0.04. The Working Group considered that the most plausible age-at-full recruitment was 8 years. The estimate of Z and the bootstrapped 95% confidence intervals were 0.14 (0.12–0.16), giving rise to a F estimate of 0.04 (0.02–0.06). These estimates are well below the current assumed value of natural mortality (Manning et al 2009).

4.2 CPUE analyses

In 2017, a summary of the recent trends in catch from the MOK 1 and MOK 3 fisheries was presented to the Southern Inshore Fishery Assessment Working Group (Langley 2018). The analysis identified three main fisheries catching blue moki:

BLUE MOKI (MOK)

1. The tarakihi bottom trawl fishery operating within the Gisborne-Mahia area (Statistical Area 013) throughout the year.
2. The target blue moki set net fishery operating between East Cape and Wairarapa (Statistical Areas 014–015) primarily during May–October.
3. The Kaikoura set net fishery (Statistical Area 018) operating during May–June and October.

For each fishery, a standardised CPUE analysis was conducted for 1989–90 to 2015–16. All three CPUE analyses modelled the positive catch of blue moki assuming a lognormal error structure, while the CPUE analysis of the tarakihi bottom trawl fishery (BT-TAR2-North) also modelled the presence of blue moki in the catch and derived delta-lognormal CPUE indices.




Figure 2: CPUE indices and 95% confidence intervals from the three main MOK 1 and MOK 3 fisheries.

The SN-MOK3 CPUE indices increased from a relatively low level in 1996–97 to 1999–2000 to reach the highest level of the time series in 2015–16. The SN-MOK1 CPUE indices increased during the same period although the CPUE indices are considerably more variable among years and are less well determined than the SN-MOK3 CPUE indices. The higher variability in the SN-MOK1 indices appears to be related to the inter-annual variation in the operation of the fishery (between Statistical Areas) and limited continuity in the core set of vessels participating in the fishery.

The SINSWG rejected the SN-MOK1 and SN-MOK3 CPUE indices as monitoring tools which could be used to determine stock status against Harvest Standard reference points, for the following reasons:

1. High inter-annual variation in the CPUE indices due to the low precision of CPUE indices derived from limited catch/effort data sets from these small fisheries and/or inter-annual variation in the catchability (availability) of migrating fish.
2. Possible hyperstability as a result of fishing directed at dense schools of migrating fish.

The WG nevertheless agreed that the SN-MOK1 and SN-MOK3 CPUE indices were likely to be broadly indicative of trends in abundance.

The two sets of SN CPUE indices are considered to represent the component (or components) of the blue moki stock migrating northward prior to spawning and then returning southward following spawning. These CPUE indices indicate that there has been a general increase in the abundance of adult blue moki within MOK 3 and the southern area of MOK 1 from the late 1990s. This is consistent with the estimates of total mortality derived from the population age structure in 2005–06 that indicated that fishing mortality on the adult population was less than natural mortality (M).

The BT-TAR2-North CPUE indices contrast the trend in the CPUE indices from the two set net fisheries. The BT-TAR2-North CPUE indices declined from 1996–97 to 2002–03 and remained at a relatively low level during 2002–03 to 2008–09. The index increased in 2009–10 and remained at about that level during 2010–11 to 2015–16. These recent indices are at a level considerably lower than the indices from 1989–90 to 1996–97 (with the exception of the low 1992–93 index).

The BT-TAR2-North CPUE indices are considered to be predominantly comprised of a component of the blue moki stock that remains in the Gisborne-Mahia area throughout the year. The trawl catch is probably comprised of both immature and mature blue moki, although limited sampling of this component of the stock was conducted during the catch sampling programme. The SINSWG considered that the BT-TAR2-North CPUE series potentially provides an index of abundance for the resident portion of the population, but did not provide a monitoring tool for the entire population.

The contrasting trends in the CPUE indices (SN-MOK1 and SN-MOK3 versus BT-TAR-North) are indicative of differences in the stock dynamics (recruitment and/or exploitation) in the two components of the stock (resident and migrating). It was not considered feasible to amalgamate the three sets of CPUE indices to derive a composite set of abundance indices for the MOK 1&3 stock as the relative proportion of the stock biomass monitored by each CPUE series is unknown. Thus, the utility of the CPUE series is limited to the monitoring each component of the stock separately.

4.3 Biomass estimates

Estimates of current and reference biomass are not available.

4.4 Yield estimates and projections

MCY for all Fishstocks combined was estimated using the equation, $MCY = cY_{AV}$ (Method 4). The national catch, and probably effort, over the period 1961–86 varied considerably (annual landings ranged from 450 to 957 t with an average value of 705 t). However, no clear trend in landings over that period is apparent. The value of *c* was set equal to 0.9 based on the estimate of $M = 0.14$.

$$MCY = 0.9 * 705 \text{ t} = 635 \text{ t}$$

The level of risk to the stock by harvesting the population at the estimated *MCY* value cannot be determined.

Yield estimates for blue moki have been made using reported commercial landings data only and therefore apply specifically to the commercial fishery. Blue moki have been caught and used as bait and not reported. Therefore, the *MCY* estimates are likely to be conservative.

No estimate of *CAY* is available for blue moki stocks.

4.5 Other factors

CPUE data from the 1970s for the main northern blue moki stock indicated that the stock had declined to a level low enough to make recruitment failure a real concern. The 1986–87 TAC was set at a level considered low enough to enable some stock rebuilding.

Blue moki forms one stock around the North Island and the east coast of the South Island north of Banks Peninsula. As other stock boundaries are unknown, any interdependence is uncertain. If only one stock exists, then blue moki from the southern waters may be moving north and rebuilding the heavily exploited northern population.

5. STATUS OF THE STOCKS

Stock Structure Assumptions

Blue moki forms one stock around the North Island and the South Island north of Banks Peninsula. The bulk of the commercial catch is taken off the east coast between Banks Peninsula and East Cape, suggesting that this is where most of the blue moki stock resides.

MOK 1&3

Stock Status	
Year of Most Recent Assessment	2017
Assessment Runs Presented	2008 – Catch-at-age 2017 – Three CPUE series

BLUE MOKI (MOK)

Reference Points	Target: Not established but $F = M$ assumed Soft Limit: 20% B_0 Hard Limit: 10% B_0 Overfishing threshold: Not established but $F = M$ assumed
Status in relation to Target	F is Very Likely ($> 90\%$) to be below M
Status in relation to Limits	Soft Limit: Unknown Hard Limit: Unknown
Status in relation to overfishing	F is Very Unlikely ($< 10\%$) to be above M

Historical Stock Status Trajectory and Current Status	-
--	---

Fishery and Stock Trends	
Recent Trend in Biomass or Proxy	Catch curve analysis from catch sampling the migratory adult population (2004–05 and 2005–06) indicated that total mortality was low, with fishing mortality well below natural mortality. The general increase in CPUE from the SN-MOK1 and SN-MOK3 fisheries suggests that the biomass of migratory adults has increased since then.
Recent Trend in Fishing Intensity or Proxy	Low estimates of fishing mortality in 2005–06 and stable catches over the previous 14 years suggest that fishing mortality had been low for more than two decades. Recent increases in CPUE suggest that adult biomass has increased since the catch-at-age study, and together with constant catch suggests that fishing mortality remains below the target.
Other Abundance Indices	-
Trends in Other Relevant Indicators or Variables	CPUE indices from three fisheries are not considered to be sufficiently reliable to represent abundance indices for the stock. Rather, the indices are considered to be indicative of general trends in abundance for components of the stock. The SN-MOK1 and SN-MOK3 CPUE indices indicate that there has been a general increase in the abundance of adult blue moki within MOK 3 and the southern area of MOK 1 from the late 1990s. By contrast the BT-TAR2N series suggests that resident MOK in the northern part of FMA2 (Mahia Peninsula) declined to the mid-2000s and then increased to 2010–11, after which it fluctuated without trend at a level approximately half of that in the early 1990s.

Projections and Prognosis	
Stock Projections or Prognosis	If catches remain at current levels then fishing mortality should remain below the target.
Probability of Current Catch or TACC causing Biomass to remain below or to decline below Limits	Soft Limit: Unknown Hard Limit: Unknown
Probability of Current Catch or TACC causing Overfishing to continue or to commence	Fishing mortality was estimated to be below the target fishing mortality level (M) in the mid-2000s. Since then, there has been a general increase in stock abundance of the migrating adult component of the stock (as indicated by the CPUE trends). It is therefore Unlikely ($< 10\%$) that fishing mortality will exceeds the overfishing threshold at current catch levels.

Assessment Methodology and Evaluation		
Assessment Type	Level 2 - Partial Quantitative stock assessment	
Assessment Method	Estimates of total mortality using Chapman-Robson estimator	
Assessment Dates	Latest assessment: 2017	Next assessment: Unknown
Overall assessment quality rank	1 – High Quality	
Main data inputs (rank)	- Age structure of setnet and trawl catches of blue moki made between Kaikoura and East Cape in 2004–05 and 2005–06 -Instantaneous rate of natural mortality (M) of 0.10 based on a maximum age of 44 years -CPUE indices for migrant components of the stock (SN-MOK1 and SN-MOK3 CPUE)	1 – High Quality 2 – Medium or Mixed Quality: uncertainty in estimate of M 2 – Medium or Mixed Quality: may not be fully representative
Data not used (rank)	N/A	
Changes to Model Structure and Assumptions	-	
Major Sources of Uncertainty	Uncertainty in the estimate of M Reliability of CPUE indices as indices of stock abundance.	

Qualifying Comments
-
Fishery Interactions
Interactions with other species are currently being characterised.

Yields and reported landings are summarised in Table 6.

Table 6: Summary of yields (t), TACCs (t), and reported landings (t) for blue moki for the most recent fishing year.


Fishstock	QMA	MCY	2016–17 Actual TACC	2016–17 Reported landings
MOK 1	Auckland (East) (West),			
MOK 3	Central (East) (West), Challenger 1, 2, 7, 8 & 9	-	403	387
MOK 4	South East (Coast) 3	-	160	162
MOK 5	South East (Chatham) 4	-	25	< 1
MOK 10	Southland, Sub-Antarctic 5 & 6	-	44	7
	Kermadec 10	-	10	0
Total		635	631	556

6. FOR FURTHER INFORMATION

- Boyd, R O; Reilly, J L (2005) 1999/2000 National marine recreational fishing survey: harvest estimates. Final Research Report for Ministry of Fisheries Research Project REC9803. (Unpublished report held by Fisheries New Zealand, Wellington.)
- Bradford, E (1998) Harvest estimates from the 1996 national marine recreational fishing surveys. New Zealand Fisheries Assessment Research Document 1998/16. 27 p. (Unpublished document held by NIWA library, Wellington.)
- Colman, J A; McKoy, J L; Baird, G G (Comp. & Ed.) (1985) Background papers for the 1985 Total Allowable Catch recommendations. Fisheries Research Division, New Zealand Ministry of Agriculture and Fisheries. 259 p. (Unpublished report, held by NIWA library, Wellington.)
- Francis, M P (1979) A biological basis for the management of New Zealand moki (*Latridopsis ciliaris*) and smoothhound (*Mustelus lenticulatus*) fisheries. (Unpublished MSc thesis, University of Canterbury, Christchurch, New Zealand).
- Francis, M P (1981a) Spawning migration of moki (*Latridopsis ciliaris*) off eastern New Zealand. *New Zealand Journal of Marine and Freshwater Research* 15: 267–273.
- Francis, M P (1981b) Age and growth of moki, *Latridopsis ciliaris* (Teleostei: Latridae). *New Zealand Journal of Marine and Freshwater Research* 15: 47–49.
- Hartill, B; Davey, N (2015) Mean weight estimates for recreational fisheries in 2011–12. *New Zealand Fisheries Assessment Report* 2015/25.
- Horn, P (1988) Blue moki. New Zealand Fisheries Assessment Research Document 1988/10. 11 p. (Unpublished document held by NIWA library, Wellington.)
- Langley, A.D. (2018). Fishery characterisation and Catch-Per-Unit-Effort analysis for blue moki (*Latridopsis ciliaris*) in MOK 1 and MOK 3. *New Zealand Fisheries Assessment Report* 2018/1. 52 p.
- Langley, A D; Walker, N (2004) Characterisation of the blue moki (*Latridopsis ciliaris*) fishery and recommendations for future monitoring of the MOK 1 Fishstock. *New Zealand Fisheries Assessment Report* 2004/33. 77 p.
- Manning, M J; Stevenson, M L; Dick, C M (2009) The length and age composition of the commercial catch of blue moki (*Latridopsis ciliaris*) in MOK 1 during the 2004/05 and 2005/06 fishing years including total and fishing mortality estimates. *New Zealand Fisheries Assessment Report* 2010/34. 65 p.
- Teirney, L D; Kilner, A R; Millar, R E; Bradford, E; Bell, J D (1997) Estimation of recreational catch from 1991/92 to 1993/94 New Zealand Fisheries Assessment Research Document 1997/15. 43 p. (Unpublished document held by NIWA library, Wellington.)
- Wynne-Jones, J; Gray, A; Hill, L; Heinemann, A (2014) National Panel Survey of Marine Recreational Fishers 2011–12: Harvest Estimates. *New Zealand Fisheries Assessment Report* 2014/67.

BLUE WAREHOU (WAR)

(*Seriolella brama*)
Warehou

**1. FISHERY SUMMARY****1.1 Commercial fisheries**

Blue (or common) warehou are caught in coastal waters of the South Island and lower North Island down to depths of about 400 m. Annual landings were generally less than 100 t up to the early 1960s, increased to about 1000 t by the early 1970s, and peaked at 4387 t in 1983–84 before declining steadily through to 1988–89 (Table 2). Figure 1 shows the the historical landings and TACC values for the main WAR stocks.

The decline was most notable in WAR 3, from which most of the catch is recorded. A TACC reduction for WAR 3, from 3357 to 2528 t, was approved for the 1990–91 fishing year. In 1990–91, total catch increased substantially. The largest increase was in WAR 3 and catches in this area exceeded 2000 t for the following three years. There is no direct correlation between WAR 3 catches and fluctuations in effort in the Snares squid fishery where blue warehou is mostly taken as bycatch. In 1996–97, total catch increased again to 1990–91 levels and total catch has been maintained at this level since. Increased catches in WAR 2, 3 and 7 contributed to the increased total catch.

Until the mid 1980s, the main domestic fishing method used to catch blue warehou was gill-netting. The majority of the landings are now taken as a bycatch from trawling. Bull & Kendrick (2006) describe the commercial fishery from 1989–90 to 2002–03.

Catches have fluctuated in most stocks but overall the total landings have increased. In 2002–03, total reported landings of blue warehou were the highest on record, with catches in WAR 3 exceeding the TACC by 983 t. From 2002–03 to 2006–07 catches in WAR 3 were well above the TACC as fishers landed catches well in excess of ACE holdings and paid deemed values for the overcatch. From 1 October 2007 the deemed values were increased to \$0.90 per kg for WAR 3 and WAR 7 stocks and differential rates were also introduced. The differential rate applied to all catch over 110% of ACE holding at which point the deemed value rate increased to \$2 per kg. The effect of these measures was seen immediately in 2007–08 as fishing without ACE was reduced and catch fell well below the TACC in WAR 3. In all other areas landings are below the TACCs.

BLUE WAREHOU (WAR)
Table 2: Reported landings (t) of blue warehou by Fishstock 1983–84 to 2016–17 and actual TACCs (t) from 1986–87 to 2016–17. QMS data from 1986–present. [Continued on next page.]

Fishstock FMA	WAR 1 1 & 9		WAR 2 2		WAR 3 3, 4, 5 & 6		WAR 7 7	
	Landings	TACC	Landings	TACC	Landings	TACC	Landings†	TACC
1983–84*	13	-	346	-	3 222	-	702	-
1984–85*	5	-	278	-	1 313	-	478	-
1985–86*	15	-	185	-	1 584	-	955	-
1986–87	7	30	190	480	1 330	3 210	780	910
1987–88	7	41	204	560	976	3 223	685	962
1988–89	12	41	177	563	672	3 348	561	969
1989–90	17	41	201	570	814	3 357	607	1 047
1990–91	14	41	250	570	2 097	2 528	758	1 117
1991–92	25	41	235	570	2 514	2 528	1 001	1 117
1992–93	15	41	199	578	2 310	2 530	539	1 120
1993–94	16	41	233	578	688	2 530	436	1 120
1994–95	15	41	203	578	1 274	2 530	468	1 120
1995–96	32	41	368	578	1 573	2 530	756	1 120
1996–97	24	41	563	578	1 814	2 531	1 428	1 120
1997–98	20	41	402	578	2 328	2 531	860	1 120
1998–99	15	41	503	578	1 978	2 531	1 075	1 120
1999–00	9	41	422	578	2 761	2 531	1 147	1 120
2000–01	12	41	388	578	1 620	2 531	1 572	1 120
2001–02	7	41	294	578	1 614	2 531	1 046	1 120
2002–03	5	41	429	578	3 514	2 531	961	1 120
2003–04	6	41	392	578	3 539	2 531	755	1 120
2004–05	6	41	402	578	2 963	2 531	756	1 120
2005–06	4	41	293	578	3 505	2 531	691	1 120
2006–07	4	41	235	578	3 326	2 531	823	1 120
2007–08	7	41	198	578	684	2 531	569	1 120
2008–09	9	41	210	578	2 021	2 531	733	1 120
2009–10	6	41	204	578	2 601	2 531	414	1 120
2010–11	11	41	102	578	2 086	2 531	633	1 120
2011–12	13	41	131	578	2 425	2 531	714	1 120
2012–13	8	41	172	578	1 847	2 531	632	1 120
2013–14	17	41	153	578	1 819	2 531	551	1 120
2014–15	24	41	123	578	2 674	2 531	823	1 120
2015–16	5	41	167	578	1 861	2 531	764	1 120
2016–17	14	41	143	578	2 357	2 531	875	1 120


Fishstock FMA	WAR 8 8		WAR 10 10		Total Total	
	Landings	TACC	Landings	TACC	Landings	TACC
1983–84*	104	-	0	-	4 387	-
1984–85*	91	-	0	-	2 165	-
1985–86*	43	-	0	-	2 782	-
1986–87	40	210	0	10	2 347	4 850
1987–88	43	218	0	10	1 915	5 014
1988–89	44	231	0	10	1 466	5 162
1989–90	57	233	0	10	1 696	5 459
1990–91	113	233	0	10	3 232	4 499
1991–92	132	233	<1	10	3 905	4 499
1992–93	152	233	<1	10	3 215	4 512
1993–94	126	233	0	10	1 500	4 512
1994–95	114	233	0	10	2 074	4 512
1995–96	186	233	0	10	2 913	4 512
1996–97	161	233	0	10	3 990	4 513
1997–98	111	233	0	10	3 720	4 513
1998–99	168	233	0	10	3 739	4 513
1999–00	116	233	0	10	4 455	4 513
2000–01	143	233	0	10	3 735	4 513
2001–02	146	233	0	10	3 107	4 513
2002–03	192	233	0	10	5 101	4 513
2003–04	129	233	0	10	4 821	4 513
2004–05	157	233	0	10	4 284	4 513
2005–06	76	233	0	10	4 569	4 513
2006–07	59	233	0	10	4 448	4 513
2007–08	72	233	0	10	1 530	4 513
2008–09	146	233	0	10	3 119	4 513
2009–10	159	233	0	10	3 384	4 513
2010–11	92	233	0	10	2 924	4 512

Table 2 [Continued]

2011–12	97	233	0	10	3 381	4 512
2012–13	111	233	0	10	2 770	4 512
2013–14	161	233	0	10	2 701	4 512
2014–15	69	233	0	10	3 713	4 512
2015–16	95	233	0	10	2 891	4 512
2016–17	59	233	0	10	3 448	4 512

* FSU data.

† Includes landings from unknown areas before 1986–87.

**Figure 1: Reported commercial landings and TACC for the four main WAR stocks. WAR 2 (Central East), WAR 3 (South East Coast) and WAR 7 (Challenger) [Continued on next page].**

BLUE WAREHOU (WAR)




Figure 1 [Continued]: Reported commercial landings and TACC for the four main WAR stocks. WAR 8 (Central Egmont).

1.2 Recreational fisheries

Estimates of recreational catch in the MAF Fisheries Central and South regions are shown in Table 3. Surveys in the North region in 1993–94 indicated that blue warehou were not caught in substantial quantities.

Table 3: Estimated harvest (t) of blue warehou by recreational fishers. Surveys were carried out in the MAF Fisheries Southern region in 1991–92 and in the Central region in 1992–93.

Fishstock	Survey	Estimated harvest	CV
1991–92			
WAR 3	Southern	10–20	-
1992–93			
WAR 2	Central	10.0	0.62
WAR 7	Central	1.7	0.65
WAR 8	Central	0.6	1.02

Blue warehou harvest estimates from the 1996 national survey were; WAR 2, 7000 fish; WAR 3, 3000 fish and WAR 7, 1000 fish. There are locally important fisheries which will not have been adequately sampled by these surveys.

The harvest estimates provided by telephone-diary surveys between 1993 and 2001 are no longer considered reliable for various reasons. A Recreational Technical Working Group concluded that these harvest estimates should be used only with the following qualifications: a) they may be very inaccurate; b) the 1996 and earlier surveys contain a methodological error; and c) the 2000 and 2001 estimates are implausibly high for many important fisheries. In response to these problems and the cost and scale challenges associated with onsite methods, a National Panel Survey was conducted for the first time throughout the 2011–12 fishing year. The panel survey used face-to-face interviews of a random sample of 30 390 New Zealand households to recruit a panel of fishers and non-fishers for a full year. The panel members were contacted regularly about their fishing activities and harvest information collected in standardised phone interviews. Harvest estimates (in numbers of fish) for blue warehou are given in Table 4 (from Wynne-Jones et al 2014).

Table 4: Recreational harvest estimates for blue warehou stocks (Wynne-Jones et al 2014). Insufficient data on fish weights were obtained from boat ramp surveys to convert numbers caught to tonnes.

Stock	Year	Method	Number of fish	Total weight (t)	CV
WAR 2	2011/12	Panel survey	1 485	-	-
WAR 3	2011/12	Panel survey	483	-	-
WAR total	2011/12	Panel survey	1 968	-	0.80

A repeat of the National Panel Survey is being conducted over the 2017–18 October fishing year. Results are expected in early 2019.

1.3 Customary non-commercial fisheries

No quantitative information is available on the current level of customary non-commercial take.

1.4 Illegal catch

No quantitative information is available on the level of illegal catch.

1.5 Other sources of mortality

No information is available on other sources of mortality.

2. BIOLOGY

Blue warehou average 40–60 cm fork length (FL) and reach a maximum of about 75 cm. Validated ageing of blue warehou shows rapid growth up to the time of first spawning (about 4–5 years), but negligible growth after about 10 years. Female blue warehou grow significantly faster and reach a larger size than males. Maximum recorded ages are 22 years for males, and 21 years for females. The best estimate of M is now considered to be 0.24 (Bagley et al 1998).

Blue warehou feed on a wide variety of prey, mainly salps but also euphausiids, krill, crabs and small squid.

Known spawning areas include the west coast of the South Island (in August–September), Kaikoura (in March, April, May), Southland (in November), and Hawke Bay (in September). Eggs are found in the surface plankton and juvenile fish are believed to occur in inshore areas.

The seasonal pattern of landings suggest that there is a coastal migration of blue warehou. There is a winter/spring fishery for blue warehou at New Plymouth and north Wairarapa, a summer fishery with a small autumn peak at Wellington and a summer/autumn fishery along the east coast South Island. The west coast South Island has a fishery in August/September which picks up again in summer. There is a summer fishery in Tasman Bay.

Biological parameters relevant to the stock assessment are shown in Table 5.

Table 5: Estimates of biological parameters for blue warehou.

Fishstock					Estimate	Source
1. Natural mortality (<i>M</i>)						
WAR 3					0.24	Bagley et al (1998)
2. Weight = a(length) ^b (Weight in g, length in cm total length).						
	Females		Males			
	a	b	a	b		
WAR 3	0.016	3.07	0.015	3.09		Bagley et al (1998)
3. Von Bertalanffy growth parameters						
	Females			Males		
	<i>L_∞</i>	<i>k</i>	<i>t₀</i>	<i>L_∞</i>	<i>k</i>	<i>t₀</i>
WAR 3	66.3	0.209	-0.79	63.8	0.241	-0.46
	Both Sexes					
WAR 1, 2, 7, 8 (part)	65.5	0.169	-1.35			Jones (1994)
WAR 8 (New Plymouth)	57.7	0.314	0.02			Jones (1994)

3. STOCKS AND AREAS

No definite stock boundaries are known; however, Bagley et al (1998), after considering known spawning grounds and seasonal fishing patterns, suggested that there may be four stocks:

- A southern population, mainly off Southland but perhaps extending into the Canterbury Bight. The main spawning time is November in inshore waters east and west of Stewart Island.
- A central eastern population, located on the northeast coast of the South Island and south east coast of the North Island (including Wellington), spawning mainly in the northern area in winter/early spring and also in autumn off Kaikoura.
- A south western population which spawns on the west coast of the South Island in winter.
- A north western population which may spawn off New Plymouth in winter/spring.

The proposed stock structure is tentative and there may be overlap between stocks. The available age and length frequency data are insufficient to compare by area and tagging studies have been minimal (about 150 fish tagged) with no returns.

For modelling WAR 3, the area on the east coast of the South Island south of Banks Peninsula including Southland was assumed to be a single stock. Movement between the west coast of the South Island and Southland is possible but there was no evidence for this from Southland seasonal trawl surveys. Also, the existence of two spawning periods, from August to September off the west coast of the South Island and from November to December in Southland, suggests two separate stocks.

4. STOCK ASSESSMENT

4.1 Estimation of fishery parameters and abundance

Biomass estimates are available from a number of early trawl surveys (Table 6) but the CVs are rather high for the *Shinkai Maru* data. From the age data from the *Tangaroa* Southland trawl surveys (1993–96) it appears that these surveys did not sample the population consistently, as apparently strong year classes did not follow through the time series of surveys.

Table 6: Trawl survey biomass indices (t) and coefficients of variation (CV) for recruited blue warehou.

Fishstock	Area	Vessel	Trip code	Date	Biomass (t)	CV (%)
WAR 3	Southland	<i>Shinkai Maru</i>	SHI8101	Jan–Mar 81	2 100	43
			SHI8201	Mar–May 82	800	62
			SHI8302	Apr–83	4 700	72
			SHI8601	Jun–86	2 000	59
WAR 3	Southland	<i>Tangaroa</i>	TAN9301	Feb–Mar 93	2 297	36
			TAN9402	Feb–Mar 94	1 629	38
			TAN9502	Feb–Mar 95	1 103	38
			TAN9604	Feb–Mar 96	1 615	40

4.2 Biomass estimates

Estimates of current and reference biomass are not available for any blue warehou Fishstocks.

4.3 Yield estimates and projections

MCY was estimated using the equation $MCY = cY_{AV}$ (Method 4) for all stocks. The value of c was set equal to 0.8 based on the revised estimate of $M = 0.24$ from the validated ageing work completed in 1997.

Auckland, Central (East) (WAR 1 and 2)

Average landings into Wellington over the period 1977 to 1983 were relatively stable at 300 t. Landings along the east coast of the North Island have shown large fluctuations. At Gisborne landings increased from 2 t in 1978 to 140 t in 1979 before declining to 2 t again in 1983. In Napier landings fluctuated from 1 t in 1960 to 87 t in 1972, decreased to less than 20 t in 1975 before peaking at 123 t in 1978 and then declining to 30–40 t. Y_{AV} for Central (East) (FMA 2) was estimated as 300–350 t.

$$\begin{aligned}
 MCY &= 0.8 \times (300 \text{ t} - 350 \text{ t}) \\
 &= 240 \text{ t} - 280 \text{ t}
 \end{aligned}$$

South-east (south of Banks Peninsula), Southland, and Sub-Antarctic (WAR 3)

The catches from 1983–84 to 1985–86 were considered to be a sustainable level of catch. $Y_{AV} = 2040$ t

$$\begin{aligned} MCY &= 0.8 \times 2040 \text{ t} \\ &= 1630 \text{ t} \end{aligned}$$

Challenger (WAR 7)

The catches from 1983–84 to 1985–86 were considered to be a sustainable level of catch. $Y_{AV} = 710$ t.

$$\begin{aligned} MCY &= 0.8 \times 710 \text{ t} \\ &= 570 \text{ t} \end{aligned}$$

Central (West) (WAR 8)

The average domestic landings in the Central (West) zone from 1977 to 1983 were 70 t, and the average (declining) catch over 1983–84 to 1985–86 was 79 t. An *MCY* of 80 t is suggested for this area. New Plymouth has a peak seasonal catch in July, the season extending from June to September.

$$MCY = 80 \text{ t}$$

The level of risk to the stock by harvesting the population at the estimated *MCY* value cannot be determined.

CAY cannot be estimated because of the lack of current biomass estimates.

4.4 Factors modifying yield estimates

No information available.

5. STATUS OF THE STOCKS

Estimates of reference and current biomass are not available.

For all Fishstocks, it is not known if recent landings or TACCs are at levels which will allow the stocks to move towards a size that will support the maximum sustainable yield.

From 2002–03 to 2006–07 catches in WAR 3 were well above the TACC as fishers landed catches well in excess of ACE holdings. Deemed values were increased from 1 October 2007 and landings in WAR 3 in 2007–08 were much reduced to 684 t, well below the current TACC. WAR 3 landings have since increased to more than 2000 t.

Yield estimates, TACCs and reported landings for the 2016–17 fishing year are summarised in Table 7.

Table 7: Summary of yield estimates (t), TACCs (t) and reported landings (t) for blue warehou for the most recent fishing year.


Fishstock		FMA	MCY	2016–17 Actual TACC	2016–17 Reported landings
WAR 1	Auckland (East) (West)	1 & 9	240–280	41	14
WAR 2	Central (East)	2		578	143
WAR 3	South-east (Coast) (Chatham), Southland & Sub-Antarctic	3,4,5 & 6	1 630	2 531	2 357
WAR 7	Challenger	7	570	1 120	875
WAR 8	Central West)	8	80	233	59
WAR 10	Kermadecs	10	0	10	0
Total				4 512	3 448

6. FOR FURTHER INFORMATION

- Bagley, N W; Ballara, S L; Horn, P L; Hurst, R J (1998) A summary of commercial landings and a validated ageing method for blue warehou, *Seriotelella brama* (Centrolophidae), in New Zealand waters, and a stock assessment of the Southern (WAR 3) Fishstock. New Zealand Fisheries Assessment Research Document 1998/20. 46 p. (Unpublished document held by NIWA library, Wellington.)
- Bagley, N W; Hurst, R J (1995) Trawl survey of middle depth and inshore bottom species off Southland, February–March 1994 (TAN 9402). *New Zealand Fisheries Data Report*: 57, 50 p.
- Bagley, N W; Hurst, R J (1996a) Trawl survey of middle depth and inshore bottom species off Southland, February–March 1995 (TAN 9502). *New Zealand Fisheries Data Report*: 73, 47 p.
- Bagley, N W; Hurst, R J (1996b) Trawl survey of middle depth and inshore bottom species off Southland, February–March 1996 (TAN 9604). *New Zealand Fisheries Data Report*: 77, 51 p.
- Bagley, N W; Hurst, R J (1997) A summary of biology and commercial landings, and a stock assessment of white warehou, *Seriotelella caerulea* Guichenot, 1848 (Stromateoidei: Centrolophidae) in New Zealand waters. New Zealand Fisheries Assessment Research Document 1997/13 34 p. (Unpublished document held by NIWA library, Wellington.)
- Bull, B; Kendrick, T H (2006) Fishery characterisations and CPUE analyses for blue warehou (*Seriotelella brama*). Final Research Report for Ministry of Fisheries Project MID2003-01. (Unpublished report held by Fisheries New Zealand, Wellington.)
- Cordue, P L (1995) MIAEL estimation of biomass and fishery indicators for the 1995 assessment of hoki stocks. New Zealand Fisheries Assessment Research Document 1995/13. 38 p. (Unpublished document held by NIWA library, Wellington.)
- Gavrilov, G M (1979) *Seriotelella* of the New Zealand plateau. TINRO, Vladivostok, 1979: 1–79. (Translation No. 204 held in NIWA library, Wellington.)
- Gavrilov, G M; Markina, N P (1970) The feeding ecology of fishes of the genus *Seriotelella* (fam. Nomeidae) on the New Zealand plateau. *Journal of Ichthyology* 19(6): 128–135.
- Horn, P L (1999) Stock assessment of blue warehou *Seriotelella brama* in New Zealand waters. New Zealand Fisheries Assessment Research Document. 1999/26. (Unpublished document held by NIWA library, Wellington.)
- Hurst, R J (1985) Common warehou. In: Colman, J A; McKoy, J L; Baird, G G (Comps. and Eds.) (1985) Background papers for the 1985 Total Allowable Catch recommendations. (Unpublished report, held in NIWA library, Wellington.) pp 63–65.
- Hurst, R J; Bagley, N W (1994) Trawl survey of middle depth and inshore bottom species off Southland, February–March 1993 (TAN 9301). *New Zealand Fisheries Data Report*: 52, 58 p.
- Jones, J B (1988) Blue warehou. New Zealand Fisheries Assessment Research Document 1988/11. 19 p. (Unpublished document held in NIWA library, Wellington.)
- Jones, J B (1994) Notes on warehou age determination. Draft New Zealand Fisheries Assessment Research Document. 15 p.
- Jones, J B; Hurst, R J (1988) Common warehou. In: Baird, G G; McKoy, J L (Comps. and Eds.). Papers from the workshop to review fish stock assessments for the 1987–88 New Zealand fishing year. Pp 67–73.
- McKoy, J L (1988) (Comp.) Blue warehou. Report from the fishery assessment meeting April–May, 1988. Pp 37–39.
- Tsarev, V T (1971) Short notes on *Seriotelella brama* (Gunther) in the south western part of the Pacific Ocean. Izvestiya TINRO [Transactions of the Pacific Ocean Scientific Research Institute for Fisheries and Oceanography] 79: 165–167. [In Russian]
- Wynne-Jones, J; Gray, A; Hill, L; Heinemann, A (2014) National Panel Survey of Marine Recreational Fishers 2011–12: Harvest Estimates. *New Zealand Fisheries Assessment Report 2014/67*.

BLUENOSE (BNS)*(Hyperoglyphe antarctica)*

Matiri

**1. FISHERY SUMMARY**

Bluenose were introduced into the QMS on 1 October 1986. A Total Allowable Catch (TAC) was set under the provisions of the 1983 Fisheries Act, initially at 1350 t. In 2010 new TACs were set for all BNS stocks along with recreational allowances, customary non-commercial allowances, and allowances for other sources of mortality. All current allowances, TACCs and TACs can be found in Table 1 below.

Table 1: Recreational and customary non-commercial allowances, TACCs and TACs by Fishstock (t) for Bluenose.

Fishstock	Recreational allowance	Customary allowance	Other mortality	TACC	TAC
BNS 1	15	2	8	400	425
BNS 2	25	2	9	438	474
BNS 3	18	2	3	171	194
BNS 7	3	2	2	62	69
BNS 8	2	1	1	29	33
BNS 10	-	-	-	10	10

1.1 Commercial fisheries

Bluenose have been landed since the 1930s, although the target line fishery for bluenose only developed in the late 1970s, with the trawl fishery on the lower east coast of the North Island developing after 1983, initially as a bycatch of the alfonsino fishery (Horn 1988a). The largest domestic bluenose fisheries occur in BNS 1 and 2. Historically, catches in BNS 2 were predominately taken in the target alfonsino and bluenose trawl fisheries, but have been primarily taken by target bottom longline fishing in recent years. There is a target line fishery for bluenose in the Bay of Plenty (BoP) and off Northland (BNS 1). Target line fisheries for bluenose also exist off the west coast of the South Island (BNS 7) and the central west coast of the North Island (BNS 8). Bluenose in BNS 7 are also taken as bycatch in the hoki trawl and ling line fisheries. The BNS 3 fishery is focussed on the eastern Chatham Rise where bottom longline catches were historically a bycatch of ling and hapuku target fisheries. Target bluenose lining has predominated since 2003–04. There has been a consistent bycatch of bluenose in the alfonsino target bottom trawl fishery and bluenose have been targeted sporadically in a mid-water trawl fishery in BNS 3 since the early 2000s. The bottom trawl fishery in BNS 3 has diminished. A small amount of target setnet fishing for bluenose occurred in the Bay of Plenty until 1999 and has occurred again since 2012. Target bluenose setnet fishing also occurs sporadically in the Wairarapa region of BNS 2. Setnet catches and off the east coast of the South Island have been a mix of target and bycatch in ling and hapuku target sets.

BLUENOSE (BNS)

Reported landings and TACCs since 1981 are given in Table 2, while the historical landings and TACC for the main BNS stocks are depicted in Figure 1.

Table 2: Reported landings (t) of bluenose by Fishstock from 1981 to 2016–17 and actual TACCs (t) from 1986–87 to 2016–17. QMS data from 1986-present. [Continued on next page]

Fish stock FMA (s)	BNS 1		BNS 2		BNS 3		BNS 7		BNS 8	
	1 & 9		2		3, 4, 5 & 6		7		8	
	Landings	TACC	Landings	TACC	Landings	TACC	Landings	TACC	Landings	TACC
1981*	146		101		36		12		-	
1982*	246		170		46		22		-	
1983†	250		352		51		47		1	
1984†	464		810		81		30		1	
1985†	432		745		73		26		1	
1986†	440		1 009		33		53		1	
1986–87	286	450	953	660	93	150	71	60	1	20
1987–88	405	528	653	661	101	166	104	62	1	22
1988–89	480	530	692	768	90	167	135	69	13	22
1989–90	535	632	766	833	132	174	105	94	3	22
1990–91	696	705	812	833	184	175	72	96	5	22
1991–92	765	705	919	839	240	175	62	96	5	22
1992–93	787	705	1 151	842	224	350	120	97	24	22
1993–94	615	705	1 288	849	311	350	79	97	27	22
1994–95	706	705	1 028	849	389	357	83	150	79	100
1995–96	675	705	953	849	513	357	140	150	70	100
1996–97	966	1 000	1 100	873	540	357	145	150	86	100
1997–98	1 020	1 000	929	873	444	357	123	150	67	100
1998–99	868	1 000	1 002	873	729	357	128	150	46	100
1999–00	860	1 000	1 136	873	566	357	114	150	55	100
2000–01	890	1 000	1 097	873	633	357	87	150	14	100
2001–02	954	1 000	1 010	873	+733	+925	70	150	17	100
2002–03	1 051	1 000	933	873	+876	+925	76	150	66	100
2003–04	1 030	1 000	933	873	915	925	117	150	96	100
2004–05	870	1 000	1 162	1 048	844	925	94	150	42	100
2005–06	699	1 000	1 136	1 048	536	925	84	150	20	100
2006–07	742	1 000	957	1 048	511	925	164	150	50	100
2007–08	585	1 000	1 055	1 048	660	925	145	150	53	100
2008–09	627	786	864	902	444	505	80	89	31	43
2009–10	665	786	845	902	419	505	94	89	36	43
2010–11	623	786	560	902	411	505	75	89	27	43
2011–12	417	571	431	629	256	248	94	89	20	43
2012–13	368	400	449	438	245	171	53	62	26	29
2013–14	382	400	435	438	248	171	60	62	28	29
2014–15	407	400	441	438	175	171	61	62	20	29
2015–16	344	400	386	438	172	171	52	62	7	29
2016–17	304	327	299	358	156	140	51	51	13	24

Table 2: [Continued]

Fish stock FMA (s)	BNS 10		Total	
	10			
	Landings	TACC	Landings	TACC
1981*	0		295	
1982*	0		484	
1983†	0		701	
1984†	0		1 386	
1985†	0		1 277	
1986†	0		1 536	
1986–87	7	10	1 411	1 350
1987–88	10	10	1 274	1 449
1988–89	10	10	1 420	1 566
1989–90	0	10	1 541	1 765
1990–91	#12	#10	1 781	1 831
1991–92	#40	#10	2 031	1 837
1992–93	#29	#10	2 335	2 016
1993–94	#3	#10	2 323	2 023
1994–95	0	10	2 285	2 161
1995–96	0	10	2 351	2 161
1996–97	#9	#10	2 846	2 480
1997–98	#30	#10	2 613	2 480
1998–99	#2	#10	2 775	2 480
1999–00	#0	#10	2 731	2 480
2000–01	#0	#10	2 721	2 480
2001–02	#0	#10	2 784	3 048
2002–03	0	10	3 002	3 058
2003–04	0	10	3 091	3 058
2004–05	0	10	3 012	3 233
2005–06	0	10	2 475	3 233

Table 2 [Continued]

Fish stock FMA (s)	BNS 10		Total	
	Landings	TACC	Landings	TACC
2006–07	0	10	2 425	3 233
2007–08	0	10	2 498	3 233
2008–09	0	10	2 046	2 335
2009–10	0	10	2 059	2 335
2010–11	0	10	1 696	2 335
2011–12	0	10	1 218	1 590
2012–13	0	10	1 142	1 110
2013–14	0	10	1 190	1 110
2014–15	0	10	1 104	1 110
2015–16	0	10	960	1 110
2016–17	0	10	823	910

* MAF data, † FSU data, # Includes exploratory catches in excess of the TAC, + An additional transitional 250 t of ACE was provided to Chatham Islands fishers, resulting in an effective commercial catch limit of 1 175 t in 2001–02 and 2002–03.

Bluenose landings prior to 1981 were poorly reported, with bluenose sometimes being recorded as bonita, or mixed with hapuku/bass/groper, and foreign licensed and charter catches in the 1970s included bluenose catches as warehou and butterfish. Landings before 1986–87 have been grouped by statistical areas which approximate the current QMAs.

TACCs were first established for bluenose upon introduction to the QMS in 1986–87, with TACCs for all bluenose stocks totalling 1350 t. From 1992 to 2009 all bluenose fishstocks were included, for at least some of the time, in Adaptive Management Programmes (AMPs). BNS 3 was the first stock to enter an AMP in October 1992, with a TACC increase from 175 t to 350 t. This was further increased within the AMP to 925 t in October 2001, plus an additional transitional 250 t of ACE provided to Chatham Islands fishers in 2001–02 and 2002–03 only. BNS 7 (TACC increase from 97 t to 150 t) and BNS 8 (TACC increase from 22 t to 100 t) entered AMPs in October 1994. BNS 1, the second largest bluenose fishery, entered an AMP in October 1996, with a TACC increase from 705 t to 1000 t. BNS 2, the largest bluenose fishery, was the most recent entry into an AMP in October 2004, with a TACC increase from 873 t to 1048 t. TACCs for all bluenose stocks were reduced on 1 October 2008: 786 (BNS 1), 902 (BNS 2), 505 (BNS 3), 89 (BNS 7) and 43 (BNS 8). AMP programmes were terminated on 30 September 2009.

Under a rebuild plan following the 2011 stock assessment, there have been further phased reductions to TACCs for bluenose stocks. On 1 October 2011, TACCs were reduced to: 571 (BNS 1), 629 (BNS 2), and 248 (BNS 3); BNS 7 and BNS 8 were not reduced at that time. On 1 October 2012, TACCs were further reduced for all bluenose stocks to: 400 (BNS 1), 438 (BNS 2), 171 (BNS 3), 62 (BNS 7) and 29 (BNS 8). The 2011 rebuild plan included a third phase of TACC reductions. For the 2016/17 fishing year, the Minister reduced the combined TACCs for bluenose stocks by 205 t as a further step towards ensuring the rebuild. He did not take stronger action as he wanted to provide the opportunity for a management procedure to be developed. As from October 2017, following the assessment being updated to include information up to the end of the 2015–16 year, the Minister noted that the stocks remained in a depleted state and he did not want to delay the rebuild any longer. Consequently, he reduced the TACCs for all BNS stocks further to ensure that BNS stocks rebuild towards the target at an appropriate rate consistent with the HSS guidelines

BLUENOSE (BNS)




Figure 1: Reported commercial landings and TACC for the five main BNS stocks. BNS 1 (Auckland East), BNS 2 (Central East), BNS 3 (South East Coast) [Continued on next page




Figure 1: [Continued] Reported commercial landings and TACC for the five main BNS stocks. BNS 7 (Challenger), BNS 8 (Central Egmont).

As a result of the TACC increases under AMPs, the combined total TACC for all bluenose stocks increased from an initial 1350 t in 1986–87 to 3233 t by 2004–05, before the reductions to 1110 t by 2012–13. Catch performance against the TACC has varied, with the combined TACC being under-caught by an average 9% (average landings 1504 t / year) over 1987–88 to 1990–91, over-caught by an average 11% (average landings 2501 t / year) over 1991–92 to 2000–01, and under-caught by an average 20% (average landings 2602 t / year) from 2004–05 to 2007–08.

1.2 Recreational fisheries

Bluenose is targeted by recreational fishers around deep offshore reefs. They are caught using line fishing methods, predominantly on rod and reel with some longline catch. The allowances within the TAC for each Fishstock are shown in Table 1.

1.2.1 Management controls

From 2012 onwards the catch limit for recreational fishers in all areas has been up to 5 bluenose per person per day as part of their multi-species (combined) individual daily bag limit.

1.2.2 Estimates of recreational harvest

There are two broad approaches to estimating recreational fisheries harvest: the use of onsite or access point methods where fishers are surveyed or counted at the point of fishing or access to their fishing activity; and, offsite methods where some form of post-event interview and/or diary are used to collect data from fishers.

The first estimates of recreational harvest for bluenose were calculated using an offsite approach, the offsite regional telephone and diary surveys. Estimates for 1996 came from a national telephone and

BLUENOSE (BNS)

diary survey (Bradford 1998). Another national telephone and diary survey was carried out in 2000 (Boyd & Reilly 2002) and a rolling replacement of diarists in 2001 (Boyd & Reilly 2004) allowed estimates for a further year (population scaling ratios and mean weights were not re-estimated in 2001). The annual recreational catch of BNS 1 was estimated from diary surveys to be 2 000 fish in 1993–94 (Teirney et al 1997), 5000 fish in 1996 (Bradford 1998) and 11 000 fish in 1999–00 (Boyd & Reilly 2004). The harvest estimates provided by these telephone diary surveys are no longer considered reliable.

A new national panel survey was developed, and implemented in the 2011–12 fishing year (Wynne-Jones et al. 2014). The panel survey used face-to-face interviews of a random sample of New Zealand households to recruit a panel of fishers and non-fishers for a full year. The panel members were contacted regularly about their fishing activities and catch information collected in standardised phone interviews. Recreational catch estimates from the national panel survey are given in Table 3. Note that the national panel survey estimates do not include recreational harvest taken under s111 general approvals on commercial vessels.

Table 3: Recreational harvest estimates for bluenose stocks (Wynne-Jones et al 2014). Mean fish weights were obtained from boat ramp surveys; for bluenose the value used was 4.473 kg (Hartill & Davey 2015).

Stock	Year	Method	Number of fish	Total weight (t)	CV
BNS 1	2011/12	Panel survey	6 287	28.15	0.40
BNS 2	2011/12	Panel survey	444	1.99	0.48
BNS 3	2011/12	Panel survey	461	2.05	0.92
BNS 7	2011/12	Panel survey	456	2.02	1.00
BNS 8	2011/12	Panel survey	137	0.61	1.03

The recreational surveys indicate that the recreational harvest of bluenose is relatively small in areas other than BNS 1. There are some locally important fisheries which will not have been adequately sampled by the national panel survey.

1.3 Customary non-commercial fishing

No quantitative information on the level of customary non-commercial take is available.

1.4 Illegal catch

No quantitative information on the level of illegal catch is available.

1.5 Other sources of mortality

There have been reports of depredation by *Orca* on bluenose caught by line fisheries.

2. BIOLOGY

Depth distribution

The depth distribution of bluenose extends from near-surface waters to about 1 200 m. Research trawl surveys record their main depth range as 250–750 m, with a peak at 300–400 m, and they regularly occur to about 800 m (Anderson et al 1998). Commercial catches recorded in logbook programmes implemented for some of the bluenose stocks under AMPs, and catch-effort data for these fisheries, confirm that bluenose catches range in depth from less than 100 m to about 1 000 m, depending on target species, but with a peak around 400 m for bluenose targeted fishing by any method.

The depth distribution of bluenose changes with size, with small juveniles known to occur at the surface under floating objects (Last et al 1993, Duffy et al 2000). Larger juveniles probably live in coastal and oceanic pelagic waters for one or two years. Fish 40–70 cm in length are caught between 200 m and 600 m, while larger fish, particularly those larger than 80 cm, are more often caught deeper than 600 m. A sequential move to deeper waters as bluenose grow has been confirmed by analysis of the stable radio-isotope ratios in otolith sections. Oxygen isotope ($\delta^{18}\text{O}$) ratios of bluenose otolith cores confirm residence of juvenile fish within surface waters. Changes in oxygen isotope ratios across otolith sections indicate changes in preferred mean depth with age of each fish (Horn et al 2008). That study hypothesised that the larger adults may be distributed below usually fished depths on underwater topographic features, but potentially available to fisheries as a result of regular vertical

feeding migrations. The largest adults appear to reside in 700–1000 m; i.e., deeper than most trawl or longline fishing for bluenose occurs. However, adult bluenose are also known to associate closely with underwater topographic features (hills and seamounts). Bluenose may undertake diurnal migrations into shallower depths to feed.

Age, growth and natural mortality

Recent ageing validation work by Horn et al (2008, 2010) substantially revised estimates of maximum age and size at maturity for bluenose which were previously considered to be moderately fast growing (Horn 1988). Radiocarbon (^{14}C) levels in core micro-samples from otoliths that had been aged using zone counts were compared with a bomb-radiocarbon reference curve which provided independent estimates of the age of the fish. Horn et al (2010) estimated a maximum age of 76 years, approximately twice the previous maximum age estimate. This maximum age is consistent with the maximum age of 85 years estimated for the closely related barrelfish (*Hyperoglyphe perciformis*) in the western North Atlantic, also determined, in part, using the bomb chronometer method (Filer & Sedberry 2008). Previous under-estimates of bluenose ages appears to have resulted from the incorrect interpretation of paired, fine ‘split rings’ as single growth zones, when they probably represent two separate growth zones.

Horn & Sutton (2011) recorded a maximum age of 71 years for BNS 1, and estimated natural mortality (M) to be in the range 0.09–0.15, based on 1% of the unfished population living to 30–50 years. Given the maximum recorded age, they commented that estimates of M less than 0.09 may be appropriate as bluenose live to at least 71 years and older fish may be poorly sampled by the line fishery. From the range of estimates resulting from recent ageing, the working group concluded that M for bluenose was unlikely to be over 0.1.

Instantaneous total mortality was estimated for five BNS 1 line fishery samples (Horn & Sutton 2011). The best estimates of Z ranged from 0.13 to 0.17, indicating that F was probably lower than M . This result was unexpected given recent strong declines in bluenose CPUE and the dramatic increase in targeting beginning in the mid-1980s. It was concluded that Z was underestimated, probably because the sampled fishing grounds did not hold closed populations, resulting in large or old fish being over-represented in the catch.

Maturity and reproduction

Biological parameters relevant to stock assessment are summarised in Table 4.

Table 4: Estimates of biological parameters for bluenose.

Fishstock			Estimate	Source
<u>1. Natural mortality (M)</u>				
BNS			0.09–0.15	Horn & Sutton (2011)
<u>2. Weight = a(length)^b (Weight in g, length in cm fork length).</u>				
BNS 2			<div>Both sexes</div> a = 0.00963 b = 3.173	Horn (1988a)
<u>3. Von Bertalanffy growth parameters</u>				
		<div>Females</div> <div>K t_0 L_{∞}</div>	<div>Males</div> <div>K t_0 L_{∞}</div>	
BNS 2	0.071	-0.5 92.5	0.125 -0.5 72.2	Horn et al (2010)
<u>3. Age at maturity (50%)</u>				
		<div>Females</div>	<div>Males</div>	
a ₅₀		17	15	Horn & Sutton (2011)

Little is known about the reproductive biology of bluenose. Maturity ogives derived from aged bluenose caught in BNS 1 from January to May indicated that ages at 50% maturity were about 15 and 17 years for males and females, respectively (Horn & Sutton 2011). Data from commercial logbook programmes implemented under AMPs indicate that bluenose sampled in QMAs 1, 3, 7 and 8 mature at between 60 cm and 65 cm. Analysis of gonad maturity stage proportions for bluenose sampled by commercial logbook programmes, primarily in BNS 1, 7 and 8, indicate that spawning probably peaks from February to April annually. No distinct spawning grounds have been identified for bluenose in New Zealand waters. The logbook programmes have sampled reproductively active

BLUENOSE (BNS)

fish around the North Island from East Cape to west of Cook Strait, and off the south west coast of the South Island. Observer data includes a small number of observations of spawning fish, but these extend from the southern half of FMA 10 to the Stewart-Snares shelf.

3. STOCKS AND AREAS

Stock boundaries are unknown, but similarity in trends in catch and CPUE across fisheries occurring in each of the five New Zealand BNS QMAs suggests the possibility that there may be a single BNS stock across all these areas, or of some close relationship between stocks in these QMAs. Tagging studies have shown that bluenose are capable of extensive migration, i.e., from the Wairarapa coast to Kaikoura, BoP, and North Cape (Horn 2003). There is a possibility that the long period of relatively stable CPUE observations in the face of increasing catches before the period of decline may be evidence of hyper-stability caused by the replenishment of adult stocks on specific areas or features. Increases in BNS targeting in some areas and increasing catches, could have exceeded the replenishment rate, causing the rapid and synchronous declines observed from about 2001–02 to 2011–12. Alternatively, there could be a simultaneous drop in recruitment due to coincident environmental factors. An environmental mechanism simultaneously affecting availability or catchability of BNS across all QMAs is considered to be less likely than the possibility of a single stock, or of correlated recruitment across sub-stocks in the various areas.

4. STOCK ASSESSMENT

The first fully quantitative stock assessment modelling for bluenose was carried out in 2011. Models were implemented in the general purpose Bayesian stock assessment program CASAL (Bull et al 2009). This assessment was updated in 2016, using standardised CPUE series and catch histories to 2014–15 (Bentley 2016). Methods for modelling CPUE were revised in 2014 (see Section 4.5)

4.1 Methods

Model structure

The 2011 assessment model (Cordue & Pomarède 2012) assumed a single New Zealand stock of bluenose, partitioned into two sexes, with 80 age groups (1–80 years with a plus group), and without maturity in the partition. The model has a single time-step, single area, two year-round fisheries (line and trawl), and mid-fishing-year spawning. The stock was assumed to be at B_0 in 1935. The maximum allowable exploitation rate in each fishery was set to 60%.

Data

The catch history in the model starts in 1936 when some bluenose were landed as groper or hapuku. The main uncertainty in the catch history is the foreign catch just prior to the implementation of the EEZ in 1978. Foreign vessels recorded bluenose catch within mixed-species groups, typically as part of a general warehou category. Catch data in the early 1980s were used to estimate the likely proportion of bluenose within a mixed warehou and bluenose group. Where possible, this was done on an area-specific basis and the proportions were applied to the pre-EEZ mixed-species catches. Due to the uncertainties in species attributions mentioned above, alternative bluenose proportions were used to construct three alternative catch histories: low, mid, and high (Figure 2, Table 5).

The catch histories for the line and trawl fisheries from 1989–90 to 2006–07 were derived from the bluenose characterisations conducted for the 2008 AMP review. From 2007–08 onwards, the total recorded catch was split between line and trawl fisheries in roughly the same proportion as the catches from the 2006–07 year. The 2009–10 catch was rounded down to provide the assumed total catch in 2010–11. Recreational and illegal catch were assumed to be zero.

Table 5: The three alternative catch (t) histories used in the BNS model runs. Trawl catch prior to 1970 was assumed to be zero.

	Line				Line				Trawl		
	Low	Mid	High		Low	Mid	High		Low	Mid	High
1936	0	75	150	1963	0	59	119				
1937	0	75	150	1964	0	66	133				
1938	0	75	150	1965	0	64	128				
1939	0	75	150	1966	0	61	123				
1940	0	56	112	1967	0	65	129				
1941	0	50	100	1968	0	57	113				
1942	0	50	100	1969	0	55	111				
1943	0	50	100	1970	0	70	140	1970	0	0	0
1944	0	50	100	1971	0	69	138	1971	0	0	0
1945	0	50	100	1972	0	59	118	1972	0	45	78
1946	0	69	138	1973	0	63	126	1973	0	42	72
1947	0	75	150	1974	0	69	137	1974	0	68	117
1948	0	81	162	1975	111	182	252	1975	0	116	204
1949	0	95	189	1976	618	692	767	1976	0	112	211
1950	0	89	177	1977	821	913	1004	1977	0	385	1505
1951	0	74	147	1978	1	81	161	1978	0	0	0
1952	0	71	142	1979	9	92	176	1979	0	0	0
1953	0	70	141	1980	15	98	180	1980	0	0	0
1954	0	69	137	1981	235	300	365	1981	0	0	0
1955	0	66	132	1982	469	511	554	1982	0	0	0
1956	0	69	138	1983	730	755	780	1983	0	0	0
1957	0	69	138	1984	951	956	962	1984	324	324	324
1958	0	75	149	1985	1013	1013	1013	1985	372	372	372
1959	0	68	137	1986	982	982	982	1986	605	605	605
1960	0	62	124	1987	744	744	744	1987	667	667	667
1961	0	60	121	1988	752	752	752	1988	522	522	522
1962	0	59	118	1989	797	797	797	1989	623	623	623

For all three catch histories

	Trawl	Line
1990	763	777
1991	577	1 192
1992	549	1 414
1993	733	1 573
1994	860	1 459
1995	904	1 382
1996	811	1 503
1997	1 060	1 765
1998	779	1 728
1999	904	1 871
2000	1 022	1 712
2001	1 082	1 638
2002	1 345	1 443
2003	1 331	1 671
2004	957	2 133
2005	1 114	1 900
2006	710	1 765
2007	424	2 001
2008	500	2 000
2009	300	1 746
2010	300	1 759
2011	300	1 700

Two CPUE indices were fitted as indices of abundance, one for line and one for trawl fisheries (Figure 3). CVs of 20% were assumed for each year. This assumption incorporates some process error as the estimated CVs for the CPUE indices are unrealistically low (as is typical for indices estimated using a GLM approach).

BLUENOSE (BNS)




Figure 2: The three alternative catch histories used in BNS model runs.




Figure 3: The line and trawl CPUE indices fitted in the 2016 BNS assessment model runs. Also presented is the CPUE series based on longline effort targeting groper (HPB).

Logbook and observer length samples were used to construct annual length frequencies for the line and trawl fisheries for each year when there were more than 500 fish measured (Line: 1993–2008; Trawl: 1995–2004). For each sample, the length frequency was scaled to the numbers of fish in the sampled catch. Catch-weighted samples were then combined with no further scaling or stratification.

Two age frequencies were fitted in each run: one from trawl caught fish on the Palliser Bank, for the single fishing-year 1985–86, and one for line caught fish in the BoP and East Northland, combined across areas for the fishing year 2000–01

Fixed and estimated parameters

In the final assessment runs, year-class strengths (YCSs) were assumed deterministic and only B_0 (uniform-log prior), the nuisance qs (for the two CPUE time series; uniform-log priors), the fishing selectivities (both double normal, uniform priors), and the CV of length at age (uniform prior) were estimated. Natural mortality (M) and steepness (h) were varied (see MPD runs below).

Fixed parameters were assigned the following values:

	Male	Female	Source
Length-weight (cm, g)			
a	0.00963	0.00963	Plenary report
b	3.173	3.173	
von Bertalanffy growth			
t_0	-0.5	-0.5	Horn et al 2010
L_∞	72.2	92.5	
k	0.125	0.071	
Maturity (logistic)			
a_{50}	15	17	Horn & Sutton 2010
$a_{95} - a_{50}$	5	10	Horn & Sutton 2010

Assessment runs

Initial assessment runs indicated that the assessment was sensitive to the assumed catch history, natural mortality, and stock-recruitment steepness. As a result the working group agreed to present results from a “grid” of MPD runs. The final set of 18 runs consisted of all combinations of:

- catch history: low, mid, high
- M : 0.06, 0.08, 0.10
- h : 0.75, 0.9

The M values cover what the working group considered a plausible range. The default assumption of $h = 0.75$ was adopted, and $h = 0.9$ was included as a sensitivity.

Iterative re-weighting was used to determine weights for the run with mid catch, $M = 0.08$ and $h = 0.75$. The CVs were unaltered from the initial assumption of 20%. These CVs and the sample-sizes, determined from the re-weighting, were fixed for all other runs. Convergence was checked for two runs (mid catch and mid M , with $h = 0.75$ and $h = 0.90$). An MCMC run was also conducted for mid catch and mid M with $h = 0.75$. This was to check that the MPD estimates were not substantially different from the medians of the posterior distributions for B_0 and stock status. As all runs had the same simple model structure, MCMCs were not conducted for other runs.

4.2 Results

The fishing selectivities for both trawl and line were estimated to be domed. However, the shapes of the fishing selectivities, especially for the line fishery, were confounded with M (Figure 4). The CV of length at age was estimated at 6% for all of the runs.

The fits to the CPUE indices were consistent with the assumed CVs of 20%. However, for both time series, a poor residual pattern was apparent, especially for the line CPUE (Figure 5). The line CPUE is flatter than the predicted values from 1990 to 2004, and then steeper than the predictions from 2005 to 2010.

The trawl and line fisheries showed different trends in exploitation rates, with the trawl fishery peaking from 2002 to 2005 and the line fishery increasing from 1980 to 2011 (Figure 6).

BLUENOSE (BNS)




Figure 4: Estimated fishing selectivities for the trawl and line fisheries for the final 18 MPD runs in the 2011 assessment. Each plot shows the results for six runs with the same value of M (which increases from 0.06 to 0.08 to 0.10 from left to right in the three plots).




Figure 5: The model fits to the line and trawl CPUE for the run with mid catch, mid M and $h = 0.75$. The fits for the other runs were almost identical.




Figure 6: Trends in fishing pressure (the maximum proportion of fish taken from any age class) for the line fishery for each of the assessment runs.

The differences between the biomass trajectories from the 18 assessment runs are driven by the value of M (Figures 7 and 8) with estimates of B_0 ranging from just over 30 000 t at an M of 0.1 to around 60 000 t with an M of 0.06.




Figure 7: Biomass trajectories (t) for the final set of 18 MPD runs.




Figure 8: Biomass trajectories (proportion of B_0) for the final set of 18 MPD runs.

Biomass trajectories, as a proportion of B_0 , all show a continuous decline in female spawner-biomass from the late 1980s to 2011, followed by a levelling off or slight increase to 2016, depending on model run (Figure 8). The runs presented are in two groups with regard to current stock status. The 6 runs with $M = 0.06$ are above 20% B_0 while the 12 runs with $M = 0.08$ or $M = 0.10$ are below 20% B_0 (Figure 8, Table 6). These results should not be interpreted as there being a 66% probability that the stock is below 20% B_0 . It is the range of the results that is important. The proportion of runs above or below 20% B_0 can be altered by including additional runs at different M values.

Table 6: Estimates of B_0 , B_{2015} and stock status (B_{2015}/B_0) for the final 18 runs. The range is given for the 6 runs at each value of M . B_0 and B_{2015} are mid-spawning season (after half the annual catch has been removed).

M	B_0 (000 t)	B_{2015} (000 t)	B_{2015}/B_0
0.06	60–62	14–17	0.24–0.27
0.08	42–44	7.2–8.3	0.17–0.19
0.10	33–34	5.9–6.1	0.17–0.18

4.3 Projections

Deterministic projections to 2050 were carried out as part of the 2011 and 2016 assessments, maintaining the 2009–10 ratio between catches from the line and trawl fisheries. For a stock below the soft limit of 20% B_0 , the time required for SSB to rebuild to 40% B_0 with no future catch is called T_{min} . Although the point estimates for some runs with low M are above 20% B_0 , the time required to rebuild to 40% B_0 was calculated for each run and is denoted as T_{min} . The estimates of T_{min} established using the 2011 assessment range from 10 to 13 years (Table 7). Catches at the level of the 2015–16 TACC were predicted (2016 assessment) to cause the stock to increase, but not nearly fast enough to attain the biomass target within the rebuild time frame (Figure 9). The maximum constant catches estimated by the 2016 assessment (and to be implemented in 2016–17) that allow a rebuild to 40% B_0 within twice the 2011 T_{min} (the maximum rebuilding time under the Harvest Strategy Standard) range from 600–840 t (Table 8).




Figure 9: Projected SSB at different catch levels from the 2016 run with alternative levels of M and h and catch histories. The short vertical lines around 40% B_0 mark $2011 + T_{min}$ and $2011 + 2 T_{min}$.

Table 7: The number of years before SSB reaches 40% B_0 when no future catch is taken (2011 Assessment). The duration, in a whole number of years, is defined as “ T_{min} ” and is shown for the six runs with the mid catch and combinations of M and h .

M	h	
	0.75	0.90
0.06	13	12
0.08	13	12
0.10	11	10

Table 8: The maximum constant catch (t) from 2016 that allows SSB to rebuild to at least 40% B_0 within twice T_{min} beginning in 2011 for the six runs with mid catch.

M	h	
	0.75	0.90
0.06	620	740
0.08	600	800
0.10	600	840

4.4 Other factors

This assessment relies on standardised catch per unit effort as an index of abundance. Members of the fishing industry have noted that bluenose fisheries have undergone a number of changes not all of which are adequately captured in the statutory catch and effort data. These include changes in quota holdings, company structures and vessel operators, and subtle shifts in fishing practice. The effect of increasing the number of hooks per line set and per day was investigated by identifying vessels that had changed their practice over time. The CPUE analysis was repeated without these vessels and the resulting standardised indices were very similar to those derived from the full dataset (Starr 2011).

Prior to 2008, CPUE was not considered to be a reliable indicator of abundance of bluenose. However, in 2008, close coincidence observed in declining trends in most trawl and line CPUE indices in recent years increased confidence in their value as indices of abundance. Standardised

BLUENOSE (BNS)

CPUE series, based on data from six fisheries spanning most major fisheries taking BNS in the NZ EEZ, declined an average of 64% over the period 2001–02 to 2006–07.

Catch at age data are limited, but suggest that the composition of catches can vary significantly on small spatial and temporal scales. The available catch-at-age data are insufficient to allow reasonable estimation of variation in year class strengths.

Information relating to bluenose stock structure is limited. In 2008, the AMP Working Group conducted full reviews of all bluenose Fishstocks which included separate CPUE abundance index standardisations for each Fishstock (Ministry of Fisheries 2008). The close coincidence between trends in the indices for all bluenose Fishstocks led the AMP Working Group to conclude that bluenose may constitute a single New Zealand-wide stock.

More complex spatial structuring of bluenose populations, such as the replenishment of the population on fished features from a wider stock pool, is also plausible and may imply a non-linear relationship between CPUE and abundance. However, preliminary modelling exploring a non-linear relationship between longline CPUE and abundance did not improve the fit to the CPUE indices.

4.5 Updated standardised CPUE indices

The approach to standardising CPUE indices for bluenose was reassessed in 2014 and the key indices were updated in 2016. For the line CPUE, effort and estimated catch data were summarised for every unique combination of vessel, date and statistical area. This reduced the higher resolution catch effort records (from LTCER and LCER forms) to lower resolution data compatible with records from the earlier CELR forms. The trawl CPUE used the higher resolution tow by tow data (from TCEPR and TCER forms) at their original resolution.

In 2014, separate CPUE indices were estimated for line fisheries targeting BNS, HPB and LIN as the high resolution data provides evidence of spatial separation in these fisheries, and they target differing depth ranges and achieve markedly different catch compositions. The BNS target line CPUE index was selected as the primary line index. The trawl CPUE index included both BT and MW trawling and BNS and BYX target tows.

The primary BLL.BNS standardisation used a Weibull error distribution and model selection retained fishing year, vessel, hooks and statistical area as explanatory variables. The influence of hook numbers was examined in detail to ensure that changes in reporting and fleet composition were dealt with appropriately in the standardisation.

Nine zones were defined, as groupings of statistical areas, which better separated the bluenose fisheries than the QMA boundaries. An amalgamated national line index was estimated by weighting the zone indices by the number of 0.1 degree cells they contained that accounted for 95% of the nationwide bluenose catch. These cells were used as a proxy for bluenose habitat.

Zone indices were calculated by fitting a zone \times year interaction (Figure 10). In general the individual zone indices show the same pattern as the overall index, with the exception of the southwest zone which has a much flatter index.




Figure 10: Zone-year indices for the line and trawl indices with the amalgamated national line index shown for reference. Zone-year combinations with less than 30 records are not shown.

The BNS target LL CPUE declined to a low point in 2011–12, increased markedly in 2012–13, but then dropped to a point in 2014–15 that remained above the 2011–12 nadir. The BNS trawl series (BNS and BYX target) had very similar overall trend to the LL series, but with general increase after the 2011–12 nadir (Figure 3). The LL BNS CPUE series based on HPB targeted effort had a similar trend to the BNS+BYX trawl series, with a gradual increase after the 2011–12 nadir, suggesting that BNS biomass had slowly increased since 2011–12; and that the spike in the BNS target LL series was probably disproportionate to abundance. All three series (BNS-BLL, HPB-BLL, and BNS/BYX-BT/MW) all have the same relative position in 2014–15.

Detailed analyses were undertaken of catch rates at the level of discrete spatial areas (“features”). No obvious, consistent changes in the distribution of catch/effort by feature since 2007–08 were apparent and there was general consistency among feature CPUE indices within a zone.

4.6 Management procedure evaluation

Four classes of management procedure were evaluated for the New Zealand bluenose fishery using the 2011 assessment as the basis for the operating model (Bentley & Middleton 2015). Evaluations were done using alternative operating model scenarios including re-estimating parameters using updated catch per unit effort (CPUE) series and different recruitment assumptions.

The MPE focussed on procedures that work to maintain a stock rebuild trajectory, and demonstrated that use of a management procedure to adjust catches provided for higher catches, for a given rebuild criterion, than maintaining a constant catch. After initial presentation of results to stakeholders, the “Trajectory Status Adjustment Restricted” (TSAR) class of management procedure (MP) became the focus for further evaluations and refinements. The TSAR class is based on a predefined CPUE trajectory with changes made to the total allowable commercial catch (TACC) when the smoothed CPUE index deviates from the defined trajectory.

BLUENOSE (BNS)

One of the performance statistics which MPs were evaluated against was the time taken to rebuild to 40% B_0 , using 25 years as an acceptance threshold (approximately two times T_{min}). Most of the TSAR instances evaluated failed to meet the 25 years to 40% B_0 rebuild criterion, but often by only a small margin.

5. STATUS OF THE STOCKS


Stock Structure Assumptions

The assessment presented here assumes that bluenose in New Zealand waters comprise a single biological stock.

BNS 1, BNS 2, BNS 3, BNS 7, BNS 8, BNS 10

Stock Status	
Year of Most Recent Assessment	2016
Assessment runs presented	Eighteen MPD runs exploring a plausible range of catch history, natural mortality rate, and stock-recruitment steepness
Reference Points	Target: Not formally established; assumed to be 40% B_0 (based on Harvest Strategy Standard Operational Guidelines, low productivity stock) Soft Limit: 20% B_0 (HSS default) Hard Limit: 10% B_0 (HSS default) Overfishing threshold: Not defined
Status in relation to Target	Unlikely (< 40%) to be at or above the default target.
Status in relation to Limits	About as Likely as Not (40–60%) to be below the Soft Limit Unlikely (< 40%) to be below the Hard Limit
Status in relation to Overfishing	-

Historical Stock Status Trajectory and Current Status



Spawning stock biomass trajectories (percentage of B_0) for the 2016 set of 18 MPD runs.

Fishery and Stock Trends	
Recent Trend in Biomass or Proxy	The MPD estimates of stock size in 2016 ranged from 17–27% B_0 . Biomass was estimated to have declined continuously from the 1980s to 2011 and then to have either levelled off or increased slightly. Biomass has been below the default 40% B_0 target since around 2000.
Recent Trend in Fishing Mortality or Proxy	Exploitation rates were estimated to have increased from 1980 as the stock declined. In 2011 exploitation rates in the trawl fishery were estimated to have declined since 2005, but remained high in the line fishery. Reduced TACCs since 2011

	have resulted in substantially reduced catches and a reduction in exploitation rates.	
Other Abundance Indices	A second BLL index based on bycatch of bluenose in the HPB LL fishery had a trend that was very similar to the Trawl index	
Trends in Other Relevant Indicator or Variables	-	
Stock Projections or Prognosis	Deterministic projections in 2011 with $M = 0.08$ and $h = 0.75$ predicted that stock abundance would decline to below the hard limit within the next 20 years (from 2010) under 2010 catch levels. The time to rebuild (T_{min}) to the assumed target (40% B_0) under zero catches ranges from 10 to 13 years, depending on model assumptions. Within the range of model runs explored, the maximum constant catch (EEZ wide) implemented in 2016 that would rebuild the stock to the target within twice T_{min} (beginning in 2011) was 600–620 t for $h = 0.75$ and 740–840 t for $h = 0.9$. A rebuilding plan to reduce catches and rebuild the stock to target levels within twice T_{min} was developed. Two stepped reductions in TACC were implemented and a third has been put on hold following a substantial increase in the standardised CPUE abundance indices. The 2016 assessment suggested that biomass had either levelled off after 2011 or increased slightly, and is projected to increase at current catches.	
Probability of Current Catch or TACC causing Biomass to remain below or to decline below Limits	Soft Limit: Unlikely (< 40%) Hard Limit: Very Unlikely (< 10%)	
Probability of Current Catch or TACC causing Overfishing to continue or to commence	Very Unlikely (< 10%)	
Assessment Methodology and Evaluation		
Assessment Type	Level 1: Full Quantitative Stock Assessment (2011)	
Assessment Method	Age-structured CASAL model with MPD estimation over a range of plausible catch histories, natural mortality rates and steepness.	
Assessment Dates	Latest assessment: 2016	Next assessment: 2021
Overall assessment quality rank	1 – High Quality	
Main data inputs (rank)	- CPUE indices derived from statutory catch and effort reporting - Length frequency data from sampling conducted under the Adaptive Management Programme, and from observer data - One age frequency distribution for each of the trawl and line fisheries	1 – High Quality 1 – High Quality 1 – High Quality
Data not used (rank)	N/A	
Changes to Model Structure and Assumptions	The longline CPUE index for the 2016 assessment is based only on BNS target fishing rather than BNS, HPB and LIN target sets (used in the 2011 assessment), and combined indices by zone weighted by a habitat proxy.	
Major Sources of Uncertainty	- Stock structure and spatial dynamics are uncertain. - The assessment assumes that CPUE indexes abundance. - Natural mortality is uncertain; the plausible range considered affects the estimate of current status, and is confounded with the estimated fishery selectivities. - Method specific selectivities are considered constant across areas. - Deterministic recruitment is assumed; variations in year class strengths are not estimated.	

BLUENOSE (BNS)

	- Catches are known and the catch history is complete.
Qualifying Comments	
Alternative plausible stock hypotheses have not been explored.	

Fishery Interactions
Bluenose is taken in conjunction with alfonso in target midwater trawl fisheries directed at the latter species and in target bluenose bottom trawl fisheries. These fisheries are frequently associated with undersea features. Bluenose is also taken by target bottom longline fisheries throughout the New Zealand EEZ. Other commercially important species taken when longlining for bluenose are ling, hapuku and bass. Incidental captures of seabirds occur in the bottom longline and setnet fisheries, including black petrel in FMA 1 and 2, that are ranked as at very high risk in the Seabird Risk Assessment. ¹ Interactions with other species are currently being characterised.

7. FOR FURTHER INFORMATION


- Anderson, O F; Bagley, N W; Hurst, R J; Francis, M P; Clark, M R; McMillan, P J (1998) Atlas of New Zealand fish and squid distributions from research bottom trawls. *NIWA Technical Report* 42. 303 p.
- Anon (2006) BNS 2 Adaptive Management Programme Report: 2004/05 fishing year. Document AMP-WG-06/17. (Unpublished manuscript available from Fisheries New Zealand, Wellington.)
- Blackwell, R G (1999) Catch sampling for size and age of bluenose (*Hyperoglyphe antarctica*) in BNS 2 during summer 1997–98. New Zealand Fisheries Assessment Research Document 1999/46. 15 p. (Unpublished document held in NIWA library.)
- Boyd, R O; Reilly, J L (2002) 1999/2000 national marine recreational fishing survey: harvest estimates. Draft New Zealand Fisheries Assessment Report. (Unpublished manuscript available from Fisheries New Zealand, Wellington.)
- Bradford, E (1998) Harvest estimates from the 1996 national recreational fishing surveys. New Zealand Fisheries Assessment Research Document 1998/16. 27 p. (Unpublished document held in NIWA library.)
- Challenger Finfish Management Company (CFMC) (2000) BNS 7 & 8 Adaptive Management Programme Proposal dated 7 May 2001. (Unpublished manuscript available from Fisheries New Zealand, Wellington.)
- Cordue, P.L.; Pomarède, M. (2012). A 2011 stock assessment of bluenose (*Hyperoglyphe antarctica*). *New Zealand Fisheries Assessment Report* 2012/06. 54p.
- Duffy, C A J; Stewart, A L; Yarrall, R (2000) First record of presettlement juvenile bluenose, *Hyperoglyphe antarctica*, from New Zealand. *New Zealand Journal of Marine and Freshwater Research* 34(2): 353–358.
- Filer, K R; Sedberry, G R (2008) Age, growth and reproduction of the barrellfish, *Hyperoglyphe perciformis* (Mitchill, 1818), in the western North Atlantic. *Journal of Fish Biology* 72: 861–882.
- Hartill, B; Davey, N (2015) Mean weight estimates for recreational fisheries in 2011–12. *New Zealand Fisheries Assessment Report* 2015/25. 37 p.
- Hoening, J M (1983) Empirical use of longevity data to estimate mortality rates. *Fisheries Bulletin* 82: 898–903.
- Horn, P L (1988a) Bluenose. New Zealand Fisheries Assessment Research Document 1988/9. 13 p. (Unpublished document held by NIWA library, Wellington.)
- Horn, P L (1988b) Age and growth of bluenose, *Hyperoglyphe antarctica* (Pisces: Stromateoidei) from the lower east coast, North Island, New Zealand. *New Zealand Journal of Marine and Freshwater Research* 22: 369–378.
- Horn, P L (2003) Stock structure of bluenose (*Hyperoglyphe antarctica*) off the north-east coast of New Zealand based on the results of a detachable hook tagging programme. *New Zealand Journal of Marine and Freshwater Research* 37: 623–631.
- Horn, P L; Massey, B R (1989) Biology and abundance of alfonso and bluenose off the lower east coast North Island, New Zealand. *New Zealand Fisheries Technical Report* 15. 32 p.
- Horn, P L; Neil, H L; Marriott, P M; Paul, L J; Francis, C (2008) Age validation for bluenose (*Hyperoglyphe antarctica*) using the bomb chronometer method of radiocarbon ageing, and comments on the inferred life history of this species. Final Research Report for Ministry of Fisheries Research Project BNS2005-01. 36 p. (Unpublished manuscript available from the Fisheries New Zealand Wellington.)
- Horn, P L; Neil, H L; Paul, L J; Marriott, P (2010) Age validation and growth of bluenose (*Hyperoglyphe antarctica*) using the bomb chronometer method of radiocarbon ageing. *Journal of Fish Biology* 77: 1552–1563.
- Horn, P L; Sutton, C P (2010) The spatial and temporal age structure of bluenose (*Hyperoglyphe antarctica*) commercial catches from Fishstock BNS 1. *New Zealand Fisheries Assessment Report* 2010/8. 22 p.
- Horn, P L; Sutton, C P (2011) The age structure of bluenose (*Hyperoglyphe antarctica*) commercial catches from the Palliser Bank (Fishstock BNS 2) in 1984–86, and estimates of mortality rates. *New Zealand Fisheries Assessment Report* 2011/30. 16 p.
- Jiang, W; Bentley, N (2008) BNS 2 Adaptive Management Programme Draft Report: 2006/07 Fishing Year. AMP-WG/2008/13, 91 p. (Unpublished manuscript available from Fisheries New Zealand, Wellington.)
- Langley, A D (1995) Analysis of commercial catch and effort data from the QMA 2 alfonso-blunose trawl fishery 1989–94. New Zealand Fisheries Assessment Research Document 1995/18. 12 p. (Unpublished document held by NIWA library.)
- Last, P; Bolch, C; Baelde, P (1993) Discovery of juvenile blue-eye. *Australian Fisheries* 52(8): 16–17.
- Mace, P M (1988) The relevance of *MSY* and other biological reference points to stock assessment in New Zealand. New Zealand Fisheries Assessment Research Document 88/6. (Unpublished document held by NIWA library.)
- Northern Inshore Finfish Management Company (NIFMC) (2001) BNS 1 Adaptive Management Programme Proposal dated 7 May 2001. (Unpublished manuscript available from Fisheries New Zealand, Wellington.)
- Paul, L J; Sparks, R J; Neil, H J; Horn, P L (2004) Maximum ages for bluenose (*Hyperoglyphe antarctica*) and rubyfish (*Plagiogeneion rubiginosum*) determined by the bomb chronometer method of radiocarbon ageing, and comments on the inferred life history of these species. Final Research Report for MFish Project INS2000/02. (Unpublished manuscript available from the Fisheries New Zealand Wellington.)

¹ The risk was defined as the ratio of the estimated annual number of fatalities of birds due to bycatch in fisheries to the Potential Biological Removal (PBR), which is an estimate of the number of seabirds that may be killed without causing the population to decline below half the carrying capacity. Richard and Abraham (2013).

- Richard, Y; Abraham, E R (2013) Risk of commercial fisheries to New Zealand seabird populations. *New Zealand Aquatic Environment and Biodiversity Report No. 109*. 58 p.
- Ryan, M; Stocker, M (1991) Biomass and yield estimates for bluenose in QMA 2 for the 1991/92 fishing year. New Zealand Fisheries Assessment Research Document 1991/8. 15 p. (Unpublished document held by NIWA library.)
- Southeast Finfish Management Company (SEFMC) (2001) BNS 3 Adaptive Management Programme Proposal dated 14 May 2001. (Unpublished manuscript available from Fisheries New Zealand, Wellington.)
- Sarr, P J (2011) Presentation to the 2011 Plenary Meeting: Problem: Increasing Number of Hooks Over Time in Bluenose Bottom Longline Fisheries. Plenary Meeting 2011/09. 23 p. Unpublished document available at the Fisheries New Zealand website: <http://cs.fish.govt.nz/forums/thread/5495.aspx>.
- Sarr, P J; Kendrick, T H (2011a) Report to the 2011 Plenary Meeting: Standardised CPUE for Total NZ Bluenose. Plenary Meeting 2011/12. 37 p. Unpublished document available at the Fisheries New Zealand website: <http://cs.fish.govt.nz/forums/thread/5777.aspx>.
- Sarr, P J; Kendrick, T H (2011b) Report To Northern Inshore Management Ltd: Review of the BNS 1 Fishery. NINSWG-2011-13. 71 p. Unpublished document available at the Fisheries New Zealand website: <http://cs.fish.govt.nz/forums/thread/5177.aspx>.
- Sarr, P J; Kendrick, T H (2011c) Report To Area 2 Inshore Finfish Management Company Ltd: Review of the BNS 2 Fishery. NINSWG-2011-14. 72 p. Unpublished document available at the Fisheries New Zealand website: <http://cs.fish.govt.nz/forums/thread/5220.aspx>.
- Sarr, P J; Kendrick, T H (2011d) Report To Southeast Finfish Management Company Ltd: Review of the BNS 3 Fishery. NINSWG-2011-15. 81 p. Unpublished document available at the Fisheries New Zealand website: <http://cs.fish.govt.nz/forums/thread/5192.aspx>.
- Sarr, P J; Kendrick, T H (2011e) Report To Challenger Fisheries Management Company Ltd: Review of the BNS 7 & BNS 8 Fisheries. NINSWG-2011-16. 59 p. Unpublished document available at the Fisheries New Zealand website: <http://cs.fish.govt.nz/forums/thread/5215.aspx>.
- Sarr, P J; Kendrick, T H; Bentley, N; Lydon, G J (2008a) 2008 Review of the BNS 1 adaptive management programme. AMP-WG-2008/11, 105 p. (Unpublished manuscript available from the Seafood New Zealand, Wellington.)
- Sarr, P J; Kendrick, T H; Bentley, N; Lydon, G J (2008b) 2008 Review of the BNS 3 adaptive management programme. AMP-WG-2008/06, 125 p. (Unpublished manuscript available from Seafood New Zealand, Wellington.)
- Sarr, P J; Kendrick, T H; Bentley, N; Lydon, G J (2008c) 2008 Review of the BNS 7 and BNS 8 adaptive management programme. AMP-WG-2008/09, 99 p. (Unpublished manuscript available from the Seafood New Zealand, Wellington.)
- Teirney, L D; Kilner, A R; Millar, R E; Bradford, E; Bell, J D (1997) Estimation of recreational catch from 1991/92 to 1993/94. New Zealand Fisheries Assessment Research Document 1997/15. 43 p. (Unpublished document held by NIWA library.)
- Vignaux, M (1997) CPUE analyses for Fishstocks in the Adaptive Management Programme. New Zealand Fisheries Assessment Research Document 1997/24. 68 p. (Unpublished document held by NIWA library, Wellington.)
- Wynne-Jones, J; Gray, A; Hill, L; Heinemann, A (2014) National Panel Survey Of Marine Recreational Fishers 2011–12: Harvest Estimates. *New Zealand Fisheries Assessment Report 2014/67*. 139 p.

BUTTERFISH (BUT)

(*Odax pullus*)
Marari



1. FISHERY SUMMARY

Butterfish was introduced into the QMS in 1 October 2002 with allowances, TACCs and TACs as follows (Table 1).

Table 1: Summary of recreational and customary non-commercial allowances, TACs, and TACCs.

Fishstock	Recreational Allowance	Customary non-commercial Allowance	TACC	Other Mortality	TAC
BUT 1	10	10	3	1	24
BUT 2	80	80	63	2	225
BUT 3	65	65	3	1	134
BUT 4	4	4	10	0	18
BUT 5	10	10	45	1	66
BUT 6	0	0	0	0	0
BUT 7	15	15	38	1	69
BUT 10	0	0	0	0	0
TOTAL	184	184	162	6	537

1.1 Commercial fisheries

Butterfish is targeted by setnets in shallow coastal waters, principally around kelp-beds. The main fishery is centred on Cook Strait, between Tasman Bay, Castlepoint, and Kaikoura. There is also a smaller fishery around Stewart Island. A minimum setnet mesh size of 108 mm and a minimum fish size of 35 cm apply to commercial and recreational fishers; additional regional netting restrictions may also apply.

Hector's dolphin setnet closure areas were introduced on 1 October 2008 as part of the implementation of a Hector's and Maui dolphin Threat Management Plan. On 18 March 2011 the Minister decided to provide an exemption to the setnet prohibition on the East Coast South Island to allow commercial fishers targeting butterfish to use setnets in a defined area at the top of the East Coast South Island.

In line with the acceptable risk of mortality associated with butterfish fishing by commercial fisheries at the top of the East Coast of the South Island, given the type of fishing gear they use and the size of the area and the numbers of Hector's dolphins, recreational fishers are also allowed to target butterfish by method of set net from 1 January–30 April (inclusive). Set netting can only be undertaken if fishers

BUTTERFISH (BUT)

stay with their nets at all times, the net is set no more than 200 m from the shore and it does not exceed 60 m in length.

Table 2: Reported domestic landings (t) and TACCs of butterfish by Fishstock from 2001–02 to 2016–17.

Fishstock FMA	BUT 1		BUT 2		BUT 3		BUT 4		BUT 5	
	1,8&9		2		3		4		5	
	Landings	TACC	Landings	TACC	Landings	TACC	Landings	TACC	Landings	TACC
2001–02	0.7	3	64	63	0.4	3	13	10	19	45
2002–03	2.0	3	58.2	63	2.8	3	4.0	10	34.6	45
2003–04	1.4	3	52.6	63	2.1	3	2.6	10	42.6	45
2004–05	1.5	3	62.9	63	2.4	3	5.3	10	35.4	45
2005–06	2.9	3	44.5	63	1.8	3	0.1	10	21.8	45
2006–07	2.4	3	55.5	63	1.8	3	0.1	10	30.1	45
2007–08	1.0	3	46.3	63	2.0	3	0	10	35.9	45
2008–09	2.1	3	55.5	63	0.6	3	0.6	10	36.9	45
2009–10	2.5	3	45.3	63	< 0.1	3	0.2	10	33.3	45
2010–11	3.1	3	42.4	63	0.1	3	0.2	10	47.0	45
2011–12	2.7	3	48.3	63	< 0.1	3	0.8	10	46.3	45
2012–13	2.1	3	53.8	63	0	3	0.1	10	34.5	45
2013–14	3.0	3	42.0	63	<1	3	<1	10	33.3	45
2014–15	2	3	36.3	63	<1	3	0	10	37.1	45
2015–16	1.4	3	38.1	63	<1	3	0	10	35.2	45
2016–17	2.8	3	44.4	63	<1	3	0	10	48.9	45

Fishstock FMA (s)	BUT 6		BUT 7		BUT 10		Total	
	6		7		10		Total	
	Landings	TACC	Landings	TACC	Landings	TACC	Landings	TACCs
2001–02	0	0	25	38	0	0	121	162
2002–03	0	0	28.5	38	0	0	130.1	162
2003–04	0	0	24.8	38	0	0	126.1	162
2004–05	0	0	24.5	38	0	0	132.0	162
2005–06	0	0	23.7	38	0	0	94.8	162
2006–07	0	0	26.9	38	0	0	116.8	162
2007–08	0	0	29.4	38	0	0	114.6	162
2008–09	0	0	26.3	38	0	0	122.0	162
2009–10	0	0	16.5	38	0	0	97.9	162
2010–11	0	0	23.3	38	0	0	116.2	162
2011–12	0	0	21.4	38	0	0	119.5	162
2012–13	0	0	19.9	38	0	0	110.4	162
2013–14	0	0	16.7	38	0	0	95.1	162
2014–15	0	0	21.8	38	0	0	97.1	162
2015–16	0	0	19.3	38	0	0	94.5	162
2016–17	0	0	18.2	38	0	0	114.3	162

Total reported landings from 1982–83 to 2000–01 ranged between 105 and 193 t. Butterfish was introduced into the QMS in 2002. Reported landings and TACCs are given in Table 2, while Figure 1 shows the historical landings and TACC values for the main BUT stocks.




Figure 1: Reported commercial landings and TACC for the four main BUT stocks: BUT 2 (Central East).




Figure 1 [Continued]: Reported commercial landings and TACC for the four main BUT stocks. From top, BUT 3 (South east coast), BUT 5 (Southland) and BUT 7 (Challenger).

1.2 Recreational fisheries

Butterfish is a popular recreational catch, and is taken mainly by setnet and spear. Recreational daily bag limits were set at 30 fish in 1986, but subsequently reduced to 20 for Northern and Central and Challenger (1995), and 15 for South (1993). Survey estimates indicate that the recreational catches appear to be of similar magnitude to those of the commercial fisheries in QMAs 1, 2, 5 and 7, and substantially higher in QMA 3 (Tables 3a and 3b).

BUTTERFISH (BUT)

Table 3a: Estimated recreational harvest of butterfish by QMA and survey.

QMA	Survey	Number caught	Survey harvest (t)	Fishstock harvest (t)
				1991–92
QMA 7	South	6 000	10	
QMA 7	South	4 000	5	15
QMA 3	South	36 000	65	65
QMA 5	South	8 000	10	10
				1993–93
QMA 2	Central	61 000	80	80
				1993–94
QMA 1 + 9	North	9 000	10	10
TOTAL		124 000		180

*Surveys were in different years: South 1991–92; Central 1992–93; and North 1993–94 (Teirney et al 1997). Many of these estimates have high CVs, and the estimate of total harvest is a guide only because of the different survey years. Line-caught ‘butterfish’ in QMA 3 and QMA 5 are excluded because of apparent species misidentification; these survey totals should be slightly higher.

Table 3b: Estimated number and weight of butterfish harvested by recreational fishers by Fishstock and survey. Surveys were carried out nationally in 1999–2000 (Boyd & Reilly 2002).

Fishstock	Survey	Number	CV%	Survey harvest (t)
BUT 1	National	1 000	71	< 1–3
BUT 2	National	23 000	39	16–36
BUT 3	National	45 000	47	27–76
BUT 5	National	17 000	42	11–27
BUT 7	National	18 000	41	12–29
BUT 8	National	1 000	100	0–2

The harvest estimates provided by telephone-diary surveys between 1993 and 2001 are no longer considered reliable for various reasons. A Recreational Technical Working Group concluded that these harvest estimates should be used only with the following qualifications: a) they may be very inaccurate; b) the 1996 and earlier surveys contain a methodological error; and c) the 2000 and 2001 estimates are implausibly high for many important fisheries. In response to these problems and the cost and scale challenges associated with onsite methods, a National Panel Survey was conducted for the first time throughout the 2011–12 fishing year. The panel survey used face-to-face interviews of a random sample of 30 390 New Zealand households to recruit a panel of fishers and non-fishers for a full year. The panel members were contacted regularly about their fishing activities and harvest information collected in standardised phone interviews. Harvest estimates for butterfish are given in Table 4 (from Wynne-Jones et al 2014 and Hartill & Davey 2015).

Table 4: Recreational harvest estimates for butterfish stocks (Wynne-Jones et al 2014). Mean fish weights were obtained from boat ramp surveys; for butterfish the values used were 1.068 kg for BUT 1 and 1.123 kg elsewhere (Hartill & Davey 2015).

Stock	Year	Method	Number of fish	Total weight (t)	CV
BUT 1	2011/12	Panel survey	27 488	29.4	0.64
BUT 2	2011/12	Panel survey	13 892	15.6	0.33
BUT 3	2011/12	Panel survey	13 637	15.3	0.42
BUT 5	2011/12	Panel survey	188	0.2	0.74
BUT 7	2011/12	Panel survey	14 625	16.4	0.94
BUT total	2011/12	Panel survey	69 831	76.9	0.34

A repeat of the National Panel Survey is being conducted over the 2017–18 October fishing year. Results are expected in early 2019.

1.3 Customary non-commercial fisheries

There is no quantitative information on the current level of customary non-commercial catch.

1.4 Illegal catch

Because this is a localised small-scale fishery, some sales from fishers directly to retailers may have gone unreported, but no quantitative estimate of this are available.

1.5 Other sources of mortality

There is no quantitative information on other sources of mortality. In the past butterfish has been used as

rock lobster bait and not reported.

2. BIOLOGY

Butterfish are endemic to New Zealand, and occur from North Cape to the Snares Islands. The species is also reported from the Chatham, Bounty and Antipodes Islands. Butterfish are more common from Cook Strait southwards. They inhabit rocky coastlines, and are commonly found among seaweed beds in moderately turbulent water. Their main depth range is 0–20 m. They occur shallower (to 10 m) in the north than in Cook Strait (to 20 m) and in southern waters they can be found as deep as 40 m.

Adult butterfish average 45–55 cm (FL) in length. Their maximum size is approximately 70 cm. Length/weight data are not available for whole fish, but as an interim measure a length/gutted weight relationship is given in Table 5.

Butterfish are almost exclusively herbivorous, feeding on several of the larger seaweeds. The diet of butterfish varies regionally and is largely determined by the species composition of the local seaweed beds. Feeding activity is greatest early in the day, and the tidal state controls the accessibility of intertidal seaweeds; fish were found to feed more actively in summer than winter (Trip 2009).

Fish were aged using sectioned sagittal otoliths, validated using daily growth (Trip 2009). Growth varies with latitude due to temperature difference, and local ecological factors such as diet and fish density.

Trip (2009) found that size and age differ significantly with latitude. Environmental temperature is the primary driver underlying the difference in life histories across latitudes, and affects growth rate, size-at-age and longevity. Butterfish living in colder temperatures (higher latitudes) grow slower, live longer, attain a greater average size and delay the onset of maturity (Trip 2009). Butterfish in Hauraki Gulf (BUT 1) reach 70% of their mean asymptotic size by the age of two, and have reached 90% of their maximum size by age 4. In the southern areas butterfish grow slower and reach a maximum size at about 75 % of their life span. The maximum age ranged from 11 years in the north (Hauraki Gulf) to 19 years in the south (Stewart Island) (Trip 2009). There are no significant differences in growth rates or mean adult body size between sexes, yet with the exception of the Hauraki Gulf, the oldest and largest fish (FL) sampled in all areas were females (Trip 2009).

Table 5: Estimates of biological parameters for butterfish.

Estimates of biological parameters for butternish.				Estimate	Source	
Fishstock						
<u>1. Natural mortality (<i>M</i>)</u>						
Cook Strait				0.30–0.45	Paul et al (2000)	
<u>2. Weight = a(length)^b (Weight in g, length in cm fork length).</u>						
	Females		Males	Juvenile		
	a	b	a	b		
Cook Strait	67.699	1 947.8	67.034	1 885.9	21.205	Ritchie (1969)
Hauraki Gulf					362.28	
Stewart Is.						
Linear regression, b = constant. Weight is gutted weight.						
<u>3. von Bertalanffy growth parameters</u>						
				Both sexes		
	<i>K</i>	<i>t</i> ₀	<i>L</i> _∞			
Cook Strait	0.23	-1.7	51.8			Paul et al (2000)
Hauraki Gulf	0.517	-0.23	457.36			Trip (2009)

Butterfish start life as female, some, but not all, undergo sex change where an estimated 50% of mature females develop into males. The size at sex change ranges between 37 to 45 cm FL. The length at which sex change occurs does not seem to differ between geographical areas, but age-at-sex change varies geographically. The mean age-at-sex change was found to be significantly lower in warmer latitudes, 2.5 yrs at the Hauraki Gulf, in comparison to 7 years old at Stewart Island. At D'Urville Island, in-between the two, fish changed sex at 5 years old (Trip 2009).

BUTTERFISH (BUT)

In the warm waters of the north females mature early and of the samples collected in the Hauraki Gulf 95% of females are sexually mature by two years old (29.7 cm FL). Females sampled at Stewart Island show delayed maturity with only 50% mature at an average age of four (25.2 cm FL) (Trip 2009).

The depth distribution of butterfish differs by size and sex. Juveniles (less than 30 cm) occur in the shallow weed beds (less than 15 m) and (outside the breeding season) males occur in deeper waters than females. Consequently, sex ratios vary with locality, but females often outnumber males.

In the North the spawning season occurs between July and November, with a peak in August. The spawning season extends from July to March in Cook Strait, peaking in September and October. In southern New Zealand the spawning season appears to be shorter (August to January, peaking in October–January).

3. STOCKS AND AREAS

There is no clear information on whether biologically distinct stocks occur, although there is some evidence of regional variation in meristic characters which suggests some separation of populations. The time larval butterfish spend in the plankton before settling out into the adult habitats as postlarvae is relatively short, a factor that may cause a high level of stock separation around coastal New Zealand. The only information on movement relates to feeding behaviour involving small-scale movements within seaweed beds. There is no information on movement along the coastline within a weed-bed habitat, or potentially longer migration between such habitats separated by open coast. However, the latter seems unlikely on any substantial scale, and as a result butterfish populations are probably quite localised. Butterfish populations at offshore islands (Chatham, Antipodes, Bounties, and Snares), have not been studied but may be distinct from the mainland population(s) simply because of their isolation.

4. STOCK ASSESSMENT

A yield per recruit analysis was undertaken in 1997 (Paul et al 2000). This report derived new estimates of growth and natural mortality from the Cook Strait which were incorporated into this analysis. Stock status was not determined by this analysis.

4.1 Estimates of fishery parameters and abundance

No information is available.

4.2 Biomass estimates

No information is available.

4.3 Yield estimates and projections

The method $MCY = cY_{av}$ (Method 4) was evaluated. However, this method was rejected due to a lack of reliable information on changes in fishing effort and/or mortality over the history of the fishery. MCY for butterfish cannot be determined.

CAY cannot be determined.

4.4 Other yield estimates and stock assessment results

A study of setnet mesh selectivity in relation to the current legal minimum fish size showed that 108 mm mesh retained few undersized fish (immature). This provides a level of protection to butterfish stocks and their recruitment. A yield per recruit analysis showed that a modest yield increase could be obtained by using a smaller mesh and taking younger (2–3 year old) fish. However, this theoretical gain would be counter-balanced by the capture of relatively more juveniles and young females, and almost certainly a higher bycatch of other reef fishes. Butterfish populations are susceptible to localised depletion.

5. STATUS OF THE STOCKS

No estimates of current and reference biomass are available. It is not known whether recent catch levels will allow the stock to move towards B_{MSY} .

Reported landings and TACCs are summarised in Table 6.

Table 6: Summary of reported landings (t) and TACCs by QMA for the most recent fishing year.

Fishstock		FMA	2016–17 Actual TACC	2016–17 Reported landings
BUT 1	Auckland (East)(West), Central (West)	1,8&9	3	2.8
BUT 2	Central (East)	2	63	44.4
BUT 3	South-east coast	3	3	<1
BUT 4	Chatham	4	10	0
BUT 5	Southland	5	45	48.9
BUT 6	Sub-Antarctic	6	0	0
BUT 7	Challenger	7	38	18.2
BUT 10	Kermadec	10	0	0
TOTAL			162	114.3


6. FOR FURTHER INFORMATION

- Boyd, R O; Reilly, J L (2002) 1999/2000 national marine recreational fishing survey: harvest estimates. Draft New Zealand Fisheries Assessment Report. (Unpublished report held by Fisheries New Zealand, Wellington.)
- Choat, J H; Clements, K D (1993) Daily feeding rates in herbivorous labroid fishes. *Marine Biology* 117(2): 205–211.
- Clements, K D; Choat, J H (1993) Influence of season, ontogeny and tide on the diet of the temperate marine herbivorous fish *Odax pullus* (Odacidae). *Marine Biology* 117(2): 213–220.
- Dunn, A; Paul, L J (2000) Estimates of butterfish (*Odax pullus*) setnet selectivity. *New Zealand Fisheries Assessment Report 2000/6*. 22 p.
- Graham, D H (1953) *A Treasury of New Zealand Fishes*. AH. & AW. Reed, Wellington. 424 p. (Revised 1956, reprinted 1974.)
- Hartill, B; Davey, N (2015) Mean weight estimates for recreational fisheries in 2011–12. *New Zealand Fisheries Assessment Report 2015/25*.
- Hickford, M J H; Schiel, D R (1995) Catch vs. count: Effects of gill-netting on reef fish populations in southern New Zealand. *Journal of Experimental Marine Biology and Ecology* 188(2): 215–232.
- Paul, L J (1997) A summary of biology and commercial landings, and a stock assessment of butterfish, *Odax pullus* (Forster in Bloch and Schneider 1801) (Labroidae: Odacidae). New Zealand Fisheries Assessment Research Document 1997/23. 25 p. (Unpublished document held by NIWA library, Wellington.)
- Paul, L J; Ó Maolagáin, C; Francis, M P; Dunn, A; Francis, R I C C (2000) Age, growth, mortality, and yield per recruit for butterfish (*Odax pullus*) in Cook Strait, New Zealand. *New Zealand Fisheries Assessment Report 2000/6*. 30 p.
- Ritchie, L D (1969) Aspects of the Biology of the Butterfish *Coriododax pullus* (Forster). Unpublished M.Sc. Thesis, Victoria University of Wellington. 145 p.
- Teirney, L D; Kilner, A R; Millar, R E; Bradford, E; Bell, J D (1997) Estimation of recreational catch from 1991/92 to 1993/94 New Zealand Fisheries Assessment Research Document 1997/15. 43 p. (Unpublished document held by NIWA library, Wellington.)
- Trip, E D L (2009) Latitudinal variation in the demography and life history of a temperate marine herbivorous fish *Odax pullus* (labridae). (Unpublished Ph.D. thesis lodged in the School of Biological Sciences, University of Auckland, Auckland, New Zealand)
- Wynne-Jones, J; Gray, A; Hill, L; Heinemann, A (2014) National Panel Survey of Marine Recreational Fishers 2011–12: Harvest Estimates. *New Zealand Fisheries Assessment Report 2014/67*.

COCKLES (COC)

(Austrovenus stutchburyi)

Tuangi



1. INTRODUCTION

Cockles are important shellfish both commercially and for non-commercial fishers.

Since 1992, Fisheries New Zealand or its predecessors has commissioned biomass surveys for cockles and pipi in the northern North Island on beaches where there is known recreational and customary fishing pressure. The objective of the surveys is to determine the distribution, abundance and size frequency of cockles and pipi on selected beaches in the Auckland Fisheries Management Area. Over the years, a total of 34 beaches have been monitored. On average, 12 beaches are sampled each year. The last survey was conducted in 2017 (see Berkenbush & Neubauer, 2017). All of the 2016–17 survey sites contained notable cockle populations, and data from the field sampling were sufficient to provide cockle population estimates with relatively low uncertainty, i.e., with a CV of less than 20%. Most (10) sites had relatively high population densities, where estimates exceeded 400 individuals per m², with two sites with particularly high density estimates (1461 individuals per m² in Ngunguru Estuary in Northland and 1125 individuals per m² in Whangamata Harbour on Coromandel Peninsula). In contrast, the lowest density estimate was at Grahams Beach, where cockles occurred at an estimated 64 individuals per m². Density estimates of large cockles highlighted their rareness in the northern populations, with values well below the density estimates for the total cockle population at each site.

1.1 Commercial fisheries

Commercial picking of cockles, *Austrovenus stutchburyi*, is carried out on Snake Bank, Whangarei Harbour (FMA 1), Papanui and Waitati Inlets, Otago (FMA 3) and Pakawau Beach, Ferry Point and Tapu Bay in Tasman and Golden Bays (FMA 7). Cockles have also been commercially harvested from Otago Harbour since August 2009 under a special permit. Cockles were introduced into the QMS on 1 October 2002. The fishing year runs from 1 October until September 30 and catches are measured in greenweight for all stocks. There is no minimum legal size for commercial or non-commercial fishers for cockles in any stock. Cockles are managed under Schedule 6 of the Fisheries Act for all stocks listed in Table 1, which allows cockles to be returned to where they were taken as soon as practicable after the cockle is taken as long as the cockle is likely to survive.

COCKLES (COC)

For assessment purposes, individual reports on the largest fisheries have been produced separately:

1. Snake Bank, Whangarei Harbour, in COC 1A.
2. Papanui Inlet, Waitati Inlet, and Otago Harbour, Otago Peninsula in COC 3.
3. Tasman and Golden Bays in COC 7A.

The landings, by stock, of these cockle fisheries are dominated by catch from COC 3 (Figure 1). Landings from COC 3 are relatively stable since 2002–03; by contrast landings from COC 1A and COC 7A have generally declined over that time period.

Information on cockles that applies to all stocks is included below rather than being repeated in the reports for each fishery.

New Zealand operates a mandatory shellfish quality assurance programme for all bivalve shellfish commercial growing or harvesting areas for human consumption. Shellfish caught outside this programme can only be sold for bait. This programme is based on international best practice and managed by Food Safety New Zealand in cooperation with the District Health Board Public Health Units and the shellfish industry¹ and is summarised below. Before any area can be used to grow or harvest bivalve shellfish, public health officials survey both the water catchment area to identify any potential pollution issues and microbiologically sampling water and shellfish over at least a 12-month period, so all seasonal influences are explored. This information is evaluated and, if suitable, the area classified and listed by Food Safety New Zealand for harvest. There is then a requirement for regular monitoring of the water and shellfish flesh to verify levels of microbiological and chemical contaminants. Management measures stemming from this testing include closure after rainfall, to deal with microbiological contamination from runoff. Natural marine biotoxins can also cause health risks, therefore testing for these also occur at regular intervals. If toxins are detected above the permissible level the harvest areas are closed until the levels fall below the permissible level. Products are also traceable so that the source and time of harvest can always be identified in case of contamination.

Table 1: TACC, Recreational, customary allowances and TAC (t) for all cockle stocks.

Code	Description	TACC	Recreational allowance	Customary allowance	TAC
COC 1A	Whangarei Harbour	346	25	25	396
COC 1B	East Northland	0	22	22	44
COC 1C	Hauraki Gulf and Bay of Plenty	5	32	32	69
COC 2	Central	0	2	2	4
COC 3	Otago	1 470	10	10	1 490
COC 3B	Part South East Coast	1	27	27	55
COC 4	South East (Chatham Rise)	0	1	1	2
COC 5	Southland and Sub-Antarctic	2	2	2	6
COC 7A	Nelson Bays	1 390	85	25	1 500
COC 7B	Marlborough	0	5	5	10
COC 7C	Part Challenger	0	3	3	6
COC 8	Central (Egmont)	0	1	1	2
COC 9	Auckland (West)	0	6	6	12

¹For full details of this programme, refer to the Animal Products (Regulated Control Scheme-Bivalve Molluscan Shellfish) Regulations 2006 and the Animal Products (Specifications for Bivalve Molluscan Shellfish) Notice 2006 (both referred to as the BMSRCS), at: <http://www.nzfsa.govt.nz/industry/sectors/seafood/bms/page-01.htm>




Figure 1: Commercial landings and the sum total (black line) of the three main commercial COC stocks throughout time. Note that this figure does not show data prior to entry into the QMS.

2. BIOLOGY

The cockle, *Austrovenus stutchburyi*, formerly known as *Chione stutchburyi*, is a shallow-burrowing suspension feeder of the family Veneridae. It is found in soft mud to fine sand on protected beaches and enclosed shores around the North and South Islands, Stewart Island, the Chatham Islands and the Auckland Islands (Morton & Miller 1973, Spencer et al 2002). Suspension feeders such as *A. stutchburyi* tend to be more abundant in sediments with a larger grain size. Cockles have been shown to be most abundant in sediments of below 12 percent mud in two separate studies (Thrush et al 2003, Anderson 2008). They are also common in eelgrass (e.g., *Zostera* sp.), which often co-occurs with sand flats.

Cockles are found from the lowest high water neap tide mark to the lowest part of the shore. Larcombe (1971) suggested that the upper limit is found where submergence is only 3.5 hours per day. *A. stutchburyi* is often a dominant species and densities as high as 4500 per m² have been reported in some areas. In Pauatahanui Inlet the cockle biomass was estimated at 80% (5000 t) of the total intertidal biomass in 1976 (Richardson et al 1979). Calculations based on laboratory measurements of filtration rates suggested that cockles over 35 mm shell length were capable of filtering 1.1×10^6 m³ of water or enough to filter all the water in Papanui Inlet every two tidal cycles (Pawson 2004).

Sexes are separate and the sex ratio is usually close to 1:1. Size at maturity has been estimated at about 18 mm shell length (Larcombe 1971). Spawning extends over spring and summer, and fertilisation is followed by a planktonic larval stage lasting about three weeks. Significant depression of larval settlement has been recorded for areas of otherwise suitable substrate from which all live cockles have been removed. This suggests the presence of some conditioning factor.

Work on Snake Bank also showed moderate differences among years in the level of recruitment of juveniles to the population. The variability of recruitment was estimated as $\sigma_R = 0.41$ using all available data (1983–1996) but as $\sigma_R = 0.31$ using data only from those years since the fishery has been considered to be fully developed (1991–96). Given the variability of most shellfish populations and the shortness of the time series, this is probably an underestimate of the real variability of recruitment in the Snake Bank population.

Small cockles grow faster than large cockles, but overall, maximum growth occurs on the first of January, and a period of no growth occurs at the beginning of July (Tuck & Williams 2012). Growth is slower in the higher tidal ranges and in high density beds. Significant increases in growth rates have been observed for individuals remaining in areas that have been ‘thinned out’ by simulated harvesting. Tagging work at Pakawau beach also highlighted the variability in growth that can occur within a beach (Osborne 2010).

Growth parameters and length weight relationships are listed in Table 2 (Stewart 2008, Williams et al 2009, Osborne 2010). However, considerable variability in growth has been seen in all three QMAs

over time. At Snake bank (1A) growth to 30 mm has been estimated as taking between 2 and 5 years in separate studies (Martin 1984, Cryer 1997). Additional tagging work on Snake Bank from 2001 to 2010 showed that on average, cockles reach maturity (18 mm; Larcombe 1971) in their second year of growth, and recruit to harvestable size (about 28 mm SL) in about 3 to 4 years, although these results showed great variability in growth rate (tabulated in table 8, Tuck & Williams 2012). At Pakawau beach (7A) K has varied between 0.36 and 0.41 and L_{∞} between 47 and 49mm (Osborne 1992, 1999). The work of Breen et al (1999) in Papanui and Waitati Inlets, Purakanui and Otago Harbour showed no significant growth after one year and modes in the length frequency distributions did not shift when measured over four sampling periods within a year. They concluded that it was unlikely that average growth is really as slow as the results indicated, but there may be high inter-annual variability in growth.

Quite extensive movements of juveniles have been documented, but individuals over 25 mm shell length remain largely sessile, moving only in response to disturbance.

Given that cockles recruit to the spawning biomass at about 18 mm shell length, but do not recruit to commercial or non-commercial fisheries until closer to 30 mm shell length, there is some protection for the stock against egg overfishing, especially as the Snake Bank and Papanui and Waitati Inlet stocks are probably not isolated as far as recruitment of juveniles is concerned. However, this generality should be treated with some caution, given that some population of adults seems to be required to stimulate settlement of spat.

Natural mortality arises from a number of sources. Birds are a major predator of cockles (up to about 23 mm shell length). Other predators include crabs and whelks. Cockles are also killed after being smothered by sediments shifted during storms or strong tides. A mass mortality that killed an estimated 56–63% of all cockles and 80–84% of cockles over 30 mm in shell length (Fisheries New Zealand unpublished data) has been reported from sites within the Whangateau harbour (north of Auckland). This mortality was attributed to a potential weakening of cockles due to heat stress then mortality from a coccidian parasite and a mycobacterium². Sediments, both suspended and deposited, both impact upon cockle fitness or survival, with terrestrial sediments having greater effects than marine sediments (Gibbs & Hewitt 2004). Increasing suspended sediment concentrations have induced increased physiological stress, decreased reproductive status and decreased juvenile growth rates (Nicholls et al 2003, Gibbs & Hewitt 2004). Sediment deposition has also been shown to negatively impact upon densities of cockles (Lohrer et al 2004). The sum of these effects is seen in the distribution of cockles which decline in abundance across a number of sites with increasing mud content in the sediments, either above zero or 11% mud content, depending upon the study (Thrush et al 2003, Anderson 2008).

Experimental work on Snake Bank led to estimates of absolute mortality of 17–30% per annum, instantaneous natural mortality (M) of 0.19–0.35, with a midpoint of $M = 0.28$. The estimated mortality rates for cockles of over 30 mm shell length were slightly greater at 19–37% per annum, (M of 0.21–0.46 with a midpoint of 0.33). This higher estimate was caused by relatively high mortality rates for cockles of over 35 mm shell length and, as these are now uncommon in the population, $M = 0.30$ (range 0.20–0.40) has been assumed for yield calculations across all three stocks (Table 2). Tagging (both notch and individual numbered tags) has been ongoing on Mair Bank from 2001 to 2009 and the last recoveries occurred in 2010 (Tuck & Williams 2012). Annualised mortality estimates (M) (averaged over 3, 6 and 9 month recoveries) were 0.356 and 0.465 from studies in 2008 and 2009.

² <http://www.biosecurity.govt.nz/media/21-08-09/cockle-death-whangateau-estuary>

Table 2: Biological parameters used for cockle assessments for different stocks. SL = shell length, within area 7A, P = Pakawau, FP = Ferry Point, TBR = Tapu Bay/Riwaka.

	1A	3	7A
1. Natural mortality (<i>M</i>)	0.3	0.3	0.3
2. Weight (grams)	= $a(\text{shell length})^b$ = $a(\text{shell length}) + b$ = $a(\text{shell length})^b$		
a	0.00014	0.7211	P = 0.000018, FP = 0.0002, TBR = 0.00015
b	3.29	11.55	P = 3.78, FP = 3.153, TBR = 3.249
3. von Bertalanffy growth parameters	Not used instead growth = $a(\ln(\text{age in years})) + b$		
<i>K</i>	0.26	0.311	a = 11.452
<i>L_∞</i> (mm)	35	40.95	b = 16.425
SL at recruitment to the fishery (mm)	28	28	30

3. STOCKS AND AREAS


Little is known of the stock boundaries of cockles. Given the planktonic larval phase, many populations may receive spat fall from other nearby populations and may, in turn, provide spat for these other areas. In the absence of more detailed knowledge, each commercial fishery area is managed as a discrete population.

4. FOR FURTHER INFORMATION

- Anderson, M J (2008) Animal-sediment relationships re-visited: Characterising species' distributions along an environmental gradient using canonical analysis and quantile regression splines. *Journal of Experimental Marine Biology and Ecology* 366(1–2): 16–27.
- Berkenbush, K; Neubauer, P (2017) Intertidal shellfish monitoring in the northern North Island region, 2016–17. *New Zealand Fisheries Assessment Report 2017/51*. 107 p.
- Breen, P A; Carbines, G C; Kendrick, T H (1999) Stock assessment of cockles in Papanui and Waitati Inlets, Otago Harbour, and Purakanui, Otago. Final Report for the Ministry of Fisheries research project COC9701 dated July 1999. (Unpublished report held by Fisheries New Zealand, Wellington.)
- Cryer, M (1997) Assessment of cockles on Snake Bank, Whangarei Harbour, for 1996. New Zealand Fisheries Assessment Research Document 97/2. 29 p. (Unpublished report held in NIWA library, Wellington.)
- Ellis, J I (2003) Habitat change in estuaries: predicting broad-scale responses of intertidal macrofauna to sediment mud content. *Marine Ecology-Progress Series* 263: 101–112.
- Gibbs, M; Hewitt, J E (2004) Effects of sedimentation on macrofaunal communities: a synthesis of research studies for ARC. *Auckland Regional Council Technical Publication* 264.
- Irwin, C (2004) The impacts of harvesting and the sustainability of a New Zealand Littleneck Clam (*Austrovenus stutchburyi*) Fishery in Papanui and Waitati Inlets, New Zealand. Marine Science. Dunedin, University of Otago. PhD.
- Larcombe, M (1971) The ecology, population dynamics, and energetics of some soft shore molluscs. . Department of Zoology, University of Auckland. PhD. 250 p.
- Lohrer, A; Thrush, S; Hewitt, J E; Berkenbusch, K; Ahrens, M; Cummings, V J (2004) Terrestrially derived sediment: response of marine macrobenthic communities to thin terrigenous deposits. *Marine Ecology Progress Series* 273: 121–138.
- McKinnon, J (1996) Studies of the age, growth and shell increment patterns in the New Zealand cockle (*Austrovenus stutchburyi*). Unpublished Msc thesis, University of Otago, Dunedin, New Zealand.
- Manly, B F J; Akroyd, J M; Walshe, K A R (2002) Two-phase stratified random surveys on multiple populations at multiple locations. *New Zealand Journal of Marine and Freshwater Research* 36: 581–591.
- Martin, N D (1984) *Chione stutchburyi* population responses to exploitation. Unpublished MSc Thesis, University of Auckland, Auckland.
- Morton, J; Miller, M (1973) *The New Zealand Sea Shore*. Collins, Auckland. 653 p.
- Nicholls, P; Hewitt, J; Halliday, J (2003) Effects of suspended sediment concentrations on suspension and deposit feeding marine macrofauna. *Auckland Regional Council Technical Publication* 211. 40 p.
- Osborne, T A (1992) Biomass survey and stock assessment of the New Zealand littleneck clam (*Chione stutchburyi*) on Pakawau Beach, Golden Bay. Unpublished report prepared for Westhaven Shellfish Co. Ltd. 27 p. (please request from Westhaven).
- Osborne, T A (1999) 1999 biomass survey and stock assessment of the New Zealand littleneck clam (*Chione stutchburyi*) on Pakawau Beach, Golden Bay. Unpublished report prepared by Osborne Research Co. Ltd. for Westhaven Shellfish Co. Ltd. 15 p.
- Osborne, T A (2010) Biomass survey and stock assessment of cockles (*Austrovenus stutchburyi*) in area COC 7A: Tapu Bay, Ferry Point, and Pakawau. *New Zealand Fisheries Assessment Report 2010/44*.
- Pawson, M (2004) The cockle *Austrovenus stutchburyi* and chlorophyll depletion in a southern New Zealand Inlet. Marine Science. Dunedin, University of Otago. MSc. 113 p.
- Richardson, J R; Alridge, A E; Main, W, deL (1979) Distribution of the New Zealand cockle, *Chione stutchburyi* at Pauatahanui Inlet. NZOI Oceanographic Field Report 14. 10 p.
- Spencer, H G; Willan, R C; Marshall, B A; Murray, T J (2002) Checklist of the Recent Mollusca described from the New Zealand Exclusive Economic Zone. <http://toroa.otago.ac.nz/pubs/spencer/Molluscs/index.html>
- Stewart, B (2008) Stock Assessment of Clams (*Austrovenus stutchburyi*) in Waitati Inlet, Otago, 2007. Report Prepared by Ryder Consulting Ltd. for Southern Clams Ltd. (please request from Southern Clams).
- Thrush, S F; Hewitt, J E; Norkko, A; Nicholls, P E; Funnell, G A; Ellis, J I (2003) Habitat change in estuaries: predicting broad-scale responses of intertidal macrofauna to sediment mud content. *Marine Ecology-Progress Series* 263: 101–112.
- Tuck, I; Williams, J (2012) Cockle growth at Snake Bank, Whangarei Harbour, 2001–10, Fisheries Research Report, Unpublished Report for Ministry for Primary Industries, 40 p.
- Williams, J R; Smith, M D; Mackay, G (2009) Biomass survey and stock assessment of cockles (*Austrovenus stutchburyi*) on Snake Bank, Whangarei Harbour, 2009. *New Zealand Fisheries Assessment Report 2009/29*. 22 p.

COCKLES (COC 1A) Snake Bank (Whangarei Harbour)

(*Austrovenus stutchburyi*)
Tuangi

**1. FISHERY SUMMARY**

COC 1A was introduced to the QMS in October 2002 with a TAC of 400 t, comprising a TACC of 346 t, customary and recreational allowances of 25 t each, and an allowance of 4 t for other fishing related mortality. These limits have remained unchanged since.

1.1 Commercial fisheries

Snake Bank is not the only cockle bed in Whangarei Harbour, but it is the only bed open for commercial fishing. Commercial fishers are restricted to hand gathering, but they routinely use simple implements such as “hand sorters” to separate cockles of desirable size from smaller animals and silt. There are several other cockle beds in the harbour, some on the mainland and some on other sandbanks, notably MacDonald Bank. Fishing on these other beds should be exclusively non-commercial.

Commercial picking in Whangarei Harbour began in the early 1980s and is now undertaken year round, with no particular seasonality. Catch statistics (Table 1) are unreliable before 1986, although it is thought that over 150 t of Snake Bank cockles were exported in 1982. There was probably some under reporting of landings before 1986, and this may have continued since. Effort and catch information for this fishery has not been adequately reported by all permit holders in the past, and there are problems interpreting the information that is available. Landed weights reported on CELRs only summed to between 52 and 91% of weights reported on LFRRs during the years 1989–90 to 1992–93. CPUE data are available but have not yet been analysed for this fishery.

Before entry of this stock to the QMS there were eight permit holders, each allowed a maximum of 200 kg (greenweight) per day by hand-gathering. If all permit holders took their quota every day a maximum of 584 t could be taken in a 365 day year. Reported landings of less than 130 t before 1988–89 rose to 537 t in 1991–92 (about 92% of the theoretical maximum). Landings for the 1992–93 fishing year were much reduced (about 316 t) following an extended closure for biotoxin contamination. Landings averaged 462 t between 1993–94 and 2000–01. Landings have decreased substantially since COC 1A entered the QMS (average of 108 t), and no landings have occurred since 2011–12, this closure (in November 2012) was due to low biomass.

COCKLES (COC 1A)

Table 1: Reported commercial landings and catch limits (t greenweight) of cockles from Snake Bank since 1986–87 (from QMR/MHR records)*. Before COC 1A entered the QMS, the fishery was restricted by daily catch limits which summed to 584 t in a 365 day year, but there was no explicit annual restriction. A TACC of 346 t was established in October 2002 when COC 1A entered the QMS.

Fishing year	Landings (t)	Limit (t)	Fishing year	Landings (t)	Limit (t)
1986–87	114	584	2001–02	405	584
1987–88	128	584	2002–03	237	346
1988–89	255	584	2003–04	218	346
1989–90	426	584	2004–05	151	346
1990–91	396	584	2005–06	137	346
1991–92	537	584	2006–07	111	346
1992–93	316	584	2007–08	151	346
1993–94	**566	584	2008–09	88	346
1994–95	501	584	2009–10	93	346
1995–96	495	584	2010–11	64	346
1996–97	457	584	2011–12	43	346
1997–98	439	584	2012–13	0	346
1998–99	472	584	2013–14	0	346
1999–00	505	584	2014–15	0	346
2000–01	423	584	2015–16	0	346
			2016–17	0	346

*Before COC 1A entered the QMS, the fishery was restricted by daily catch limits which summed to 584 t in a 365 day year, but there was no explicit annual restriction. A TACC of 346 t was established in October 2002 when COC 1A entered the QMS. ** The figure of 566 t for 1993–94 may be unreliable.

The low catch in recent years may partly reflect reduced effort on the bank because of temporary fishery closures during incidents of sewage and stormwater overflows which adversely affected harbour water quality. The fishery was closed for these reasons for 101, 96, 167 and 96 days for the 2006–7, 2007–8, 2008–9 and 2009–10 fishing years, respectively¹. Figure 1 shows the recent landings and TACC values of COC 1A.

The mean length of the commercial harvest is about 29.5 mm and cockles smaller than 25 mm are less attractive to both commercial and non-commercial fishers.




Figure 1: Reported commercial landings and TACC for COC 1A (Whangarei Harbour).

1.2 Recreational fisheries

The recreational fishery is harvested entirely by hand digging, and large cockles (30 mm shell length or greater) are preferred. No recreational harvest estimates specific to the Snake Bank fishery are available. A regional telephone and diary survey in 1993–94, and national recreational diary surveys in 1996, 1999–2000, and 2000–01 estimated the numbers of cockles harvested in QMA 1 to be 0.57–2.4 million (Table 2). It is not clear to what extent these estimates include customary take. No mean harvest weight for cockles was available, but an assumed mean weight of 25 g (as for cockles 30 mm SL or more from the 1992 Snake Bank survey) led to a QMA 1 recreational harvest of 14–59 t (Table 2).

The harvest estimates provided by telephone-diary surveys between 1993 and 2001 are no longer considered reliable for various reasons. A Recreational Technical Working Group concluded that these

¹ Statistics supplied by New Zealand Food Safety Authority in Whangarei.

harvest estimates should be used only with the following qualifications: a) they may be very inaccurate; b) the 1996 and earlier surveys contain a methodological error; and c) the 2000 and 2001 estimates are implausibly high for many important fisheries. In response to these problems and the cost and scale challenges associated with onsite methods, a National Panel Survey was conducted for the first time throughout the 2011–12 fishing year. The panel survey used face-to-face interviews of a random sample of 30 390 New Zealand households to recruit a panel of fishers and non-fishers for a full year. The panel members were contacted regularly about their fishing activities and harvest information collected in standardised phone interviews. Harvest estimates for cockles in FMA 1 are given in Table 2 (from Wynne-Jones et al 2014).

Table 2: Estimated numbers of cockles harvested by recreational fishers in QMA 1, and the corresponding harvest tonnage based on an assumed mean weight of 25 g. Figures were extracted from telephone-diary surveys in 1993–94, 1996, 1999–00, and 2000–01 and the National Panel Survey in 2011–12.

Year	QMA 1 harvest (number of cockles)	CV (%)	QMA 1 harvest (t)	Source
1993–94	2 140 000	18	55	Bradford (1997)
1996	569 000	18	14	Bradford (1998)
1999–00	2 357 000	24	59	Boyd & Reilly (2002)
2000–01	2 327 000	27	58	Boyd et al (2004)
2011–12	299 765	68	7	Wynne-Jones et al (2014)

1.3 Customary fisheries

In common with many other intertidal shellfish, cockles are very important to Maori as a traditional food. The MFish customary catch database contained no records of Maori customary harvest of cockles from COC 1A. Patuharakeke gazetted their rohe moana which covers the southern shoreline of the Whangarei harbour in 2009. Reporting of customary permits is now required. However, a full understanding of Maori customary take will not occur until such time as all iwi operate under the Fisheries (Kaimoana Customary Fishing) Regulations 1998.

1.4 Illegal catch

Anecdotal evidence suggests that there was a significant illegal catch from Snake Bank in the 1990s, with some fishers greatly exceeding their catch limits. Commercial landings, therefore, may have been under-reported. There is also good evidence that illegal commercial gathering has occurred on MacDonald Bank on a reasonable scale in the past, which could have resulted in some over-reporting of catch from Snake Bank in some years. However, no quantitative information on the level of illegal catch is available.

1.5 Other sources of mortality

No quantitative information on the level of other sources of mortality is available. It has been suggested that some methods of harvesting such as brooms, rakes and “hand sorters” cause some mortality, particularly of small cockles, but this proposition has not been tested.

2. BIOLOGY

Biological parameters used in this assessment are presented in the general cockle section.

3. STOCKS AND AREAS

This is covered in the general cockle section.

4. STOCK ASSESSMENT

Stock assessment for Snake Bank cockles has been conducted periodically using absolute biomass surveys, yield per recruit (YPR), and spawning stock biomass per recruit (SSBPR) modelling. The stock

assessments were used to estimate *CAY* and *MCY*. A length-based stock assessment model was developed for cockles but was not successful.

4.1 Estimates of fishery parameters and abundance

Estimated and reference fishing mortality rates, estimates of total mortality and exploitation rate are available for Snake Bank (Table 3, Figure 2). Exploitation rate in 2012 and 2013 was 0% and had generally had a downward trend since 1991 (70%) with the exception of a large peak around 2001 (93%). Exploitation rate is likely to be overestimated in the calculation below as the size of cockles commercially harvested is believed to have decreased from over 30 mm to over 28 mm shell length over time.

Table 3: Estimates of fishery parameters.

Population and years	Estimate	Source
<u>1. Estimated Fishing Mortality (F_{est}, recruited size classes only)</u>		
Snake Bank, 1991–92	1.55	Cryer (1997)
Snake Bank, 1992–93	0.62	Cryer (1997)
Snake Bank, 1995–96	0.50	Cryer (1997)
Snake Bank, 1991–96	0.89	Cryer (1997)
<u>2. Reference Fishing Mortality (F_{ref}, recruited size classes only)</u>		
Snake Bank, $F_{0.1}$	0.41	Cryer (1997)
Snake Bank, F_{max}	0.62	Cryer (1997)
Snake Bank, $F_{50\%}$	4.52	Cryer (1997)
<u>3. Total Instantaneous Mortality (Z, all size classes)</u>		
Snake Bank, 1992–93	0.46	Cryer & Holdsworth (1993)
<u>4. Exploitation rate percentage (≥ 30 mm shell length)</u>		
Year*	%	
1991	71	
1992	41	
1995	34	
1996	57	
1998	54	
1999	38	
2000	74	
2001	93	
2002	51	
2003	21	
2004	28	
2005	14	
2006	14	
2007	11	
2008	8	
2009	11	
2012	0	
2013	0	

* Exploitation rate is only given in years when biomass surveys were completed and catch reporting was considered reliable (apart from in 2012 and 2013 where no catch was reported, therefore exploitation rate percentage must be zero).

4.2 Biomass estimates

Biomass estimates for the Snake Bank cockle population from 1982–96 were made using grid surveys. Surveys done from 1998 used a stratified random approach (Table 4, Figure 3). The data given here differ from those in reports before 1997 because the assumptions made when estimating biomass have changed. The surveys conducted in 1985 and 1991 did not cover the whole area of the bank, and results from these surveys have been corrected in the table by assuming that the cockle population occupied the same area of the bank in these years as it did in 1982 (the first and largest survey). It has been further assumed for the estimation of variance for the grid based surveys that samples have been taken at random from the bank, although variance estimators not requiring this assumption gave very similar results in 1995 and 1996. The post 1997 surveys also incorporated a large area of low density cockles not included in previous surveys, although this adds only a small tonnage of biomass to the total figure. In 1998 and 2000, biomass surveys were undertaken at MacDonald Bank using a stratified random approach (Table 5). Cryer et al (2003) reported biomass estimates for several locations in Whangarei Harbour in 2002, including a new MacDonald Bank stratum (Table 5). Northland Regional Council completed a survey in 2014 but only reported total biomass (Griffiths & Eyre 2014), this is included as it gives a recent indication of biomass in the absence of commercial fishing.




Figure 2: Exploitation rate (≥ 30 mm shell length).

Table 4: Estimates of biomass (t) of cockles on Snake Bank for surveys (n , number of stations) between 1982 and 2015. Biomass estimates for the ≥ 18 mm shell length component and those marked with an asterisk (*) were made using length frequency distributions and length-weight regressions, the other size fractions were generated by direct weighing of samples. Two alternative estimates are presented for 1988 because the survey was abandoned part-way through, “a” assuming the distribution of biomass in 1988 was the same as in 1991, and “b” assuming the distribution in 1988 was the same as in 1985. The 2001 result comes from the second of two surveys, the first having produced unacceptably imprecise results. The 2007 and 2008 results differ slightly from those reported previously because they were estimated using an analytical approach more consistent with that used in other years. The column “% $B_{recruited}$ ” compares the biomass in the ≥ 30 mm SL to the defined B_0 for that size (22 340 t in 1982).

Year	n	Total		≥18 mm SL		≥ 30 mm SL		≥ 35 mm SL		% $B_{recruited}$
		Biomass	CV	Biomass	CV	Biomass	CV	Biomass	CV	
1982	199	2 556	-	-	-	*2 340	-	1 825	~ 0.10	100
1983	187	2 509	-	2460	0.06	*2 188	-	1 700	~ 0.10	94
1985	136	2 009	0.08	1360	0.07	1 662	0.08	1 174	~ 0.10	71
1988 a	53	-	-	-	-	1 140	> 0.15	-	-	-
1988 b	53	-	-	-	-	744	> 0.15	-	-	-
1991	158	1 447	0.09	1069	0.08	761	0.10	197	0.12	33
1992	191	1 642	0.08	1355	0.07	780	0.08	172	0.11	33
1995	181	2 480	0.07	2380	0.07	1 478	0.07	317	0.12	63
1996	193	1 755	0.07	-	-	796	0.08	157	0.11	34
1998	53	2 401	0.18	-	-	880	0.17	114	0.20	38
1999	47	3 486	0.12	2645	0.11	1 321	0.14	194	0.32	56
2000	50	1 906	0.23	2609	0.18	570	0.25	89	0.32	24
2001	51	1 405	0.17	1382	0.17	435	0.17	40	0.29	19
2002	53	1 618	0.14	-	-	466	0.19	44	0.29	20
2003	60	2 597	0.11	2385	0.31	1 030	0.12	121	0.14	44
2004	65	1 910	0.15	1096	0.14	546	0.14	59	0.22	23
2005	57	2 592	0.18	2035	0.15	967	0.20	111	0.20	41
2006	57	2 412	0.13	2039	0.13	792	0.13	103	0.20	34
2007	73	2 883	0.13	2681	0.13	1 434	0.15	329	0.42	61
2008	70	2 510	0.10	-	-	1 165	0.11	193	0.43	50
2009	75	1 686	0.15	-	-	815	0.13	88	0.19	35
2014	63	1 794	0.14	-	-	-	-	-	-	-

Virgin biomass, B_0 , is assumed to be equal to the estimated biomass of cockles above a certain shell length in 1982. For example, if a length at recruitment of 30 mm or more was used then a biomass of 2340 t resulted. This biomass was estimated using length frequency distributions, a length weight regression, and a direct estimate of the biomass of cockles ≥ 35 mm shell length in 1982 (1 825 t).

Between the start of the commercial fishery in 1982 and the survey in 1992, there was a consistent decline in the biomass of large cockles (≥ 30 mm shell length) on Snake Bank. The biomass of these large individuals declined to 33% of its virgin level in 1991. A decrease in the proportion and biomass of large, old individuals can be expected with the development of a commercial fishery. The biomass of mature cockles has fluctuated since then without trend between 63 and 19% of virgin levels. The recruited biomass is likely to be underestimated in the calculation below as the size of cockles

commercially harvested is believed to have decreased from over 30 mm to over 28 mm shell length over time. There was no survey that has allowed calculation of percent B_0 since 2009.




Figure 3: Recruited biomass (≥ 30 mm shell length) over time as a percentage of B_0 in relation to the hard and soft limits.

Table 5: Biomass estimates (t) and approximate CVs by shell length size classes for cockles on MacDonald Bank. n = the number of samples in the survey.

Year	n	Total		< 30 mm SL		≥ 30 mm SL		≥ 35 mm SL	
		Biomass	CV	Biomass	CV	Biomass	CV	Biomass	CV
1998	33	6 939	0.19	5 261	0.18	1 678	0.31	128	0.41
2000	30	6 037	0.28	4 899	0.29	1 137	0.30	34	0.37
2002	24	2 548	0.12	2 010	0.14	538	0.36	61	0.46

4.3 Yield estimates and projections

A range of sizes are taken commercially, selectivity seems to vary between years and MCY estimates are sensitive to the assumed size at recruitment to the fishery (Table 6). These are presented over time for two different shellfish lengths at recruitment into the fishery (when available), 30 mm the historic size at recruitment, and 28 mm the more recently accepted size at recruitment (Table 7). All of these estimates include commercial and all non-commercial catch.

Table 6: Sensitivity of biomass and CAY estimates to shell length at recruitment (L_{RECR}) for Snake Bank cockles

L_{reer} (mm)	Rationale	B_{av} (1991–2009) (t)	B_{curr} (2009) (t)	M	$F_{0.1}$	MCY (t)	CAY (t)
25	Smallest in catch	1 877	1 596	0.3	0.34	385	401
28	Fisher selectivity	1 409	1 265	0.3	0.38	289	349
30	Historical assumption	890	815	0.3	0.41	182	239
35	Largest cockles	145	88	0.3	1.00	30	49

As fishing is conducted year round on Snake Bank, the Baranov catch equation is appropriate (Method 1, see Plenary introduction). This approach assumes that, between the start of the fishing year and when the biomass survey is started, productivity and catch cancel each other. The estimate includes non-commercial catch.

A range of sizes are taken commercially, selectivity seems to vary between years and CAY estimates are sensitive to the assumed size at recruitment to the fishery (Table 6). The level of risk to the stock by harvesting the population at the estimated CAY value cannot be determined.

4.4 Other yield estimates and stock assessment results

$F_{0.1}$ was estimated using a yield per recruit (YPR) model using quarterly (rather than the more usual annual) increments and critical sizes (rather than ages) for recruitment to the spawning stock and to the fishery. The following input information was used: growth rate parameters from a MULTIFAN analysis of 1991–96 length frequencies; an estimate of $M = 0.30$ (range 0.20–0.40) from a tagging study in 1984;

length weight data from 1992, 1995 and 1996 combined; size at maturity of 18 mm; and size at recruitment of 30 mm from an analysis of fisher selectivity. For the base case analysis, $F_{0.1} = 0.41$. Estimates were neither sensitive to the length weight regression used, nor to the value of M chosen ($F_{0.1} = 0.38$ – 0.45 for $M = 0.20$ – 0.40), but were more sensitive to the assumed length at recruitment ($F_{0.1} = 0.34$ for $L_{recr} = 25$ mm).

Table 7: *MCY* and *CAY* estimates (t) for different shell lengths at recruitment (L_{RECR}). *MCY* is calculated using the equation for developing fisheries prior to 1995 and developed fisheries after 1995. A value for 2010 is not shown as no survey was completed in COC 1A in 2010. Year labels as given in Table 4.

Year	<i>MCY</i> ≥ 28 mm SL	<i>MCY</i> ≥ 30mm SL	<i>CAY</i> ≥ 28 mm SL	<i>CAY</i> ≥ 30mm SL
1982		240		687
1983		240		642
1985		240		488
1988 a		240		335
1988 b		240		218
1991		240		223
1992		240		229
1995		206		434
1996		196		234
1998		192		258
1999		206		388
2000		193		167
2001		180		128
2002		171		137
2003	269	175	255	302
2004		169		160
2005	238	171	389	284
2006	254	171	329	233
2007	243	179	516	421
2008	293	183	584	342
2009	268	182	349	239

4.5 Other factors

Biomass and yield estimates will differ for different sizes of recruitment. Maori and recreational fishers prefer cockles of 30 mm shell length and greater whereas commercial fishers currently prefer cockles of 25 mm and greater. Therefore, yield has been estimated for sizes of recruitment between 25 and 30 mm. As cockles become sexually mature at around 18 mm, using a size of recruitment between 25 mm and 30 mm should provide some protection against egg overfishing under most circumstances. However, using the smaller size of recruitment to estimate yield will confer a greater risk of overfishing.

As the Snake Bank cockle population may receive spat from spawnings in other parts of Whangarei Harbour, it may not be realistic to assume that the Snake Bank stock is discrete and that reduced egg production (as a result of heavy fishing mortality on medium and large sized individuals) would necessarily lead to recruitment overfishing. Spawning stock biomass per recruit (SSBPR) analysis suggests that $F_{50\%} > F_{max} > F_{0.1}$ ($F_{50\%}$ is that fishing mortality which would lead to egg production from the population at equilibrium being half of egg production from the virgin stock), except where the size at recruitment is reduced to 25 mm. Substantial reduction of egg production is therefore unlikely if fishing mortality is restrained to within $F_{0.1}$ or F_{max} , and the fishery concentrates on cockles over 30 mm in length.

However, it has been demonstrated for this bank that recruitment of juvenile cockles can be reduced by the removal of a large proportion of adult cockles from a given area of substrate. Conversely, there did not seem to be heavy recruitment to the population during the years when adult biomass was close to virgin (1982–85). This would suggest that there is some optimal level of adult biomass to facilitate recruitment, although its value is not known. It would appear prudent, therefore, to exercise some caution in reducing the biomass of adult cockles. If adult biomass is driven too low, then recruitment overfishing of this population could still occur despite high levels of egg production. In addition, sporadic recruitment of juveniles will probably lead to a fluctuating biomass, suggesting that a *CAY* approach may be more appropriate than a constant catch approach.

A length-based stock assessment model developed in 2000 allowed for more of the natural variability of the system to be incorporated in the stock assessment. This first model did not adequately capture the detail of cockle dynamics. Further work in 2002 (McKenzie et al 2003) did not resolve all of these

COCKLES (COC 1A)

problems and substantial conflict remained in the model. Additional information on growth and the length frequency of cockles taken by the fishery was collected in 2003 and 2004 and updated in the model. Several additions and enhancements to the model were also made in an attempt to resolve the above-mentioned conflict (Cryer et al 2004, Watson et al 2005). As a result, the model showed an improved fit to the observed data. However, there still remained some conflict, primarily relating to annual variability in the growth increment data, in which only two years of observations were available (2002 and 2004). This was thought to be due to the existence of annual variability in recruitment, and possibly mortality, which are presently not explicitly modelled. Watson et al (2005) therefore concluded that no further development of the model should be undertaken for three to five years, and that resources be concentrated more on data collection, and in particular, growth and recruitment data. Consequently, a tag-recapture experiment was started in March 2005, and additional large samples of cockles have been notch-tagged and released annually from 2005 to 2010. Tagged individuals were recovered and measured on a quarterly basis, and preliminary results suggested there may be strong seasonal variability in growth.

Although the Shellfish Working Group considered that the development of a length-based stock assessment model would be of considerable benefit to the stock assessment, the problems with the model were such that the current approach used to estimate yield for this fishery that had been agreed to by the Shellfish Fishery Assessment Working Group since 1992, would remain.


5. STATUS OF THE STOCKS

Stock structure assumptions

Snake bank is assumed to be a single stock.

COC 1A

Stock Status																															
Year of Most Recent Assessment	2009																														
Assessment Runs Presented	Survey biomass estimate for ≥ 30 mm shell length																														
Reference Points	Target: Not defined, but B_{MSY} assumed Soft Limit: 20% B_0 Hard Limit: 10% B_0 Overfishing Threshold: -																														
Status in relation to Target	About as Likely as Not (40–60%) to be at or above the target																														
Status in relation to Limits	Unlikely (< 40%) to be below both soft and hard limits																														
Historical Stock Status Trajectory and Current Status <table border="1"> <caption>Approximate data points for Historical Stock Status Trajectory</caption> <thead> <tr> <th>Year</th> <th>%B0</th> </tr> </thead> <tbody> <tr><td>1982</td><td>100</td></tr> <tr><td>1984</td><td>95</td></tr> <tr><td>1986</td><td>70</td></tr> <tr><td>1988</td><td>55</td></tr> <tr><td>1990</td><td>40</td></tr> <tr><td>1992</td><td>35</td></tr> <tr><td>1994</td><td>60</td></tr> <tr><td>1996</td><td>35</td></tr> <tr><td>1998</td><td>55</td></tr> <tr><td>2000</td><td>25</td></tr> <tr><td>2002</td><td>20</td></tr> <tr><td>2004</td><td>45</td></tr> <tr><td>2006</td><td>35</td></tr> <tr><td>2008</td><td>60</td></tr> </tbody> </table>		Year	%B0	1982	100	1984	95	1986	70	1988	55	1990	40	1992	35	1994	60	1996	35	1998	55	2000	25	2002	20	2004	45	2006	35	2008	60
Year	%B0																														
1982	100																														
1984	95																														
1986	70																														
1988	55																														
1990	40																														
1992	35																														
1994	60																														
1996	35																														
1998	55																														
2000	25																														
2002	20																														
2004	45																														
2006	35																														
2008	60																														
Recruited biomass (≥ 30 mm shell length) over time as a percentage of B_0 in relation to the hard and soft limits.																															
Fishery and Stock Trends																															
Recent Trend in Biomass or Proxy	The stock status in 2009 was at 35% of B_0 and has varied between 19 and 63% of B_0 since 1988, following a decline from 1982–1991.																														

Recent Trend in Fishing Mortality or Proxy	<p>Exploitation rate (≥ 30 mm shell length) generally trended downward from 1991 (70%) until 2012 (0%), with the exception of a large peak in rate around 2001 (up to 93%). It is Exceptionally Unlikely ($< 1\%$) that overfishing is occurring.</p>  <table border="1"> <caption>Exploitation rate (%) data (approximate values from graph)</caption> <thead> <tr> <th>Year</th> <th>Exploitation rate (%)</th> </tr> </thead> <tbody> <tr><td>1991</td><td>70</td></tr> <tr><td>1992</td><td>40</td></tr> <tr><td>1993</td><td>35</td></tr> <tr><td>1994</td><td>30</td></tr> <tr><td>1995</td><td>35</td></tr> <tr><td>1996</td><td>55</td></tr> <tr><td>1997</td><td>50</td></tr> <tr><td>1998</td><td>55</td></tr> <tr><td>1999</td><td>40</td></tr> <tr><td>2000</td><td>70</td></tr> <tr><td>2001</td><td>93</td></tr> <tr><td>2002</td><td>50</td></tr> <tr><td>2003</td><td>20</td></tr> <tr><td>2004</td><td>25</td></tr> <tr><td>2005</td><td>15</td></tr> <tr><td>2006</td><td>15</td></tr> <tr><td>2007</td><td>10</td></tr> <tr><td>2008</td><td>5</td></tr> <tr><td>2009</td><td>10</td></tr> <tr><td>2010</td><td>5</td></tr> <tr><td>2011</td><td>0</td></tr> <tr><td>2012</td><td>0</td></tr> <tr><td>2013</td><td>0</td></tr> </tbody> </table>	Year	Exploitation rate (%)	1991	70	1992	40	1993	35	1994	30	1995	35	1996	55	1997	50	1998	55	1999	40	2000	70	2001	93	2002	50	2003	20	2004	25	2005	15	2006	15	2007	10	2008	5	2009	10	2010	5	2011	0	2012	0	2013	0
Year	Exploitation rate (%)																																																
1991	70																																																
1992	40																																																
1993	35																																																
1994	30																																																
1995	35																																																
1996	55																																																
1997	50																																																
1998	55																																																
1999	40																																																
2000	70																																																
2001	93																																																
2002	50																																																
2003	20																																																
2004	25																																																
2005	15																																																
2006	15																																																
2007	10																																																
2008	5																																																
2009	10																																																
2010	5																																																
2011	0																																																
2012	0																																																
2013	0																																																
Other Abundance Indices	-																																																
Trends in Other Relevant Indicators or Variables	-																																																

Projections and Prognosis	
Stock Projections or Prognosis	-
Probability of Current Catch or TACC causing Biomass to remain below or to decline below Limits	Fishing at present levels is Exceptionally unlikely ($< 1\%$) to cause declines below soft or hard limits.
Probability of Current Catch or TACC causing Overfishing to continue or to commence	-

Assessment Methodology and Evaluation		
Assessment Type	Level 2: Partial quantitative stock assessment	
Assessment Method	Absolute biomass estimates from quadrant surveys	
Assessment Dates	Latest assessment: 2009	Next assessment: Unknown
Overall assessment quality rank		
Main data inputs (rank)	- Abundance - Length frequency	
Data not used (rank)	-	
Changes to Model Structure and Assumptions	-	
Major sources of Uncertainty	- The estimate of B_0 was from 1982 and is not necessarily a good estimate of average unfished biomass. - Maturity at length.	

Qualifying Comments
Water quality issues have influenced the amount of time when cockles can be harvested from the bank in recent years, e.g. the fishery was closed for 96 days in the 2009–10 year due to poor water quality.

The % $B_{recruited}$ and the exploitation rate are likely to be underestimate and overestimate, respectively as they are based on a 30 mm shell length and the size limit for commercial harvest is believed to have decreased from 30 to 28 mm over time.

Fishery Interactions

Interactions with other species are currently being characterised.


7. FOR FURTHER INFORMATION

- Anderson, M J (2008) Animal-sediment relationships re-visited: Characterising species' distributions along an environmental gradient using canonical analysis and quantile regression splines. *Journal of Experimental Marine Biology and Ecology* 366(1–2): 16–27.
- Boyd, R O; Gowing, L; Reilly, J L (2004) 2000–2001 national marine recreational fishing survey: diary results and harvest estimates. Final Research Report for Ministry of Fisheries project REC2000/03. 81 p. (Unpublished report held by Fisheries New Zealand, Wellington.)
- Boyd, R O; Reilly, J L (2002) 1999/2000 National marine recreational fishing survey: harvest estimates. Final Research Report for Ministry of Fisheries project REC98/03. 28 p. (Unpublished report held by Fisheries New Zealand, Wellington.)
- Bradford, E. (1997) Estimated recreational catches from Ministry of Fisheries North region marine recreational fishing surveys, 1993–94. New Zealand Fisheries Assessment Research Document 97/17. 16 p. (Unpublished report held in NIWA library, Wellington.)
- Bradford, E (1998) Harvest estimates from the 1996 national marine recreational fishing surveys. New Zealand Fisheries Assessment Research Document 98/16. 27 p. (Unpublished report held in NIWA library, Wellington.)
- Cryer, M (1997) Assessment of cockles on Snake Bank, Whangarei Harbour, for 1996. New Zealand Fisheries Assessment Research Document 1997/2. 29 p. (Unpublished report held in NIWA library, Wellington.)
- Cryer, M; Holdsworth, J (1993) Productivity estimates for Snake Bank cockles, August 1992 to August 1993. (Unpublished Internal Report, held by Fisheries New Zealand, Wellington.)
- Cryer, M; Smith, M; Parkinson, D; MacKay, G; Tasker, R (2003) Biomass surveys of cockles in Whangarei Harbour, 2002. Final Research Report for Ministry of Fisheries Project COC2001/01, Objective 3. 6 p. (Unpublished report held by Fisheries New Zealand, Wellington.)
- Cryer, M; Watson, T G; Smith, M D; MacKay, G; Tasker, R (2004) Biomass survey and stock assessment of cockles on Snake Bank, Whangarei Harbour, 2003. Final Research Report for Ministry of Fisheries Research Project COC2002/01, (Unpublished report held by Fisheries New Zealand, Wellington.)
- Griffiths, R, Eyre, R (2014) Population and Biomass Survey of Cockles (*Austrovenus stutchburyi*) in Whāngārei Harbour 2014. Report by Northland Regional Council (available on request) 19 p.
- Holdsworth, J; Cryer, M (1993) Assessment of the cockle, *Chione stutchburyi*, resource and its associated fishery in Whangarei Harbour. Unpublished Report held at NIWA, Auckland.
- Larcombe, M F (1971) The ecology, population dynamics, and energetics of some soft shore molluscs. Unpublished PhD thesis. University of Auckland, Auckland, New Zealand. 250 p.
- Martin, N D (1984) *Chione stutchburyi* population responses to exploitation. Unpublished MSc Thesis, University of Auckland, Auckland.
- McKenzie, J R; Cryer, M; Breen, P A; Kim, S (2003) A length-based model for cockles on Snake Bank, Whangarei Harbour, 2002. Final Research Report for Ministry of Fisheries Research Project COC2001/01, Objective 2. (Unpublished report available from Fisheries New Zealand, Wellington.)
- Morrison, M; Cryer, M (1999) Stock assessment of cockles on Snake and McDonald Banks, Whangarei Harbour, 1998. New Zealand Fisheries Assessment Research Document 99/7. (Unpublished report held by NIWA library, Wellington.)
- Morrison, M; Parkinson, D (2001) Stock assessment of cockles on Snake Bank and McDonald Banks, Whangarei Harbour, 2000. *New Zealand Fisheries Assessment Report 2001/19*.
- Morton, J; Miller, M (1968) *The New Zealand Sea Shore*. Liverpool, Collins.
- Sullivan, K J; Mace, P M; Smith, N W M; Griffiths, M H; Todd, P R; Livingston, M E; Harley, S J; Key, J M; Connell, A M (Comps.) (2005) Report from the Fishery Assessment Plenary, May 2005: stock assessments and yield estimates. 792 p. (Unpublished report held in NIWA library, Wellington.)
- Teirney, L D; Kilner, A R; Millar, R E; Bradford, E; Bell, J D (1997) Estimation of recreational catch from 1991–92 to 1993–94 New Zealand Fisheries Assessment Research Document 97/15. 43 p. (Unpublished report held by NIWA library, Wellington.)
- Thrush, S F; Hewitt, J E; Norkko, A; Nicholls, P E; Funnell, G E; Ellis, J I (2003) Habitat change in estuaries: predicting broad-scale responses of intertidal macrofauna to sediment mud content. *Marine Ecology-Progress Series* 263: 101–112.
- Watson, T G; Cryer, M; Smith, M D; MacKay, G; Tasker, R (2005) Biomass survey and stock assessment of cockles on Snake Bank, Whangarei Harbour, 2004. *New Zealand Fisheries Assessment Report 2005/15*. 39 p.
- Williams, J R; Cryer, M; McKenzie, J R; Smith, M D; Watson, T G; MacKay, G; Tasker, R (2006a) Biomass survey and stock assessment of cockles (*Austrovenus stutchburyi*) on Snake Bank, Whangarei Harbour, 2005. *New Zealand Fisheries Assessment Report 2006/21*. 21 p.
- Williams, J R; Smith, M D; MacKay, G (2006b) Biomass survey and stock assessment of cockles (*Austrovenus stutchburyi*) on Snake Bank, Whangarei Harbour, 2006. *New Zealand Fisheries Assessment Report 2006/38*. 21 p.
- Williams, J R; Smith, M D; MacKay, G (2008) Biomass survey and stock assessment of cockles (*Austrovenus stutchburyi*) on Snake Bank, Whangarei Harbour, 2007. *New Zealand Fisheries Assessment Report 2008/43*. 22 p.
- Wynne-Jones, J; Gray, A; Hill, L; Heinemann, A (2014) National Panel Survey of Marine Recreational Fishers 2011–12: Harvest Estimates. *New Zealand Fisheries Assessment Report 2014/67*.

COCKLES (COC 3) Otago Peninsula

(*Austrovenus stutchburyi*)

Tuaki



1. FISHERY SUMMARY

COC 3 was introduced into the Quota Management System in October 2002 with a TAC of 1500 t; comprising of a customary allowance of 10 t, a recreational allowance of 10 t, an allowance for other fishing related mortality of 10 t, and a TACC of 1470 t. Historical catch limits can be seen in Table 1.

1.1 Commercial fisheries

Cockles are present at various locations around the Otago Peninsula but are only commercially fished from Papanui Inlet, Waitati Inlet, and Otago Harbour (under a current special permit). Commercial fishing in Papanui and Waitati Inlets began in 1983. A limit of 104 t was in effect for Papanui and Waitati Inlets combined from 1986–87 until 1991–92. From 1992–93 to 1998–99, the catch limits were 90 t for Papanui Inlet and 252 t for Waitati Inlet. In April 2000, the catch limits were increased to 427 t for Papanui Inlet and 746 t for Waitati Inlet. In 2002 when cockles entered the QMS spatial restrictions upon harvest within COC 3 were removed. Commercial landings from Papanui and Waitati Inlets are shown in Table 1. Since August 2009 cockles have been taken from Otago Harbour under a special permit in order to investigate the ecosystem effects of commercial cockle harvesting in this location. This permit stated no explicit limit to the tonnage able to be taken but delimited the area where harvest would be taken, and expired on January 31st 2017.

In 1992, 35 mm shell length was the minimum size for commercial cockles. However, commercial fishers currently target cockles 28 mm or more, therefore 28 mm is used as the effective minimum size in yield calculations. CPUE data are available for this fishery, but have not been analysed.

COCKLES (COC 3)

Table 1: Reported landings (t) of cockles from Papanui and Waitati Inlets, Otago, combined (FMA 3), from 1986–87 to 2016–17 based on Licensed Fish Receiver Returns (LFRR). Catch splits are provided by Southern Clams Ltd and are partially from Stewart (2006). N/A = Not Applicable.

Year	Papanui catch (t)	Papanui limit (t)	Waitati catch (t)	Waitati limit (t)	Otago Harbour catch (t)	Total catch (t)	Total limit (t)
1986–87	14	–	–	–	–	14	104
1987–88	8	–	–	–	–	8	104
1988–89	5	–	–	–	–	5	104
1989–90	25	–	–	–	–	25	104
1990–91	90	–	16	–	–	106	104
1991–92	90	–	14	–	–	104	104
1992–93	90	90	92	252	–	182	342
1993–94	90	90	109	252	–	199	342
1994–95	90	90	252	252	–	342	342
1995–96	90	90	252	252	–	342	342
1996–97	90	90	252	252	–	342	342
1997–98	90	90	252	252	–	342	342
1998–99	90	90	293	252	–	383	342
1999–00	118	427	434	746	–	552	1 273
2000–01	90	427	606	746	–	696	1 273
2001–02	49	N/A	591	N/A	–	640	1 273
2002–03	52	N/A	717	N/A	–	767	1 470
2003–04	73	N/A	689	N/A	–	762	1 470
2004–05	91	N/A	709	N/A	–	800	1 470
2005–06	68	N/A	870	N/A	–	943	1 470
2006–07	0*	N/A	907	N/A	–	907	1 470
2007–08	–	N/A	760	N/A	–	760	1 470
2008–09	–	N/A	751	N/A	24	775	1 470
2009–10	–	N/A	379	N/A	441	820	1 470
2010–11	–	N/A	240	N/A	596	836	1 470
2011–12	–	N/A	358	N/A	437	795	1 470
2012–13						790	1 470
2013–14						800	1 470
2014–15						815	1 470
2015–16						923	1 470
2016–17						967	1 470

*No catches have been taken from Papanui Inlet since 2006–07 because of water quality problems.




Figure 1: Reported commercial landings and TACC for COC 3 (Otago).

1.2 Recreational fisheries

Cockles are taken by recreational fishers in many areas of New Zealand. The recreational fishery is harvested entirely by hand digging. Relatively large cockles are preferred.

No recreational harvest estimates specific to the COC 3 commercial fishery areas are available. Amateur harvest levels in FMA 3 were estimated by telephone-diary surveys in 1993–94 (Teirney et al 1997), 1996 (Bradford 1998) and 2000 (Boyd & Reilly 2002) and are given in Table 2. Harvest weights are estimated using an assumed mean weight of 25 g (for cockles over 30 mm).

The harvest estimates provided by telephone-diary surveys between 1993 and 2001 are no longer considered reliable for various reasons. A Recreational Technical Working Group concluded that these harvest estimates should be used only with the following qualifications: a) they may be very inaccurate; b) the 1996 and earlier surveys contain a methodological error; and c) the 2000 and 2001 estimates are implausibly high for many important fisheries. In response to these problems and the cost and scale challenges associated with onsite methods, a National Panel Survey was conducted for the first time throughout the 2011–12 fishing year. The panel survey used face-to-face interviews of a random sample of 30 390 New Zealand households to recruit a panel of fishers and non-fishers for a full year. The panel members were contacted regularly about their fishing activities and harvest information collected in standardised phone interviews. Harvest estimates for cockles in FMA 3 are given in Table 2 (from Wynne-Jones et al 2014).

Table 2: Estimated numbers of cockles harvested by recreational fishers in FMA 3, and the corresponding harvest tonnage. Figures were extracted from telephone-diary survey in 1993–94, 1996 and 2000, and from the National Panel Survey in 2011–12. Harvest weights were estimated using an assumed mean weight of 25 g.

Survey	Harvest (N)	% CV	Harvest (t)	Reference
1993–94 South	106 000	51	2.7	Teirney et al (1997)
1996	144 000	–	3.6	Bradford (1998)
2000	1 476 000	45	36.9	Boyd & Reilly (2002)
2011–12	300 158	67	7.5	Wynne-Jones et al (2014)

1.3 Customary non-commercial fisheries

Many intertidal bivalves, including cockles, are very important to Maori as traditional food, particularly to Huirapa and Otakou Maori in the Otago area. Tangata tiaki issue customary harvest permits for cockles in Otago. The number of cockles harvested under customary permits is given in Table 3, and is likely to be an underestimate of customary harvest.

On 1 October 2010, on the recommendation of the Taiapure Committee, the Minister of Fisheries introduced new regulations for the East Otago Taiāpure¹. These included a new amateur daily bag limit of 50 for shellfish, including cockles, and a ban on the commercial take of cockles from any part of the Taiapure, except for the existing sanitation areas within Waitati Inlet. The new regulations reflect the Committee's concern about fishing pressure on shellfish stocks, including cockles, within the Taiāpure.

A long-running time series of surveys suggest that there are no sustainability concerns for cockles within the Taiāpure. However, they do indicate a shift in some beds towards smaller size classes of cockle. Larger cockles are preferred by both customary and recreational fishers. The Committee hopes that reducing the bag limit and limiting the spatial extent of commercial harvest will lead to an increase in the number of large cockles.

Table 3: Number of cockles harvested under customary fishing permits.

Year	Number of cockles
1998	750
1999	0
2000	1 109
2001	1 090
2002	0
2003	2 750
2004	4 390
2005	5 699

¹ The Kati Huirapa Runanga ki Puketeraki application for a taiāpure-local fishery was gazetted as the East Otago Taiāpure-Local Fishery in 1999. A management committee, made up of representatives from the Runanga and various recreational, environmental, commercial, community and scientific groups, was appointed in 2001.

COCKLES (COC 3)

Fisheries New Zealand reports indicate that for the years since 2007 customary take under the regulations for COC 3 (which includes Otago Harbour) have been:

Year	Number of cockles
2007	7 680
2008	800
2009	23 565
2010	19 450
2011	11 800
2012	3 600
2014	2 100
2016	6 310

It is understood that local customary fishers generally utilise the daily amateur bag for their customary needs.

1.4 Illegal catch

No quantitative information is available on the magnitude of illegal catch but it is thought to be insignificant.

1.5 Other sources of mortality

No quantitative information is available on the magnitude of other sources of mortality. It has been suggested that some harvesting implements, such as brooms, rakes, “hand-sorters”, bedsprings and “quickfeeds” cause some incidental mortality, particularly of small cockles, but this proposition has not been scientifically investigated. High-grading of cockles is also practised, with smaller sized cockles being returned to the beds. The mortality from this activity is unknown, but is likely to be low.

2. STOCKS AND AREAS

Each inlet is assumed to be an independent fishery within the stock.

3. STOCK ASSESSMENT

Stock assessments for Papanui Inlet and Waitati Inlet have been conducted using absolute biomass surveys, yield-per-recruit analyses, and Method 1 for estimating CAY (Annala et al 2003). Breen et al (1999) also estimated biomasses and yields for Otago Harbour and Purakanui. Stewart (2006, 2008a) estimated biomass and yields for Papanui and Waitati Inlets in 2004 and Waitati Inlet in 2007.

3.1 Estimates of fishery parameters and abundance

A project to estimate growth and mortality in Papanui and Waitati Inlets, Purakanui and Otago Harbour was undertaken in the late 1990s. Notched clams did not exhibit significant growth when recovered after one year, and modes in the length frequency distributions did not shift when measured over four sampling periods within a year (Breen et al 1999).

Yield-per-recruit modelling has been conducted for Papanui and Waitati inlets separately (Stewart 2006, 2008a, Jiang et al 2011). The most recent parameters used in this modelling are detailed in table 2 of the cockle introductory section. Estimates of $F_{0.1}$ from these studies are given in Table 4 below. Exploitation rate is below 7% for Waitati, Papanui Inlet and Otago harbour (Table 4a, Figure 2).

Table 4: Estimates of fishery parameters (recruitment to this fishery is at ≥ 28 mm)

M	$F_{0.1}$ 2004	$F_{0.1}$ 2007	$F_{0.1}$ 2011	
			Waitati	Papanui
0.2	0.2321	0.2899	0.2600	0.2900
0.3	0.3412	0.3863	0.3900	0.4400
0.4	0.4767	0.5537	0.5300	0.6000

Table 4a: Exploitation rate % (for cockles ≥ 30 mm across each entire inlet)*

Year	Papanui	Waitati	Otago Hbr
1998	2	0	
2002	1	5	
2004	2	6	
2007	0	7	0
2011	0	2	4

* This measure is likely to overestimate exploitation as harvest occurs down to a size limit of 28mm.




Figure 2: Exploitation rate as calculated by landings divided by biomass (≥ 30 mm) from whole inlets. Note: This measure is likely to overestimate exploitation as harvest occurs down to a size limit of 28 mm.

3.2 Biomass estimates

Biomass surveys have been undertaken periodically in COC 3 since 1984. The methods for the calculation of biomass have changed over time² which means that comparison of biomass values between times of different calculation methodologies should be done cautiously.

The Spawning stock biomass (19 mm or more, shell length) has been stable around the level of virgin biomass in Waitati Inlet (Table 5, Figure 3). In Papanui Inlet the spawning stock biomass (19 mm or more shell length) has shown a trend of gradual decline from 1984 until 2011, when it was at 73% of virgin biomass (notably no commercial harvesting has occurred in Papanui Inlet since 2006–07). The recruited biomass (30 mm or more shell length) in the sanitation areas (beds 1804 and 1805) in Otago Harbour decreased before the start of harvesting in 2008 and has decreased more since then (to 60% of virgin biomass). A new survey was conducted in January 2017. From 164 stations at bed 1804 and 176 stations at bed 1805 the total clam biomass for each bed was estimated to be 4549 tonnes for 1804 and 4829 tonnes for 1805.

Table 5: Current ($\pm 95\%$ CI) and previous biomass estimates from COC 3*.

Papanui Inlet							
Size Class	1984	1992	1998	2002	2004	2004	2011
					Total inlet	Commercial area	Total inlet
>2 to 18 mm (juveniles)	65	139	33	17 \pm 1.7	36 \pm 2.2	13 \pm 1.3	8 \pm 1.4
19 – 34 mm (adults)	3 705	3 721	3 435	1 970 \pm 192	2 415 \pm 151	825 \pm 88	1 400 \pm 168
≥ 35 mm	2 370	1 706	2 231	2 579 \pm 252	2 301 \pm 273	1 847 \pm 208	3 048 \pm 429
≥ 30 mm			3 990	3 860 \pm 365	3 677 \pm 367	2 420 \pm 271	4 025 \pm 542
Total (t)	6 140	5 567	5 699	4 565 \pm 424	4 752 \pm 425	2 685 \pm 298	4 457 \pm 601

² Wildish (1984a and b) and Stewart et al (1992) separated cockles by sieving into three size classes. Breen et al (1999) measured random samples of cockles from each inlet to calculate length-weight relationships. The first method only allows estimation of biomass from predetermined size classes. By calculating size structure of populations using length to weight data, a more flexible approach is allowed where data can be matched to current commercial needs as well as to future survey results. The 1998 survey used random samples from each inlet to calculate length to weight relationships (Breen et al 1999). This method was once again used in the 2002 survey (Wing et al 2002). In the 2004 and 2007 surveys random samples from each shellfish bed were weighed and their longest axis measured (Stewart 2006, 2008a). These data were then used to generate length to weight relationships. The 2017 survey replicated the method used in the 2004 and 2007 surveys.

COCKLES (COC 3)

Table 5 [Continued]

Waitati Inlet.**

Size Class	1984	1992	1998	2002	2004 Total Inlet	2004 Commercial area	2007 Total Inlet	2007 Commercial area	2011 Total Inlet
>2 to 18 mm (juveniles)	619	1 210	304	153 ± 20	257 ± 14	77 ± 4	335 ± 26	102 ± 7.5	220 ± 14
19 to 34 mm (adults)	7 614	5 198	8 519	6 653 ± 652	7 272 ± 403	2 735 ± 129	7 673 ± 591	1 284 ± 95* ³	7 348 ± 501
≥35 mm	3 844	4 620	4 381	4 298 ± 298	4 535 ± 508	3 872 ± 384	3 941 ± 462		6 323 ± 643
≥30 mm			7 235	7 183 ± 463	7 993 ± 720	5 612 ± 681	7 107 ± 548	4 726 ± 352	11 441 ± 946
Total (t)	12 080	11 027	13 204	11 103 ± 848	12 064 ± 925	6 685 ± 517	11 948 ± 921	6 112 ± 456	13 892 ± 1149

Purkaunui Inlet

Size Class	1998	2008	2012
≥30 mm	1 825		

Otago Harbour

Size Class	1998	2008	2012
≥30 mm	32 975		

Otago Harbour (sanitation area, 1804)

Size Class	1998	2008	2012	2017
≥30 mm	8 901*	5 473	4 169	4 549

Otago Harbour (sanitation area 1805)

Size Class	1998	2008	2012	2017
, ≥30 mm	5 546*	3 526	4 093	4 829

*Wildish 1984a; Stewart et al 1992; Breen et al 1999; Wing et al 2002; Stewart, 2006; Stewart 2008a, Stewart 2008b; Jiang et al 2011; Stewart 2013. Area of current commercial beds, Papanui Inlet = 815 811 m². **Area of current commercial beds, Waitati Inlet = 943 986 m². *³ = this value is only for ≥19 mm to <30 mm cockles. *⁴The survey of Breen et al 1999 covered a larger extent on these beds than the two subsequent surveys of Stewart 2008b and 2013.




Figure 3: Biomass as a proportion of B_0 for Waitati and Papanui Inlets, this is estimated from biomass >19mm. Note: No catch has been taken from Papanui Inlet since 2006–07. Virgin biomass was taken from the Stewart 2008b survey for Otago harbour as this is the extent that has been subsequently surveyed.

3.3 Yield estimates and projections

Estimates of MCY are given in Table 6.

For Waitati Inlet, CAY was estimated (Table 7) using Method 1 ($CAY = (F_{0.1}/Z) (1 - \exp(-Z)) B_{BEG}$) (Annala et al 2003) and biomass estimates at different times. CAY has been estimated at times for both the entire inlet area and a subset area where the commercial fishery has been operating for the past several years. This approach assumes that, between the start of the fishing year and when the biomass survey is started, productivity and catch cancel each other.

Table 6: Estimates of $MCY(t)$ for COC 3 generated using Method 1 (Annala et al 2003) $MCY = 0.5F_{0.1}B_{AV}$, an average biomass ≥ 30 mm as B_0 and the 2011 estimates of $F_{0.1}$. This calculation is likely to underestimate the true MCY .

Location	M	1998	2002	2004	2007	2011
Waitati Inlet	0.2	941	934	1039	924	1487
Waitati Inlet	0.3	1411	1401	1559	1386	2231
Waitati Inlet	0.4	1917	1903	2118	1883	3032
Waitati Inlet (commercial)	0.2			730	614	894
Waitati Inlet (commercial)	0.3			1094	922	1342
Waitati Inlet (commercial)	0.4			1487	1252	1823
Papanui Inlet	0.2	579	560	533		584
Papanui Inlet	0.3	878	849	809		886
Papanui Inlet	0.4	1197	1158	1103		1208
Papanui Inlet (commercial)	0.2			351		259
Papanui Inlet (commercial)	0.3			532		392
Papanui Inlet (commercial)	0.4			726		535

Table 7: CAY estimates (t) for COC 3. WI = Waitati Inlet, PI = Papanui Inlet, WIc and PIc are estimates for commercial areas only, B_{beg} = Projected biomass at the beginning of the fishing year.

Year	M	$F_{0.1}$	$\geq SL$ (mm)	WI		WIc		PI		PIc		Reference
				B_{beg}	CAY	B_{beg}	CAY	B_{beg}	CAY	B_{beg}	CAY	
2011	0.2	0.26	30	11 441	2 385	6881	1434					Jiang et al 2011
2011	0.3	0.39	30	11 441	3 223	6881	1938					Jiang et al 2011
2011	0.4	0.53	30	11 441	3 948	6881	2374					Jiang et al 2011
2011	0.2	0.29	30					4 026	923	1784	409	Jiang et al 2011
2011	0.3	0.44	30					4 026	1 252	1784	555	Jiang et al 2011
2011	0.4	0.60	30					4 026	1 527	1784	677	Jiang et al 2011
2007	0.2	0.2899	28	8 378	1 920	5 261	1 206					Stewart 2008a
2007	0.3	0.3863	28	8 378	2 342	5 261	1 471					Stewart 2008a
2007	0.4	0.5537	28	8 378	2 990	5 261	1 878					Stewart 2008a
2007	0.2	0.2899	30	7 106	1 629	4 725	1 083					Stewart 2008a
2007	0.3	0.3863	30	7 106	1 986	4 725	1 321					Stewart 2008a
2007	0.4	0.5537	30	7 106	2 536	4 725	1 686					Stewart 2008a
2004	0.2	0.2321	30	9 399	1 771	6 081	1 146	4 119	776	2 454	462	Stewart 2006
2004	0.3	0.3412	30	9 399	2 367	6 081	1 532	4 119	1 038	2 454	618	Stewart 2006
2004	0.4	0.4767	30	9 399	2 984	6 081	1 930	4 119	1 308	2 454	779	Stewart 2006
2002	0.2	0.2017	30	7 183	1 193	5 364	891	3 860	641	2 322	386	Wing et al 2002
2002	0.3	0.3015	30	7 183	1 627	5 364	1 215	3 860	874	2 322	526	Wing et al 2002
2002	0.4	0.3956	30	7 183	1 960	5 364	1 464	3 860	1 053	2 322	634	Wing et al 2002
1999	0.2	0.258	30	7 235	1 498			3 990	826			Breen et al 1999
1999	0.3	0.357	30	7 235	1 848			3 990	1 019			Breen et al 1999
1999	0.4	0.457	30	7 235	2 221			3 990	1 225			Breen et al 1999

3.4 Other factors

Commercial, customary and recreational fishers target different sized cockles. Biomass and yield estimates will differ for different sizes of recruitment to the fishery. Maori and recreational fishers prefer larger cockles (45 mm shell length and greater) whereas commercial fishers currently prefer cockles of around 28–34 mm. Estimates of yields have been estimated for size of recruitment at 28 mm; however, these estimates do not consider multiple fisheries preferring different sized cockles. Depending on the management approach taken in the future in COC 3, the appropriateness of the current methods to estimate yield may need to be reviewed.

The yield estimates use information from yield-per-recruit analyses that assume constant recruitment and constant growth and mortality rates. Yield estimates will be improved when growth, mortality and recruitment variation are better known.

As cockles become sexually mature at around 18 mm, using a size of recruitment of 30 mm should provide some protection against egg overfishing under most circumstances. Certainly the increase in the biomass of small cockles (2 to 18 mm) seen in both inlets in 2004 suggests that the very poor recruitment observed by Wing et al (2002) may have been due to natural variability, and supports the conjecture that significant recruitment might occur only sporadically in the Otago fishery, as suggested by John Jillett (*pers. comm.*) and Breen et al (1999). The possibility that fishing has an effect on recruitment remains an unknown.

COCKLES (COC 3)

In other cockle fisheries it has been shown that recruitment of juvenile cockles can be reduced by the removal of a large proportion of adult cockles from a given area of substrate. This would suggest that there is some optimal level of adult biomass to facilitate recruitment, although its value is not known. To date it has not been determined whether the cockles being targeted by commercial harvesting in the Otago fishery comprise the bulk of the spawning stock or if disturbance of the cockle beds is influencing settlement.

The distribution of very small size classes (2 to 10 mm) across the various beds is variable and no consistent differences exist for this size of shellfish between commercial and non-commercial beds (Stewart 2008a). A comparison of the size/frequency histograms with fishing history for each bed would be a worthwhile exercise and may reveal more. The fact that the relationship between spawning stock and recruitment in this fishery is poorly understood remains a concern.

The very slight decrease in biomass recorded in the Stewart (2008a) survey suggests that the current level of harvest is sustainable. What is not known is if the decrease in biomass is the beginning of a long-term trend or simply the result of natural variability.

The effects of the illegal catch, the Maori traditional catch and incidental handling mortality are unknown, although illegal catch is thought to be insignificant. The impacts of the recreational fishery are probably minor compared with those from the commercial fishery.

4. STATUS OF THE STOCKS


Stock structure assumptions

Each inlet is assessed separately.

COC 3

Stock Status	
Year of Most Recent Assessment	2011
Assessment Runs Presented	Survey biomass estimate for ≥ 19 mm shell length
Reference Points	Target: Not defined, but B_{MSY} assumed Soft Limit: 20% B_0 Hard Limit: 10% B_0 Overfishing threshold: Not defined
Status in relation to Target	Likely ($> 60\%$) to be at or above the target
Status in relation to Limits	Very Unlikely ($< 10\%$) to be below both soft and hard limits
Status in relation to overfishing	Exploitation rate has never exceeded 7% at any of the harvested sites. It is Very Unlikely ($< 10\%$) that overfishing is occurring.

Historical Stock Status Trajectory and Current Status



Biomass as a proportion of B_0 for Waitati and Papanui Inlets, this is estimated from biomass > 19 mm. Note: No catch has been taken from Papanui Inlet since 2006–07. Virgin biomass was taken from the Stewart 2008b survey for Otago harbour as this is the extent that has been subsequently surveyed.


Fishery and Stock Trends

Recent Trend in Biomass or Proxy

Biomass at Waitati Inlet has been stable or increasing and has never decreased below 85% of B_0 . At Papanui Inlet biomass generally decreased to approximately 70% of B_0 in 2004 but little commercial catch has come out of this inlet since. In Otago Harbour biomass has declined, but most of this occurred before harvesting starting.

Recent Trend in Fishing Intensity or Proxy

Exploitation rate has never exceeded 7% at any of the harvested sites. It is Very Unlikely (< 10%) that overfishing is occurring.



Exploitation rate as calculated by landings divided by biomass (≥ 19 mm) from whole inlets (excluding Otago Harbour where it was taken from commercial beds only).

Other Abundance Indices

Trends in Other Relevant Indicators or Variables

-

Projections and Prognosis

Stock Projections or Prognosis

-

Probability of Current Catch or TACC causing Biomass to remain below or to decline below Limits

Fishing at recent levels is Very Unlikely (< 10%) to cause declines below soft or hard limits

Probability of Current Catch or TACC causing Overfishing to continue or to commence

-

Assessment Methodology and Evaluation		
Assessment Type	Level 2: Partial quantitative stock assessment	
Assessment Method	Absolute biomass estimates from quadrat surveys	
Assessment Dates	Latest assessment: 2010 or 2011 (depending upon location)	Next assessment: unknown
Overall assessment quality rank	-	
Main data inputs (rank)	- Abundance - Length frequency	
Data not used (rank)	-	
Changes to Model Structure and Assumptions	-	
Major Sources of Uncertainty	-	

Qualifying Comments
Water quality issues have influenced the amount of time when cockles can be harvested from Papanui Inlet in recent years.

Fishery Interactions
- The effects of harvesting on the composition of communities and substrate associated with the clam beds are under investigation.

5. FOR FURTHER INFORMATION


- Annala, J H; Sullivan, K J (1996) Report from the mid-year Fishery Assessment Plenary, November 1996: stock assessments and yield estimates. (Unpublished report held by Fisheries New Zealand, Wellington.)
- Annala, J H; Sullivan, K J; O'Brien, C J; Smith N W McL; Grayling, S M (2003) Report from the Fishery Assessment Plenary, May 2003: stock assessments and yield estimates. (Unpublished report held by Fisheries New Zealand.)
- Boyd, R O; Gowing, L; Reilly, J L (2004) 2000–2001 national marine recreational fishing survey: diary results and harvest estimates. Final Research Report for Ministry of Fisheries project REC2000/03. 81 p. (Unpublished report held by Fisheries New Zealand, Wellington.)
- Boyd, R O; Reilly, J L (2002) 1999/2000 National marine recreational fishing survey: harvest estimates. Final Research Report for Ministry of Fisheries project REC98/03. 28 p. (Unpublished report held by Fisheries New Zealand, Wellington.)
- Bradford, E (1998) Harvest estimates from the 1996 national recreational fishing surveys. New Zealand Fisheries Assessment Research Document 1998/16. 27 p. (Unpublished report held by NIWA library, Wellington.)
- Breen, P A; Carbines, G C; Kendrick, T H (1999) Stock assessment of cockles in Papanui and Waitati Inlets, Otago Harbour, and Purakanui, Otago. Final Report for Ministry of Fisheries research project COC9701. (Unpublished report held by Fisheries New Zealand, Wellington.)
- Cameron, T (1997) Aspects of growth and condition in *Austrovenus stutchburyi* (Finlay 1927 (Bivalvia: Veneridae) at Waitati Inlet: Influence of shore height and distance from mouth. Unpublished MSc thesis, University of Otago, Dunedin, New Zealand.
- Cranfield, H J; Michael, K P; Francis, R I C C (1996) Growth rates of five species of subtidal clam on a beach in the South Island, New Zealand. *Marine and Freshwater Research* 47: 773–784.
- Francis, R I C C (1984) An adaptive strategy for stratified random trawl surveys. *New Zealand Journal of Marine and Freshwater Research* 18: 59–71.
- Hilborn, R; Walters, C J (1992) *Quantitative Fisheries Stock Assessment: Choice, dynamics and uncertainty*. Chapman and Hall, London.
- Irwin, C R (1999) The effects of harvesting on the reproductive and population biology of the New Zealand Littleneck Clam (*Austrovenus stutchburyi*) in Waitati Inlet. Unpublished Msc thesis, University of Otago, Dunedin, New Zealand.
- Jiang, W; Goodwin, E; Stewart, B (2011) Stock Assessment of Clams (*Austrovenus stutchburyi*) in Papanui Inlet and Waitati Inlet, Otago, 2010–2011. Prepared for Southern Clams Ltd. Cawthron Report No. 1926. 34 p.
- McKinnon, J (1996) Studies of the age, growth and shell increment patterns in the New Zealand cockle (*Austrovenus stutchburyi*). Unpublished MSc thesis, University of Otago, Dunedin, New Zealand.
- Manly, B F J; Akroyd, J M; Walshe, K A R (2002) Two-phase stratified random surveys on multiple populations at multiple locations. *New Zealand Journal of Marine and Freshwater Research* 36: 581–591.
- Morton, J; Miller, M (1973) *The New Zealand Sea Shore*. Collins, Auckland. 653 p.
- Stewart, B (2004) Warrington treated Sewage Discharge: Waitati Inlet Ecological Survey, 2004. Prepared for Dunedin City Council. 21 p.
- Stewart, B (2006) Stock assessment of Cockles (*Austrovenus stutchburyi*) in Papanui and Waitati Inlets, Otago, 2004. Final Research Report for the Ministry of Fisheries Research Project COC2004/02. 54 p. (Unpublished report held by Fisheries New Zealand, Wellington.)
- Stewart, B (2008a) Stock assessment of clams (*Austrovenus stutchburyi*) in Waitati Inlet, Otago, 2007. Final Research Report for Southern Clams Ltd. 24 p.
- Stewart, B G (2008b) Clam (*Austrovenus stutchburyi*) resource and habitat survey in Otago Harbour (COC 3), Otago, 2008. Report prepared for Southern Clams Ltd. By Ryder Consulting.
- Stewart, B G (2013) Investigations into the Effects of Commercial Harvest of Clams (*Austrovenus stutchburyi*) in Otago Harbour (COC3), Otago: Report on Phase II Harvesting, 2012 Report prepared for Southern Clams Ltd. By Ryder Consulting. 66 p.
- Stewart, B; Keogh, J; Fletcher, D; Mladenov, P (1992) Biomass survey of the New Zealand littleneck clam (*Chione stutchburyi*) in Papanui and Waitati Inlets, Otago during 1991/1992. Marine Science and Aquaculture research Centre, University of Otago, Dunedin, New Zealand. 37 p.
- Sukhatme, P V (1954) *Sampling theory of surveys: with applications*. New Delhi : Indian Society of Agricultural Statistics. 49 p.

- Teirney, L D; Kilner, A R; Millar, R E; Bradford, E; Bell, J D (1997) Estimation of recreational catch from 1991–92 to 1993–94 New Zealand Fisheries Assessment Research Document 1997/15. 43 p. (Unpublished document held by NIWA library, Wellington.)
- Wildish, K (1984a) The cockle resource in the Otago region: 1. An initial study of the New Zealand cockle (*Chione stutchburyi*) resource in Papanui and Waitati Inlets. Report to the Ministry of Agriculture and Fisheries.
- Wildish, K (1984b) The cockle resource in the Otago region: further analysis of results from the survey of *Chione stutchburyi* (the New Zealand cockle) populations at Papanui and Waitati Inlets. Report to the Ministry of Agriculture and Fisheries.
- Wing, S; Irwin, C; Granger, G (2002) Biomass survey and yield estimates for the New Zealand littleneck clam *Austrovenus stutchburyi* in Papanui and Waitati Inlets, Otago. Final Report for the Ministry of Fisheries Research Project COC2001/02. 52 p. (Unpublished report held by Fisheries New Zealand, Wellington.)
- Wynne-Jones, J; Gray, A; Hill, L; Heinemann, A (2014) National Panel Survey of Marine Recreational Fishers 2011–12: Harvest Estimates. *New Zealand Fisheries Assessment Report 2014/67*.
- Zar, J H (1996) *Biostatistical Analysis*. Third Edition. Prentice Hall International Inc.

COCKLES (COC 7A) Tasman and Golden Bays

(*Austrovenus stutchburyi*)

Tuangi



1. FISHERY SUMMARY

COC 7A was introduced into the Quota Management System in October 2002 with a TAC of 1 510 t; comprising a customary allowance of 25 t, a recreational allowance of 85 t, an allowance for other fishing related mortality of 10 t, and a TACC of 1 390 t. These limits have remained unchanged since.

1.1 Commercial fisheries

Commercial harvesting at Pakawau Beach in Golden Bay began in 1984, but with significant landings taken only since 1986. Harvesting at Pakawau Beach has occurred every year since 1984. Cockles have also been taken commercially from Tapu Bay-Riwaka (in Tasman Bay) since 1992–93, and Ferry Point (in Golden Bay) since 1998–99. Catch statistics (Table 1) are derived from company records and QMS returns. All commercial landings have been taken by mechanical harvester. Historical landings and TACC for this stock are depicted in Figure 1.

Table 1: Reported landings (t) of cockles from all commercially harvested areas in COC 7A/7B. Landings from 1983–84 to 1991–92 are based on company records.

Fishing Year	Total Landings	TACC
1983–84	2	225
1984–85	38	225
1985–86	174	225
1986–87	230	225
1987–88	224	225
1988–89	265	300
1989–90	368	300
1990–91	535	300
1991–92	298	300
1992–93	300	336
1993–94	440	336
1994–95	326	336
1995–96	329	336
1996–97	325	336
1997–98	513	949
1998–99	552	1 130
1999–00	752	1 130
2000–01	731	1 134
2001–02	556	1 134
2002–03	569	1 390
2003–04	553	1 390
2004–05	428	1 390
2005–06	460	1 390
2006–07	337	1 390
2007–08	237	1 390
2008–09	307	1 390
2009–10	301	1 390
2010–11	348	1 390
2011–12	220	1 390
2012–13	269	1 390
2013–14	290	1 390
2014–15	263	1 390
2015–16	263	1 390
2016–17	238	1 390

COCKLES (COC 7A)

At Pakawau Beach, the fishery operated up to October 1988 under a special permit constraining annual landings to 225 t. From 1988–89 to 1997–98, the fishery operated under a commercial permit allowing an annual catch of 300 t. In 1997–98, the fishery was re-assessed and a catch limit of 913 t was set based on a CAY harvest strategy. This level of harvest was changed to 760 t from the 1998–99 fishing year and then 764 t for the 2000–01 fishing year. The harvest is taken from an area of about 500 ha.

The Ferry Point fishery, initiated in 1998–99, has an annual allowable catch of 334 t based on an MCY harvest strategy. The harvested area is about 40 ha. Reportedly, the area has not been fished since 2004. The Tapu Bay-Riwaka fishery, which was developed in 1990–91, has operated under a commercial permit limiting catches to 36 t annually. This fishery has been only lightly harvested owing largely to water quality issues and the area from which catches have been taken is probably less than 100 ha.




Figure 1: Total reported landings and TACC for COC 7A (Nelson Bays).

1.2 Recreational fisheries

Cockles are taken by recreational fishers, generally using hand digging. The catch limit is currently 150 cockles per person per day. Relatively large cockles (i.e., shell length over 30 mm) are generally preferred. Specific areas for recreational fishing are set aside from the commercial fishery by regulation and these include the area north of Ferry Point opposite Totara Ave and the area of Tapu Bay itself north of the fishery.

No estimates of recreational harvest of cockles from COC 7A are available. Early estimates of the amateur cockle harvest from the whole of FMA 7 are available from telephone-diary survey in 1992–93 (Teirney et al 1997), 1996 (Bradford 1998), and 2000 (Boyd & Reilly 2002). Harvest weights were estimated assuming a mean weight of 25 g per cockle (Table 2).

The harvest estimates provided by telephone-diary surveys between 1993 and 2001 are no longer considered reliable for various reasons. A Recreational Technical Working Group concluded that these harvest estimates should be used only with the following qualifications: a) they may be very inaccurate; b) the 1996 and earlier surveys contain a methodological error; and c) the 2000 and 2001 estimates are implausibly high for many important fisheries. In response to these problems and the cost and scale challenges associated with onsite methods, a National Panel Survey was conducted for the first time throughout the 2011–12 fishing year. The panel survey used face-to-face interviews of a random sample of 30 390 New Zealand households to recruit a panel of fishers and non-fishers for a full year. The panel members were contacted regularly about their fishing activities and harvest information collected in standardised phone interviews. Harvest estimates for cockles in FMA 7 are given in Table 2 (from Wynne-Jones et al 2014). The estimate for 2011–12 is lower than expected, potentially because of the number of toxic algal blooms in that year.

Table 2: Estimated numbers of cockles harvested by recreational fishers in FMA 7, and the corresponding harvest tonnage estimates calculated using an assumed mean weight of 25 g per cockle.

Survey	Number	CV	(t)	Reference
1992–93 telephone-diary	166 000	-	4	Teirney et al (1997)
1996 telephone-diary	325 000	-	8	Bradford (1998)
2000 telephone-diary	499 000	-	12.5	Boyd & Reilly (2002)
2011–12 national panel survey	78 751	0.45	2	Wynne-Jones et al (2014)

A repeat of the National Panel Survey is being conducted over the 2017–18 October fishing year. Results are expected in early 2019.

1.3 Customary non-commercial fisheries

Cockles are an important Maori traditional food, but no quantitative information on the level of customary take in COC 7A/7B is available. However, Kaitiaki are now in place in many areas and estimates of customary harvest can be expected to improve.

1.4 Illegal catch

No quantitative information on the level of illegal catch is available.

1.5 Other sources of mortality

The extent of any other sources of mortality is unknown. Incidences of unexplained large-scale die-off in localised areas have been noted (e.g., at Pakawau Beach and Ferry Point in 1999). Mortality of unrecruited cockles during the mechanical harvesting process was found to be very low (Bull 1984), and disturbance and mortality of other invertebrates in the harvested areas is slight (Wilson et al 1988).

2. BIOLOGY

All references to “shell length” in this report refer to the maximum linear dimension of the shell (in an anterior-posterior axis). General cockle biology has been summarised earlier in this Plenary report. Some aspects of biology with particular relevance to COC 7A follow.

Estimates of growth and mortality have been made for cockles from Pakawau Beach (Osborne 1992, 1999, 2010), and the two early studies are summarised in Table 3. The 1992 investigation used a Walford plot of tag recapture data (Bull 1984), and measured growth after about 18 months on translocated cockles, to produce the growth parameters. A MIX analysis of the scaled length-frequency distribution from the 1992 survey enabled calculation of the proportional reduction of the 4+ and 5+ age classes to produce estimates of instantaneous natural mortality, M (after removal of estimated fishing mortality, F).

The 1999 investigation used a MIX analysis of length-frequency data from two strata in comparable surveys in 1997, 1998 and 1999 to estimate mean lengths (and proportion in the population) of the first 8 year classes. Von Bertalanffy parameters were estimated for each survey. Mean natural mortality rates were estimated (for age classes 4–7) between 1997 and 1998, and 1998 and 1999.

Table 3: Estimates of biological parameters.

Population & years	Estimate		Source
<u>1. Natural mortality (<i>M</i>)</u>			
Pakawau Beach (1992)	0.45 for 4+; 0.30 for 5+		Osborne (1992, 1999)
Pakawau Beach (1998)	0.4		Osborne (1999)
Pakawau Beach (1999)	0.52		Osborne (1999)
<u>2. Weight = <i>a</i> (shell length)^b (weight in g, shell length in mm)</u>			
	<i>a</i>	<i>b</i>	
Pakawau Beach (1992)	0.000017	3.78	Osborne (1992)
Ferry Point (1996)	0.00020	3.153	Forrest & Asher (1997)
Tapu Bay-Riwaka (1991)	0.000150	3.249	Stark & Asher (1991)

COCKLES (COC 7A)

Table 3 [Continued]

Population & years	Estimate			Source
<u>3. von Bertalanffy growth parameters</u>				
	K	t_0	L_∞	
Pakawau Beach (1984–92)	0.36	0.3	49	Osborne (1992)
Pakawau Beach (1997)	0.38	0.68	48.3	Osborne (1999)
Pakawau Beach (1998)	0.4	0.68	47.4	Osborne (1999)
Pakawau Beach (1999)	0.41	0.66	47	Osborne (1999)

It was acknowledged that none of the MIX analyses converged, but the results presented were the best available fits (Osborne 1992, 1999). However, all four analyses produced very similar von Bertalanffy parameters. There is a trend of a reducing L_∞ and increasing K over the period 1992–1999, which might be expected as a result of fishing. In 2009 growth was modeled by the equation $y = 11.452\ln(x) + 16.425$, where y is shell width and x is age in years, this equation is only applicable to individuals 23–55 mm in shell width.

3. STOCKS AND AREAS

Little is known of the stock boundaries of cockles. The planktonic larval phase of this shellfish has a duration of about three weeks, so dispersal of larvae to and from a particular site could be considerable. Cockles are known to be abundant and widely distributed throughout Golden and Tasman Bays, and although nothing is known about larval dispersion patterns, cockles in these areas are likely to comprise a single stock. However, in the absence of any detailed information on stocks, the three currently fished sites in COC 7A are all managed as one stock.

4. STOCK ASSESSMENT

This report summarizes estimates of absolute biomass and yields for exploited and unexploited cockle populations in Tasman and Golden Bays. Stock assessments have been conducted using absolute biomass surveys, yield-per-recruit analyses, Methods 1 and 2 for estimating *MCY*, and Method 1 for estimating *CAY* (Ministry of Fisheries 2010).

Recruited cockles are considered to be those with a shell length of 30 mm or greater. This is the minimum size of cockles generally retained by the mechanical harvesters used in the COC 7A fishery. Where possible, estimates of yields from surveys are based on recruited biomass not occurring in areas of eel grass (*Zostera*), as the disturbance of these *Zostera* beds by mechanical harvesters has detrimental effects on intertidal ecology.

4.1 Estimates of fishery parameters and abundance

None available.

4.2 Biomass estimates

Biomass estimates from surveys are available for the three commercially fished areas and three other sites.

On Pakawau Beach, the surveys done in 1992 and 1997–2008 used a stratified random approach (Table 4, Figure 2). An additional southern stratum was added to the survey area in 1997 after legal definition of the fishery area, accounting for the greater survey area relative to 1992. The surveys in 1984 and 1988 covered smaller areas still. The survey area was reduced further in 2008 and 2014 to remove areas that were observed to be consistently unsuitable habitat for cockles or cockle harvesting (sand banks, soft mud and *Zostera* areas). The eight comparable surveys show total and recruited biomass to have fluctuated with no consistent trend, but the lowest value in this time series was recorded in 2014. In addition to recruited biomass (>30mm size), and vulnerable biomass (outside *Zostera* beds), reference biomass levels used for *MCY* calculation this year and in previous years are shown in Table 4.

Estimates of biomass are available for Tapu Bay-Riwaka in 1991 using a fixed transect approach (Stark & Asher 1991) and Ferry Point in 1996 using a stratified random approach (Forrest & Asher 1997). Both these surveys were conducted about two years prior to the commencement of commercial harvesting in those areas. The cockle resource on three other beaches in Golden Bay was assessed using stratified random surveys in 1993 (Osborne & Seager 1994). Since then both Riwaka and Ferry Point have been surveyed in 2004 and 2008 using stratified random survey designs. Results from all these surveys are listed in Table 5 and shown in Figure 2. The biomass at Riwaka and Ferry Point have generally decreased over time.

Table 4: Estimates of biomass with 95% confidence intervals where available for Pakawau Beach. Values are recruited (>30mm) and vulnerable biomass (not occurring in *Zostera* beds) and reference levels of biomass used for calculating MCY (B_0 virgin biomass, B_{av} average biomass). In 2014 vulnerable biomass was calculated differently (see Osborne 2014 for details).

	Recruited biomass				Vulnerable biomass				Assessed reference levels		
	Area	tonnes	95 % CI	CV	Area	tonnes	95 % CI	CV	B_0	B_{av}	95 % CI
1984	326	4604	1562	-	-	-	-	-	-	-	-
1988	510	5640	-	-	-	-	-	-	-	-	-
1992	588	6784	929			3586	612	8.7	3293	-	-
1997	642	7796	1628	10.7	275	3723	1331	18.2	-	3655	134
1998	642	6768	1221	9.0	317	3412	827	12.4	-	3574	176
1999	642	7502	1294	8.8	246	3058	727	12.1	-	3445	282
2000	642	7128	1237	8.9	266	2139	555	13.2	-	3184	556
2001	642	9117	1519	8.5	254	3111	712	11.7	-	3172	455
2004	642	9421	1195	6.5	307	5747	909	8.1	-	3539	817
2008	407	8285	1599	9.8	299	4954	1025	10.6	-	3716	788
2014	358	3363	561	8.5	358	3363	561	8.5	-	5686	1137

Table 5: Estimates of biomass (t) with 95% confidence intervals (CI) where available, and mean density (kg/m^2) for cockles at various sites in Golden and Tasman Bays. Where possible, values are given for the total and recruited (≥ 30 mm) populations. n = number of samples in the survey.

Site	Date	Area (ha)	n	Total biomass			Recruited biomass		
				t	CI	kg/m^2	t	CI	kg/m^2
Tapu Bay-Riwaka	Mar-91	306	321	~3 900	-	1.28	-	-	-
Riwaka	Feb-04	122.7	144	1 423	269	1.16	1 076	235.6	0.88
Riwaka	Mar-08	103	82	1475	257	1.44	939	178	0.9
Riwaka (excl. Tapu Bay)*	Mar-91	-	-	-	-	-	1 880	450	-
Ferry Point	Dec-96	40	552	2 617	190	5.99	2 442	191	5.6
Ferry Point	Feb-04	40	126	646	99.8	1.63	443	79	1.12
Ferry Point	Jan-08	28.2	75	662	112	2.35	470	83	1.7
Collingwood Beach	Mar-93	176	70	334	148	0.19	292	139	0.17
Takaka Beach	Mar-93	338	107	1 850	671	0.55	796	395	0.24
Rangihaeata Beach	Mar-93	197	75	473	345	0.24	438	320	0.22

* Recalculated by Breen (1996) from data in Stark & Asher (1991).

Surveys reporting on cockle abundance have also been produced for Motupipi, Golden Bay, in June 1995 (transect survey, 50 ha, 30 samples, mean density of 87 cockles per m^2 , no sizes or weights recorded), and at various sites in the Marlborough Sounds in August 1986 (diver survey below mean low water only, 9 sites, main densities in Kenepuru and inner Pelorus Sounds).

Absolute virgin biomass, B_0 , are assumed to be equal to estimated biomass of cockles 30 mm or over shell length from surveys conducted before, or in the early stages of, any commercial fishing. These are listed above in Tables 4 and 5. Absolute current biomass can be estimated similarly from current surveys.

COCKLES (COC 7A)




Figure 2: Recruited biomass (≥ 30 mm shell length) over time. Notably, the area surveyed over time has changed (see Tables 4 and 5) and decreased at the last time of survey (compared to previous occasions) at all three sites.

The biomass that will support the maximum sustainable yield (B_{MSY}) is not known for any of the areas fished in COC 7A.

4.3 Yield estimates and projections

Estimates of MCY have been made for populations of cockles in various areas, and at various times, using the equation $MCY = 0.25 * F_{ref} * B_0$ (Method 1), where F_{ref} is either $F_{0.1}$ or F_{max} . This method applies to new fisheries, or to those with only very low past levels of exploitation. The value of F_{ref} is dependent on M , so owing to the uncertainty of M a range of MCY estimates have been given for each stock (Table 6). For all estimates in Table 6, B_0 was taken as recruited biomass available for fishing (i.e. not in *Zostera* beds) in the survey area.

Estimates of MCY for Pakawau Beach have also been produced from $MCY = 0.5 * F_{REF} * B_{AV}$ (Method 2), using $F_{0.1}$, and with B_{AV} being the average of the available recruited biomass from the previous comparable surveys. For a range of M values, the latest estimates of MCY are as follows:

M	0.2	0.3	0.4
MCY	665	996	1 312

Table 6: Estimates of MCY (t, using $0.25 * F_{REF} * B_0$) for various cockle stocks in Tasman and Golden Bays, assuming a range of values for M .

Site	Date	F_{ref}	M			
			0.2	0.3	0.4	0.5
Pakawau Beach	1992	$F_{0.1}$	230	324	434	554
Pakawau Beach	1997	$F_{0.1}$	397	559	751	957
Pakawau Beach	2001	F_{MAX}	1 182	2 418	4 658	
Pakawau Beach	2004	$F_{0.1}$	482	683	924	
Pakawau Beach	2008	$F_{0.1}$	340	481	651	
Pakawau Beach	2014	$F_{0.1}$	665	996	1 312	
Ferry Point	1996	$F_{0.1}$	127	170	223	284
Ferry Point	1996	F_{MAX}	264	453	789	1 493
Ferry Point	2004	$F_{0.1}$	122	173	234	
Ferry Point	2008	$F_{0.1}$	111	157	212	
Riwaka	1991	$F_{0.1}$	167	224	286	-
Riwaka	2004	$F_{0.1}$	81	115	156	
Riwaka	2008	$F_{0.1}$	118	167	226	
Collingwood Beach	1993	$F_{0.1}$	20	28	37	48
Takaka Beach	1993	$F_{0.1}$	53	74	100	127
Rangihaeata Beach	1993	$F_{0.1}$	23	32	43	55

The level of risk of harvesting the populations at the estimated MCY levels cannot be determined for any of the surveyed areas. However, yield estimates are substantially higher when based on F_{MAX} than on $F_{0.1}$, so risk would be greater at MCY s based on F_{MAX} .

Estimates of CAY have been made in the past for cockle stocks at Pakawau Beach, Ferry Point and Riwaka, using $CAY = F_{REF}/(F_{REF} + M) * (1 - e^{-(F_{REF} + M)}) * B_{BEG}$ (Method 1), where beginning of season biomass (B_{BEG}) is current recruited biomass available to the fishery, and F_{REF} is either $F_{0.1}$ or F_{max} . Estimates of current biomass that allow updated calculations are available in 2008 for Pakawau Beach, Ferry Point and Tapu Bay (Riwaka). The most recent estimates of CAY available for all stocks are listed in Table 7.

4.4 Other yield estimates and stock assessment results

$F_{0.1}$ and CAY were estimated from a yield per recruit (YPR) analysis using the age and length-weight parameters for Pakawau Beach cockles from Osborne (1992), and assuming size at recruitment to the fishery of either 30 or 35 mm shell length. A range of M values was used to produce the latest estimates in Table 8 (Osborne 2014).

Table 7: Estimates of CAY (t) for various cockle stocks in Tasman and Golden Bays, assuming a range of values for M .

Site	Date	F_{REF}	M			
			0.2	0.3	0.4	0.5
Pakawau Beach	2001	$F_{0.1}$	778	996	1 210	1 396
Pakawau Beach #	2001	$F_{0.1}$	1 964	2 514	3 053	3 522
Pakawau Beach	2004	$F_{0.1}$	1 202	1 555	1 910	
Pakawau Beach	2008	$F_{0.1}$	1 161	1 501	1 845	
Pakawau Beach	2014	$F_{0.1}$	638	844	1 040	
Ferry Point	1996	$F_{0.1}$	407	501	600	696
Ferry Point	2004	$F_{0.1}$	69	89	109	
Ferry Point	2008	$F_{0.1}$	88	114	140	
Riwaka	1993	$F_{0.1}$	507	615	708	
Riwaka	2004	$F_{0.1}$	138	179	220	
Riwaka	2008	$F_{0.1}$	1 161	1 501	1 845	

Calculations using total recruited biomass, rather than available recruited biomass.

Table 8: Latest estimates of $F_{0.1}$ from a yield per recruit analysis and CAY at different levels of minimum size at harvest (MSH) and natural mortality (M) (Osborne 2014).

	MSH	B_{beg}	M		
			0.20	0.30	0.40
$F_{0.1}$	30		0.23	0.34	0.46
CAY		3363	638	844	1040
$F_{0.1}$	35		0.28	0.40	0.54
CAY		2409	541	696	838
$F_{0.1}$	37		0.31	0.43	0.56
CAY		2026	489	617	732

4.5 Other factors

The areas of Golden Bay and Tasman Bay currently commercially fished for cockles are very small with respect to the total resource. Recruitment overfishing is unlikely owing to the extent of the resource protected from the fishery in *Zostera* beds, in sub-tidal areas, and in the protected areas adjacent to Farewell Spit and in other areas of Golden Bay. Cockle larvae are planktonic for about three weeks, so areas like Golden Bay and Tasman Bay probably constitute single larval pools.

Consequently, fisheries in relatively small areas (like Pakawau Beach) are likely to have little effect on recruitment. It is noted, however, that recruitment of juvenile cockles can be reduced by the removal of a large proportion of adult cockles from the area (i.e., successful settlement occurs only in areas containing a population of adult cockles).

COCKLES (COC 7A)

It is also likely that growth and mortality of cockles are density-dependent. A reduction in density due to fishing could enhance the growth and survival of remaining cockles.


Because cockles begin to spawn at a shell length of about 18 mm, and the larval pools in Tasman and Golden Bays are probably massive and derive from a wide area (most of which is closed to commercial fishing), there is a low risk of recruitment overfishing at any of the exploited sites.

5. STATUS OF THE STOCKS

Stock structure assumptions

Little is known of the stock boundaries of cockles. Given differences in growth and mortality within and between different beds and in the absence of more detailed knowledge regarding larval connectivity, this commercial fishery area is managed as a discrete population.

COC 7A

Stock Status	
Year of Most Recent Assessment	2014
Assessment Runs Presented	Survey biomass estimates for ≥ 30 mm shell length
Reference Points	Target(s): Not defined, but B_{MSY} assumed Soft Limit: 20% B_0 Hard Limit: 10% B_0 Overfishing threshold: - Undefined
Status in relation to Target	About as Likely as Not (40–60%) to be at or above the target (except for local depletion in some bays)
Status in relation to Limits	Unlikely (< 40%) to be below the soft limit and Very Unlikely (< 10%) to be below the hard limit
Status in relation to Overfishing	Overfishing is Very Unlikely (<10%) to be occurring
Historical Stock Status Trajectory and Current Status	
 <p>Recruited biomass (≥ 30 mm shell length) over time. Notably, the area surveyed over time has changed (see Tables 4 and 5) and decreased at the last time of survey (compared to previous occasions) at all three sites.</p>	
Fishery and Stock Trends	
Recent Trend in Biomass or Proxy	The recruited biomass estimates of cockles from Pakawau beach have shown a general trend of increase until 2004, with the lowest value in 1992 (5299 t) and the highest value in 2004 (8803 t); followed by a decline to historically low levels in 2014 (3363 t). The Ferry Point recruited biomass estimates

	declined from 2442 t in 1996 to 443 t and 470 t in 2004 and 2008, respectively. Riwaka total biomass estimates decreased from 1991 (1880 t) to 2008 (939 t). Notably, the area surveyed has changed over time and decreased at the last survey (compared to previous surveys) at all three sites.
Recent Trend in Fishing Mortality or Proxy	Landings since 2004–05 are intermediate compared to the history of the fishery and have fluctuated without trend between 220 and 460 t.
Other Abundance Indices	-
Trends in Other Relevant Indicators or Variables	-

Projections and Prognosis		
Stock Projections or Prognosis	-	
Probability of Current Catch or TACC causing Biomass to remain below or to decline below Limits	Fishing at present levels is Very Unlikely (< 10%) to cause declines below the soft or hard limits.	
Probability of Current Catch or TACC causing Overfishing	Very Unlikely (< 10%)	
Assessment Methodology and Evaluation		
Assessment Type	Level 2: Partial quantitative stock assessment	
Assessment Method	Absolute biomass estimates from quadrant surveys	
Assessment Dates	Latest assessment: 2014	Next assessment: Unknown
Overall assessment quality rank	1 – High Quality	
Main data inputs (rank)	- Abundance - Length frequency	1 – High Quality 1 – High Quality
Data not used (rank)		
Changes to Model Structure and Assumptions	-	
Major Sources of Uncertainty	-	


Qualifying Comments
Water quality issues have influenced the amount of time when cockles can be harvested from Ferry Point in recent years.

Fishery Interactions
Interactions with other species are currently being characterised.

6. FOR FURTHER INFORMATION

- Belton, R J (1986) A resource survey of the *Chione stutchburyi* stocks in the Marlborough Sounds. Unpublished report prepared for M. Hones & B. Huntley. Contact Southern Clams for more information.
- Boyd, R O; Reilly, J L (2002) 1999/2000 National marine Recreational Fishing Survey: harvest estimates. New Zealand Fisheries Research Report. (Unpublished document held by Fisheries New Zealand, Wellington.)
- Bradford, E (1998) Harvest estimates from the 1996 national recreational fishing surveys. New Zealand Fisheries Assessment Research Document 1998/16. 27 p. (Unpublished document held by NIWA library, Wellington.)
- Breen, P A (1996) Yield estimates for the Tapu Bay - Riwaka fishery for cockles (*Austrovenus stutchburyi*). Unpublished report prepared for Talley's Fisheries Ltd. (NIWA Client Report WLG1996/27).
- Breen, P A (1997a) Yield calculations for cockles (*Austrovenus stutchburyi*) at Ferry Point, Golden Bay. Unpublished report prepared for Talley's Fisheries Ltd. (NIWA Client Report WLG1997/17).
- Breen, P A (1997b) An evaluation of a biomass survey for cockles on Pakawau Beach, consideration of reasonable CAY estimates and consideration of increased catch. Unpublished report prepared for Westhaven Shellfish Co. Ltd. (NIWA Client Report WLG97/32).
- Bull, M (1984) A brief study of intertidal clam beds in Western Golden Bay associated with trials of a mechanical clam harvester. Unpublished Ministry of Fisheries report. 11 p.
- Forrest, B; Asher, R (1997) Ferry Point cockle survey: revised report. Unpublished report prepared for Talley's Fisheries Ltd. (Cawthron Report No. 411).
- Grange, K (1995) Biological communities of the intertidal sandflats, Motupipi, Golden Bay. Unpublished report prepared for Sealife Investments Ltd. (NIWA Client Report 1995/NEL911/1).
- Hartill, B; Cryer, M; Morrison, M (2005) Estimates of biomass, sustainable yield, and harvest: neither necessary nor sufficient for the management of non-commercial urban intertidal shellfish fisheries. *Fisheries Research* 71: 209–222.
- Ministry of Fisheries (2010) Report from the Fisheries Assessment Plenary, May 2010: stock assessments and yield estimates. 1158 p. (Unpublished report held by Fisheries New Zealand, Wellington.)
- Osborne, T A (1992) Biomass survey and stock assessment of the New Zealand littleneck clam (*Chione stutchburyi*) on Pakawau Beach, Golden Bay. Unpublished report prepared for Westhaven Shellfish Co. Ltd. 27 p.
- Osborne, T A (1997) Biomass survey and stock assessment of the New Zealand littleneck clam (*Chione stutchburyi*) on Pakawau Beach, Golden Bay. Unpublished report prepared for Westhaven Shellfish Co. Ltd. 16 p.
- Osborne, T A (1998) 1998 biomass survey and stock assessment of the New Zealand littleneck clam (*Chione stutchburyi*) on Pakawau Beach, Golden Bay. Unpublished report prepared by Osborne Research Co. Ltd. for Westhaven Shellfish Co. Ltd. 23 p.
- Osborne, T A (1999) 1999 biomass survey and stock assessment of the New Zealand littleneck clam (*Chione stutchburyi*) on Pakawau Beach, Golden Bay. Unpublished report prepared by Osborne Research Co. Ltd. for Westhaven Shellfish Co. Ltd. 15 p.
- Osborne, T A (2001) Annual biomass survey and stock assessment of cockles on Pakawau Beach, Golden Bay - 2001. Unpublished report prepared by Osborne Research Co. Ltd. for Westhaven Shellfish Co. Ltd. 23 p.
- Osborne, T A (2004) Annual biomass survey and stock assessment of cockles on Pakawau Beach, Golden Bay - 2001. Unpublished report prepared by Osborne Research Co. Ltd. for Westhaven Shellfish Co. Ltd..
- Osborne, T A (2010) Biomass survey and stock assessment of cockles (*Austrovenus stutchburyi*) in area COC 7A:Tapu Bay, Ferry Point, and Pakawau. *New Zealand Fisheries Assessment Report 2010/44*.
- Osborne, T. (2014). Biomass survey and stock assessment of cockles (*Austrovenus stutchburyi*) at Pakawau Beach in area COC 7A 2014. An ORCO report produced for Westhaven Shellfish Company Ltd: 18 p.
- Osborne, T A; Seager, V (1994) Biomass survey and stock assessment of the New Zealand littleneck clam (*Chione stutchburyi*) on three beaches in Golden Bay. Unpublished report prepared for Wakatu Incorporation and Te Tau Ihu Ltd Joint Venture.
- Stark, J D; Asher, R A (1991) Cockle resource survey Tapu Bay - Motueka River (February 1991). Unpublished report prepared by Cawthron Institute for Talley's Fisheries Ltd.
- Teirney, L D; Kilner, A R; Millar, R E; Bell, J D (1997) Estimation of recreational catch from 1991/92 to 1993/94. New Zealand Fisheries Assessment Research Document 1997/15. 43 p. (Unpublished document held by NIWA library, Wellington.)
- Wilson, N; Stevens, J; Alspach, P; Butler, D (1988) Environmental Impact Assessment. Effects of mechanical harvesting of cockles from the Pakawau/Puoponga intertidal area. Report prepared by DSIR Division of Horticulture and Processing, Fish Technology Section, Nelson, for Westhaven Shellfish Co. Ltd.
- Wynne-Jones, J; Gray, A; Hill, L; Heinemann, A (2014) National Panel Survey of Marine Recreational Fishers 2011–12: Harvest Estimates. *New Zealand Fisheries Assessment Report 2014/67*. 139 p.

DEEPWATER (KING) CLAM (PZL)

(Panopea zelandica)

1. FISHERY SUMMARY

Deepwater clams (*Panopea zelandica*), commonly referred to as geoducs or geoducks, were introduced into the Quota Management System on 1 October 2006 with a total TAC of 40.5 t, consisting of 31.5 t TACC and a 9 t allowance for other sources of mortality (Table 1). No changes have occurred to the TAC since. The fishing year is from 1 October to 30 September and commercial catches are measured in greenweight. Deepwater clams are harvested by divers using underwater breathing apparatus and a hydraulic jet.

Table 1: Current TAC, TACC and allowances for other sources of mortality for *Panopea zelandica*.

Fishstock	TAC (t)	TACC (t)	Other sources of mortality
PZL 1	1.5	1.2	0.3
PZL 2	1.5	1.2	0.3
PZL 3	1.5	1.2	0.3
PZL 4	1.5	1.2	0.3
PZL 5	1.5	1.2	0.3
PZL 7	30.0	23.1	6.9
PZL 8	1.5	1.2	0.3
PZL 9	1.5	1.2	0.3
Total	40.5	31.5	9.0

1.1 Commercial fisheries

The largest landings since 1989 were reported between 1989 and 1992 (Table 2), almost all taken in the Nelson-Marlborough region under a special permit for investigative research. Targeted fishing was also carried out under a special permit in PZL 7 between 2004 and 2005. Rare catches have also been made by trawlers. The largest catch since 1993 (10.885 t) occurred in 2011–12 and was mainly taken from the Nelson-Marlborough region (Table 2).

1.2 Recreational fisheries

There are no estimates of recreational take for this surf clam. Recreational take is likely to be very small or non-existent.

1.3 Customary fisheries

This clam is harvested for customary use when washed ashore after storms but there are no estimates of this use of this clam. Customary take is likely to be very small or non-existent.

DEEPWATER (KING) CLAM (PZL)

Table 2: TACCs and reported landings (t) of deepwater clam by Fishstock from 1988–89 to present, taken from CELR and CLR data. There have never been any reported landings in PZL 2, 4, 5, 8, or 9.

Fishstock	PZL 1		PZL 3		PZL 7		Total	
	Landings	TACC	Landings	TACC	Landings	TACC	Landings	TACC
1989–90	0.315	-	0	-	95.232	-	95.547	-
1990–91	0	-	0	-	29.293	-	29.293	-
1991–92	0	-	0.725	-	31.394	-	32.119	-
1992–93	0	-	0.053	-	0	-	0.053	-
1993–94	0	-	0	-	0	-	0	-
1994–95	0	-	0	-	0	-	0	-
1995–96	0	-	0	-	0	-	0	-
1996–97	0	-	0	-	0	-	0	-
1997–98	0	-	0	-	0	-	0	-
1998–99	0	-	0	-	0	-	0	-
1999–00	0	-	0	-	0	-	0	-
2000–01	0	-	0.146	-	0	-	0.146	-
2001–02	0.003	-	0.068	-	0	-	0.071	-
2002–03	0	-	0.001	-	0	-	0.001	-
2003–04	0	-	0	-	1.444	-	1.444	-
2004–05	0	-	0	-	2.944	-	2.944	-
2005–06	0	-	0	-	0	-	0	-
2006–07	0	1.2	0	1.2	0	23.1	0	31.5
2007–08	0	1.2	0.132	1.2	0.320	23.1	0.450	31.5
2008–09	0	1.2	0.016	1.2	5.100	23.1	5.116	31.5
2009–10	0	1.2	0	1.2	4.578	23.1	4.578	31.5
2010–11	0	1.2	0.076	1.2	7.880	23.1	7.956	31.5
2011–12	0	1.2	0.036	1.2	10.849	23.1	10.885	31.5
2012–13	0	1.2	0	1.2	1.746	23.1	1.746	31.5
2013–14	0	1.2	0	1.2	6.072	23.1	6.072	31.5
2014–15	0	1.2	0.003	1.2	3.927	23.1	3.93	31.5
2015–16	0	1.2	0	1.2	4.686	23.1	4.686	31.5
2016–17	0	1.2	0	1.2	3.260	23.1	3.260	31.5

1.4 Illegal catch

There is no documented illegal catch of this clam.

1.5 Other sources of mortality

There is little information on other sources of mortality, although the clam has on rare occasions been captured during trawling operations. Adults show poor reburial after being dug out (Gribben & Creese 2005).

2. BIOLOGY

There are two similar *Panopea* species in New Zealand, *P. zelandica* and *P. smithae*, both of which are endemic and occur around the North, South and Stewart Islands. *P. smithae* has also been reported from the Chatham Islands. Their distributions overlap, but *P. zelandica* occurs mainly in shallow waters (5–25 m) in sand and mud off sandy ocean beaches, while *P. smithae* lives mainly at greater depths (110–130 m) on coarse shell bottoms, and is also thought to burrow deeper in the substrate. In samples of commercial and exploratory catches, *P. zelandica* is more abundant than *P. smithae*, and in the early 1990s it comprised virtually all of the catch.

Deepwater clams are broadcast spawners with separate sexes. Protandric development (where an organism begins life as a male and then becomes a female) is considered likely for a proportion of the population (Gribben & Creese 2003). Fifty percent sexual maturity was calculated at 55 and 57 mm length for populations in Wellington and on the Coromandel Peninsula, respectively. Samples taken from three locations between the Coromandel Peninsula and Nelson showed spawning between spring and late summer (Gribben et al 2004). Spawning may be temperature controlled because it occurred at the Coromandel and Wellington sites when water temperature reached approximately 15°C (Gribben et al 2004). The larval life is thought to be about two to three weeks (Gribben & Hay 2003), and there is evidence of significant recruitment variation between years.

The oldest *P. zelandica* based on annual ring counts in Golden Bay, Shelly Bay and Kennedy Bay were 34, 34 and 85 years respectively (Breen 1991, Gribben & Creese 2005); ring counts were validated from Shelly Bay only. Growth in shell length appeared to be rapid for the first 10–12 years in these populations and total weight increased rapidly until at least 12–13 years of age. Differences in growth

rates were seen between the Kennedy and Shelly Bay populations: estimates of K varied between 0.16 and 0.29, t_0 between 1.67 and 3.8 and L_∞ between 103.6 and 116.5 mm, respectively (Breen 1991, Gribben & Creese 2005)¹.

Estimates of M , instantaneous natural mortality, from catch curve analysis, estimates of maximum age, and the Chapman-Robson estimator from Kennedy Bay and Shelly Bay populations were all between 0.02 and 0.12 (Gribben & Creese 2005). The estimate by Breen (1991) for Golden Bay was 0.15, but in modeling this parameter was varied from 0.1 to 0.2.

3. STOCKS AND AREAS

For management purposes stock boundaries are based on FMAs, however, there is little information on stock structure, recruitment patterns, or other biological characteristics to determine fishstock boundaries.

4. STOCK ASSESSMENT

No stock assessments have been carried out for any deepwater clam stocks. Sustainable fishing rate estimates were made by Breen (1994).

4.1 Estimates of fishery parameters and abundance

No abundance estimates are available for any geoduc stocks. Sustainable fishing rate estimates were made by Breen (1994).

4.2 Biomass estimates

Biomass has not been estimated for any deepwater clam stocks.

4.3 Yield estimates and projections

MCY has not been estimated for any deepwater clam stocks. However, an age-structured stochastic model suggested that sustainable yields for this species, with realistic management constraints, appear to be on the order of 2% to 4% of virgin biomass (Breen 1994).

CAY has not been estimated for any deepwater clam stocks.

5. STATUS OF THE STOCKS

PZL 7 - *Panopea zelandica*

Stock Status	
Year of Most Recent Assessment	No formal assessment done for any stock
Assessment Runs Presented	-
Reference Points	Target: Not defined, but B_{MSY} assumed Soft Limit: 20% B_0 Hard Limit: 10% B_0 Overfishing threshold: -
Status in relation to Target	Because of the relatively low levels of exploitation of <i>P. zelandica</i> , it is likely that this stocks is still effectively in a virgin state, therefore it is Very Likely (> 60%) to be at or above the target.
Status in relation to Limits	Very Unlikely (< 40%) to be below the soft or hard limit
Historical Stock Status Trajectory and Current Status	-

¹ No confidence intervals were available for these estimates.

DEEPWATER (KING) CLAM (PZL)

Fishery and Stock Trends	
Recent Trend in Biomass or Proxy	Unknown
Recent Trend in Fishing Mortality or Proxy	In 1989–92 the landings for PZL 7 averaged 52 t; however, since that time fishing has been light in all QMAs with a maximum of only 10.9 t taken across all QMAs in the 2011–12 fishing year.
Other Abundance Indices	-
Trends in Other Relevant Indicators or Variables	-

Projections and Prognosis	
Stock Projections or Prognosis	-
Probability of Current Catch or TACC causing Biomass to remain below or to decline below Limits	Current catches are Unlikely (< 40%) to cause declines below soft or hard limits.
Probability of Current Catch causing Overfishing to continue or to commence	-

Assessment Methodology and Evaluation		
Assessment Type	-	
Assessment Method	-	
Assessment Dates	Latest assessment: -	Next assessment: -
Overall assessment quality rank	-	
Main data inputs (rank)		
Data not used (rank)		
Changes to Model Structure and Assumptions	-	
Major Sources of Uncertainty	-	

Qualifying Comments
Early surveys show that density is generally low compared with North American species but that productivity is higher.

Fishery Interactions
Interactions with other species are currently being characterised.

7. FOR FURTHER INFORMATION

- Beentjes, M P; Baird, S J (2004) Review of dredge fishing technologies and practice for application in New Zealand. *New Zealand Fisheries Assessment Report 2004/37*. 40 p.
- Breen, P A (1991) The New Zealand deepwater clams (geoducs), *Panopea zelandica* and *P. smithae*. New Zealand Fisheries Assessment Research Document 1991/5. 12 p. (Unpublished report held by NIWA library, Wellington.)
- Breen, P A (1994) Sustainable fishing patters for geoduc clam (*Panopea zelandica*) populations in New Zealand. New Zealand Fisheries Assessment Research Document 1994/4. 34 p. (Unpublished report held by NIWA library, Wellington.)
- Breen, P A; Gabriel, C; Tyson, T. (1991) Preliminary estimates of age, mortality, growth, and reproduction in the hiatellid clam *Panopea zelandica* in New Zealand. *New Zealand Journal of Marine and Freshwater Research* 25: 231–237.
- Cranfield, H J; Michael, K P; Stotter, D R (1993) Estimates of growth, mortality, and yield per recruit for New Zealand surf clams. New Zealand Fisheries Assessment Research Document 1993/20. 46 p. (Unpublished report held by NIWA library, Wellington.)
- Cranfield, H J; Michael, K P; Stotter, D; Doonan, I J (1994) Distribution, biomass and yield estimates of surf clams off New Zealand beaches. New Zealand Fisheries Assessment Research Document 1994/1. 27 p. (Unpublished report held by NIWA library, Wellington.)
- Cranfield, H J; Michael, K P (2001) The surf clam fishery in New Zealand: description of the fishery, its management, and the biology of surf clams. *New Zealand Fisheries Assessment Report 2001/62*. 24 p.
- Gribben, P E; Creese, R G (2003) Protandry in the New Zealand geoduck, *Panopea zelandica* (Mollusca, Bivalvia). *Invertebrate Reproduction & Development* 44(2–3): 119–129.
- Gribben, P E; Creese, R G (2005) Age, growth, and mortality of the New Zealand geoduck clam, *Panopea zelandica* (Bivalvia : Hiatellidae) in two north island populations. *Bulletin of Marine Science* 77(1): 119–135.
- Gribben, P E; Hay, B E (2003) Larval development of the New Zealand geoduck *panopea zelandica* (Bivalvia : Hiatellidae). *New Zealand Journal of Marine and Freshwater Research* 37(2): 231–239.


DEEPWATER (KING) CLAM (PZL)

- Gribben, P E; Helson, J; Jeffs, A (2004) Reproductive cycle of the New Zealand geoduck, *Panopea zelandica*, in two north island populations. *Veliger* 47(1): 53–65.
- Morton, J; Miller, M (1968) *The New Zealand sea shore*. Collins, Auckland. 638 p.
- Powell, A W B (1979) *New Zealand Mollusca. Marine, land and freshwater shells*. Collins, Auckland. 500 p.

ELEPHANT FISH (ELE)

(Callorhinchus milii)

Reperepe



- Commercial fisheries

From the 1950s to the 1980s, landings of elephant fish of around 1000 t/year were common. Most of these landings were from the area now encompassed by ELE 3, but fisheries for elephant fish also developed on the south and west coasts of the South Island in the late 1950s and early 1960s, with average catches of around 70 t per year in the south (in the 1960s to the early 1980s) and 10–30 t per year on the west coast. Total annual landings of elephant fish dropped considerably in the early 1980s (between 1982–83 and 1994–96 they ranged between 500 and 700 t) but later increased to the point that they have annually exceeded 1 000 t since the 1995–96 fishing season. Reported landings since 1931 are shown in Tables 1 and 2, while an historical record of landings and TACC values for the three main ELE stocks are depicted in Figure 1. ELE 3 has customary, recreational and other mortality allowances of 5 t, 5 t, and 50 t respectively, and ELE 5 has allowances 5 t, 5 t, and 7 t respectively.

Table 1: Reported total landings of elephant fish for calendar years 1936 to 1982. Sources: MAF and FSU data.

Year	Landings (t)	Year	Landings (t)	Year	Landings (t)	Year	Landings (t)	Year	Landings (t)
1936	116	1946	235	1956	980	1966	1 112	1976	705
1937	184	1947	188	1957	1 069	1967	934	1977	704
1938	201	1948	230	1958	1 238	1968	862	1978	596
1939	193	1949	310	1959	1 148	1969	934	1979	719
1940	259	1950	550	1960	1 163	1970	1 128	1980	906
1941	222	1951	602	1961	983	1971	1 401	1981	690
1942	171	1952	459	1962	1 156	1972	1 019	1982	661
1943	220	1953	530	1963	1 095	1973	957		
1944	270	1954	853	1964	1 235	1974	848		
1945	217	1955	802	1965	1 111	1975	602		

The TACC for ELE 3 has, with the exception of 2002–03, been consistently exceeded since 1986–87. The ELE 3 TACC was consequently increased to 500 t for the 1995–96 fishing year, and then increased twice more under an Adaptive Management Programme (AMP): initially to 825 t in October 2000 and then to 950 t in October 2002. This new TACC combined with the allowances for customary and recreational fisheries (5 t each), increased the new TAC for the 2002–03 fishing year in ELE 3 to 960 t. For the 2009–10 fishing year, the TACC was increased from 960 t to 1 000 t where it presently remains. ELE 3 fishing is seasonal, mostly occurring in spring and summer in inshore waters. Most of the increase in catch from the early 2000s in the ELE 3 trawl fishery has been taken as a bycatch of the flatfish target fishery and an emerging target

ELEPHANT FISH (ELE)

ELE fishery (Starr & Kendrick 2013). During the 1990s, the level of elephant fish bycatch from the RCO 3 trawl fishery increased from around 80 t/year to greater than 400 t in 2000–01 (Starr & Kendrick 2013). There was a steady increase in the level of ELE 3 bycatch from the FLA 3 trawl fishery, with catches increasing from around 70 t in 1994–95 to 300 t in 1999–00. There is also a significant setnet fishery in ELE 3, largely directed at rig and elephant fish.

The fishery in ELE 5 is mainly a trawl fishery targeted at flatfish and to a lesser extent giant stargazer. Very little catch in ELE 5 is taken by target setnet fisheries. Catches have been increasing consistently since 1992–93, exceeding the TACCs since 1995–96. The ELE 5 TACC was increased from 71 t to 100 t under an AMP in October 2001. The TACC was further increased under the AMP to 120 t in October 2004 and catches have exceeded this TACC by 70% in 2007–08 and 2008–09. For the 2009–10 fishing season, the TACC was increased by 17% up from 120 t to 140 t. All AMP programmes ended on 30 September 2009. The ELE 5 TACC was further increased to 170 t in 2012–13.

From 1 October 2008, a suite of regulations intended to protect Maui's and Hector's dolphins was implemented for all of New Zealand by the Minister of Fisheries. For ELE 3, commercial and recreational set netting was banned in most areas to 4 nautical miles offshore of the east coast of the South Island, extending from Cape Jackson in the Marlborough Sounds to Slope Point in the Catlins. Some exceptions were allowed, including an exemption for commercial and recreational set netting to only one nautical mile offshore around the Kaikoura Canyon, and permitting setnetting in most harbours, estuaries, river mouths, lagoons and inlets except for the Avon-Heathcote Estuary, Lyttelton Harbour, Akaroa Harbour and Timaru Harbour. In addition, trawl gear within 2 nautical miles of shore was restricted to flatfish nets with defined low headline heights. For ELE 7, both commercial and recreational setnetting were banned to 2 nautical miles offshore, with the recreational closure effective for the entire year and the commercial closure restricted to the period 1 December to the end of February. The closed area extends from Awarua Point north of Fiordland to the tip of Cape Farewell at the top of the South Island. Some interim relief to these regulations was provided in ELE 5 from 1 October 2008 to 24 December 2009.

Table 2: Reported landings (t) for the main QMAs from 1931 to 1990.

Year	ELE 1	ELE 2	ELE 3	ELE 5	ELE 7
1931–32	0	0	0	0	0
1932–33	0	0	0	0	0
1933–34	0	0	0	0	0
1934–35	0	0	0	0	0
1935–36	0	0	0	0	0
1936–37	0	0	79	0	1
1937–38	0	0	183	0	0
1938–39	0	0	194	1	2
1939–40	0	1	190	1	1
1940–41	0	1	243	8	1
1941–42	0	0	220	1	0
1942–43	0	0	163	6	0
1943–44	0	0	219	1	0
1944	0	0	251	10	0
1945	0	2	205	3	3
1946	0	0	228	3	4
1947	0	2	176	0	10
1948	0	2	227	0	9
1949	0	1	296	2	13
1950	0	1	522	14	13
1951	0	2	585	6	10
1952	0	0	440	9	5
1953	0	3	514	13	3
1954	0	2	839	5	7
1955	0	3	771	4	25
1956	0	1	933	16	29
1957	0	2	992	28	46
1958	0	0	1 140	47	51
1959	0	0	1 066	37	44
1960	0	1	1 099	38	27
1961	0	0	913	43	27
1962	0	4	1 066	73	14
1963	0	2	976	111	8
1964	0	3	1 109	107	16
1965	0	7	983	88	34
1966	0	1	985	99	27
1967	0	1	812	77	45

ELEPHANT FISH (ELE)

Table 2: [Continued]

1968	0	1	757	54	52
1969	0	1	824	75	33
1970	0	3	987	87	53
1971	0	0	1243	103	37
1972	0	0	928	70	15
1973	0	0	864	73	21
1974	0	0	766	97	41
1975	0	1	557	55	28
1976	0	0	622	91	52
1977	0	0	601	114	45
1978	0	0	552	49	26
1979	0	0	661	63	18
1980	0	0	794	129	34
1981	0	1	543	114	16
1982	0	0	584	85	34

Notes:

1. The 1931–1943 years are April–March but from 1944 onwards are calendar years.
2. Data up to 1985 are from fishing returns: Data from 1986 to 1990 are from Quota Management Reports.
3. Data for the period 1931 to 1982 are based on reported landings by harbour and are likely to be underestimated as a result of under-reporting and discarding practices. Data includes both foreign and domestic landings. Data were aggregated to FMA using methods and assumptions described by Francis & Paul (2013).

Table 3: Reported landings (t) of elephant fish by Fishstock from 1983–84 to 2016–17 and actual TACCs (t) from 1986–87 to 2016–17. QMR data from 1986 – present. No landings have been reported from ELE 10.

Fishstock	ELE 1		ELE 2		ELE 3		ELE 5		ELE 7			Total
FMA (s)	1 & 9		2 & 8		3 & 4		5 & 6		7			
	Landings	TACC	Landings	TACC	Landings	TACC	Landings	TACC	Landings	TACC	Landings	
1983–84*	< 1	-	5	-	605	-	94	-	60	-	765	-
1984–85*	< 1	-	3	-	517	-	134	-	50	-	704	-
1985–86*	< 1	-	4	-	574	-	57	-	46	-	681	-
1986–87	< 1	10	2	20	506	280	48	60	29	90	584	470
1987–88	< 1	10	3	20	499	280	64	60	44	90	610	470
1988–89	< 1	10	1	22	450	415	49	62	43	100	543	619
1989–90	< 1	10	3	22	422	418	32	62	55	101	510	623
1990–91	< 1	10	5	22	434	422	55	71	59	101	553	636
1991–92	< 1	10	11	22	450	422	58	71	78	101	597	636
1992–93	< 1	10	5	22	501	423	39	71	61	102	606	638
1993–94	< 1	10	6	22	475	424	46	71	41	102	568	639
1994–95	< 1	10	5	22	580	424	60	71	39	102	684	639
1995–96	< 1	10	7	22	688	500	72	71	93	102	862	715
1996–97	< 1	10	9	22	734	500	74	71	94	102	912	715
1997–98	< 1	10	12	22	910	500	95	71	66	102	1 082	715
1998–99	< 1	10	9	22	842	500	129	71	117	102	1 098	715
1999–00	< 1	10	6	22	950	500	105	71	87	102	1 148	715
2000–01	2	10	7	22	956	825	153	71	90	102	1 207	1 040
2001–02	< 1	10	9	22	852	825	105	100	88	102	1 053	1 057
2002–03	1	10	9	22	950	950	106	100	59	102	1 125	1 194
2003–04	< 1	10	10	22	984	950	102	100	42	102	1 139	1 194
2004–05	< 1	10	13	22	972	950	125	120	74	102	1 184	1 214
2005–06	< 1	10	14	22	1 023	950	147	120	76	102	1 260	1 214
2006–07	< 1	10	17	22	960	950	158	120	116	102	1 251	1 214
2007–08	< 1	10	16	22	1 092	950	202	120	125	102	1 435	1 214
2008–09	1	10	21	22	1 063	950	208	120	91	102	1 384	1 214
2009–10	< 1	10	21	22	1 089	1 000	176	140	86	102	1 372	1 274
2010–11	< 1	10	14	22	1 123	1 000	153	140	93	102	1 384	1 283
2011–12	< 1	10	16	22	1 074	1 000	157	140	130	102	1 377	1 283
2012–13	< 1	10	16	22	1 140	1 000	157	170	123	102	1 436	1 304
2013–14	< 1	10	16	22	1 110	1 000	173	170	96	102	1 394	1 304
2014–15	< 1	10	11	22	1 048	1 000	179	170	102	102	1 340	1 304
2015–16	< 1	10	9	22	1 159	1 000	137	170	95	102	1 400	1 304
2016–17	< 1	10	12	22	1 051	1 000	182	170	81	102	1 326	1 304

ELEPHANT FISH (ELE)




Figure 1: Reported commercial landings and TACC for the three main ELE stocks. From top: ELE 3 (South East Coast and Chatham Rise), ELE 5 (Southland and Sub-Antarctic), and ELE 7 (Challenger).

- **Recreational fisheries**

Catches of elephant fish by recreational fishers are low compared with those of the commercial sector. The National Panel Survey (NPS) in 2011–12 (Wynne-Jones et al 2014) generated estimates of recreational harvest of 4853 fish from ELE 3, 960 fish from ELE 7, and about 200 fish from each of ELE 2 and ELE 5. These estimates are quite uncertain; the CV on the national harvest of 6198 fish was 34%. Regional surveys in the early 1990s (Teirney et al 1997) and national surveys in 1996, 1999, and 2000 (Bradford 1998, Boyd & Reilly 2002) showed similarly low number of fish harvested and similar geographical patterns. No estimates of mean weight are available to convert these estimates of harvested fish to harvested weights.

- **Customary non-commercial fisheries**

Quantitative information on the current level of customary non-commercial catch is not available.

- **Illegal catch**

There are reports of discards of juvenile elephant fish by trawlers from some areas. However, no quantitative estimates of discards are available.

- **Other sources of mortality**

The significance of other sources of mortality has not been documented.

2. BIOLOGY

Elephant fish are uncommon off the North Island and occur south of East Cape on the east coast and south of Kaipara on the west coast. They are most plentiful around the east coast of the South Island.

Males mature at a length of 50 cm fork length (FL) at an age of 3 years, females at 70 cm FL at 4 to 5 years of age. The maximum age of elephant fish is unknown. However a tagged, 73 cm total length, Australian male was at liberty for 16 years, suggesting a longevity for males of at least 20 years (Coutin 1992, Francis 1997). Females probably also live to at least 20 years. A longevity of 20 years suggests that M is about 0.23. This results from use of the equation $M = \log_e 100/\text{maximum age}$, where maximum age is the age to which 1% of the population survives in an unexploited stock.

Mature elephant fish migrate to shallow inshore waters in spring and aggregate for mating. Eggs are laid on sand or mud bottoms, often in very shallow areas. They are laid in pairs in large yellow-brown egg cases. The period of incubation is at least 5–8 months, and juveniles hatch at a length of about 10 cm FL. Females are known to spawn multiple times per season. After egg laying the adults are thought to disperse and are difficult to catch; however, juveniles remain in shallow waters for up to 3 years. During this time juveniles are vulnerable to incidental trawl capture, but are of little commercial value.

Von Bertalanffy growth curves based on MULTIFAN analysis of length-frequency data are available for Pegasus Bay and Canterbury Bight in 1966–68 and 1983–88. However, the ages of the larger fish were probably underestimated and the growth curves are only reliable to about 4–5 years (Francis 1997). New empirical growth curves were developed by fitting a Von Bertalanffy growth function to a dataset consisting of (a) the first six length-frequency modes from the study by Francis (1997) and (b) an approximate maximum size and age for male and female elephant fish. The latter points ‘anchor’ the curves at the right hand ends and generate more plausible curve shapes, L_∞ estimates, and therefore length-at-age. The largest measured fish in the ELE 3 samples from 1966–68 and 1983–88 (i.e. 76 cm FL for males and 97 cm FL for females) were considered to be reasonable estimates of the mean maximum lengths of elephant fish in an unfished population. The following data points were therefore used in fitting the growth curves: 76 cm and 20 years for males, and 97 cm and 20 years for females. The best fitting growth model had separate male and female coefficients for K and L_∞ and a common coefficient for t_0 (M. Francis, unpubl. data).

Biological parameters relevant to the stock assessment are shown in Table 4.

ELEPHANT FISH (ELE)

Table 4: Estimates of biological parameters for elephant fish.

Fishstock	Estimate		Source
<u>1. Natural mortality (<i>M</i>)</u>			
All	0.23		See text
<u>2. Weight = a (length)^b (Weight in g, length in cm fork length)</u>			
	Both sexes		
	a	b	
ELE 3	0.0091	3.02	Gorman (1963)
<u>3. von Bertalanffy Growth Function</u>			
	Females		
	<i>L</i> _∞	<i>k</i>	<i>t</i> ₀
ELE 3	97.88	0.26	-0.55
	Males		
	<i>L</i> _∞	<i>k</i>	<i>t</i> ₀
ELE 3	75.03	0.34	-0.55
	See text		

3. STOCKS AND AREAS

There are no data that would alter the current stock boundaries. Results from tagging studies conducted during 1966–69 indicate that elephant fish tagged in the Canterbury Bight remained in ELE 3. Separate spawning grounds to maintain each ‘stock’ have not been identified. The boundaries used are related to the historical fishing pattern when this was a target fishery.

4. STOCK ASSESSMENT

Estimates of fishery parameters and abundance

1.1.1 Trawl survey biomass indices

ECSI Trawl Survey

The ECSI winter surveys from 1991 to 1996 in 30–400 m were replaced by summer trawl surveys (1996–97 to 2000–01) which also included the 10–30 m depth range, but these were discontinued after the fifth in the annual time series because of the extreme fluctuations in catchability between surveys (Francis et al 2001). The winter surveys were reinstated in 2007 and this time included additional 10–30 m strata in an attempt to index elephant fish and red gurnard which were included in the target species. Only the 2007, 2012, 2014 and 2016 surveys provide full coverage of the 10–30 m depth range (Figure 2).

Total biomass in the core strata increased markedly in 1996 and although it has fluctuated since then it has remained high with the post-1994 average (including 2014) about three-fold greater than that of the early 1990s (Figure 2). The 2016 biomass was more than six-fold greater than this average, but the CV around the estimate was 68%, very high compared to previous surveys. In the core plus shallow strata, biomass increased markedly in 2016, but the CV was also very high. The additional elephant fish biomass captured in the 10–30 m depth range accounted for 44%, 64%, 41% and 7% of the biomass in the core plus shallow strata (10–400 m) for 2007, 2012, 2014 and 2016 respectively, indicating the importance of shallow strata for elephant fish biomass (Table 5, Figure 2). Further, the addition of the 10–30 m depth range had a significant effect on the shape of the length frequency distributions with the appearance of strong 1+ and 2+ cohorts, otherwise poorly represented in the core strata, particularly in 2007 and 2012. The proportion of pre-recruit biomass in the core plus shallow strata was also greater than that of the core strata alone, indicating that younger fish are more common in shallow water (Table 5). For the four core plus shallow strata surveys the juvenile biomass (based on the length-at-50% maturity) has varied from about one third to three quarters of the total biomass in the first three surveys, but was much lower in 2016 at 9%. The distribution of elephant fish hot spots varies, but overall this species is consistently well represented over the entire survey area from 10 to 100 m, but is most abundant in the shallow 10 to 30 m.

WCSI Trawl Survey

For WCSI Trawl Surveys, elephant fish (ELE 7) total biomass estimates are variable between successive surveys and the biomass estimates are frequently imprecise, particularly for the higher biomass estimates (Table 5). The last three trawl surveys (2009, 2011 and 2013) have estimated relatively high levels of recruited biomass compared to the biomass estimates from the earlier surveys (Figure 3). However, of the three recent surveys, only the 2013 survey provided a biomass estimate with a reasonable level of precision (CV 26%). The survey estimates of pre-recruit biomass are also poorly determined.




Figure 2: Elephant fish total biomass and 95% confidence intervals for all ECSI winter surveys in core strata (30–400 m), and core plus shallow strata (10–400 m) in 2007, 2012, 2014 and 2016.




Figure 3: Elephant fish trawl survey pre-recruit and recruited biomass estimates for the west coast South Island area of the WCSI trawl survey, with associated confidence intervals. Recruited fish were defined as fish above 40 cm F.L.

ELEPHANT FISH (ELE)

Table 5: Relative biomass indices (t) and coefficients of variation (CV) for elephant fish for east coast South Island (ECSI) - summer and winter, west coast South Island (WCSI) and the Stewart-Snares Island survey areas*. Biomass estimates for ECSI in 1991 have been adjusted to allow for non-sampled strata (7 and 9 equivalent to current strata 13, 16 and 17). The sum of pre-recruit and recruited biomass values do not always match the total biomass for the earlier surveys because at several stations length frequencies were not measured, affecting the biomass calculations for length intervals. – , not measured; NA, not applicable. Recruited is defined as the size-at-recruitment to the fishery (50 cm).

Region	Fishstock	Year	Trip number	Total Biomass estimate	CV (%)	Total Biomass estimate	CV (%)	Pre- recruit	CV (%)	Pre- recruit	CV (%)	Recruite d	CV (%)	Recruite d	CV (%)
ECSI(winter)	ELE 3			30–400 m		10–400 m		30–400 m		10–400 m		30–400 m		10–400 m	
		1991	KAH9105	300	40	-	-	NA	NA	-	-	NA	NA	-	-
		1992	KAH9205	176	32	-	-	54	83	-	-	122	28	-	-
		1993	KAH9306	481	33	-	-	60	56	-	-	421	34	-	-
		1994	KAH9406	152	33	-	-	22	51	-	-	142	34	-	-
		1996	KAH9606	858	30	-	-	338	40	-	-	520	26	-	-
		2007	KAH0705	1 034	32	1 859	24	516	59	1 201	36	518	21	658	20
		2008	KAH0806	1404	35	-	-	627	57	-	-	777	27	-	-
		2009	KAH0905	596	23	-	-	210	38	-	-	387	25	-	-
		2012	KAH1207	1 351	39	3 781	31	66	46	581	25	1 285	39	3 199	36
		2014	KAH1402	951	34	1600	21	174	32	429	25	777	40	1 171	28
		2016	KAH1605	6 812	68	7 299	63	62	43	167	30	6 750	68	7 132	64
ECSI(summer)	ELE 3	1996–97	KAH9618	1 127	31	-	-	-	-	-	-	-	-	-	-
		1997–98	KAH9704	404	18	-	-	-	-	-	-	-	-	-	-
		1998–99	KAH9809	1 718	28	-	-	-	-	-	-	-	-	-	-
		1999–00	KAH9917	1 097	25	-	-	-	-	-	-	-	-	-	-
		2000–01	KAH0014	693	18	-	-	-	-	-	-	-	-	-	-
WCSI	ELE 7	1992	KAH9204	38	42	-	-	-	-	-	-	-	-	-	-
		1994	KAH9404	167	33	-	-	-	-	-	-	-	-	-	-
		1995	KAH9504	85	35	-	-	-	-	-	-	-	-	-	-
		1997	KAH9701	94	33	-	-	-	-	-	-	-	-	-	-
		2000	KAH0004	42	63	-	-	-	-	-	-	-	-	-	-
		2003	KAH0304	49	34	-	-	-	-	-	-	-	-	-	-
		2005	KAH0503	59	33	-	-	-	-	-	-	-	-	-	-
		2007	KAH0704	28	53	-	-	-	-	-	-	-	-	-	-
		2009	KAH0904	185	83	-	-	-	-	-	-	-	-	-	-
		2011	KAH1104	170	53	-	-	-	-	-	-	-	-	-	-
		2013	KAH1305	110	26										
Stewart-Snares	ELE 5	1993	TAN9301	219	33	-	-	-	-	-	-	-	-	-	-
		1994	TAN9402	177	47	-	-	-	-	-	-	-	-	-	-
		1995	TAN9502	69	49	-	-	-	-	-	-	-	-	-	-
		1996	TAN9604	137	46	-	-	-	-	-	-	-	-	-	-

*Assuming area availability, vertical availability and vulnerability equal 1.0. Biomass is only estimated outside 10 m depth except for COM9901 and CMP0001. Note: because trawl survey biomass estimates are indices, comparisons between different seasons (e.g., summer and winter ECSI) are not strictly valid.

1.1.2 CPUE biomass indices

ELE 3 and ELE 5

Three standardised CPUE series for ELE 3 were prepared for 2012, with each series based on the bycatch of elephant fish in bottom trawl fisheries defined by different target species combinations. Initially, the Working Group accepted a series based solely on the bycatch of elephant fish when targeting red cod. It then requested two further analyses: one [ELE 3(MIX)] where the target species definition was expanded to include STA, BAR, TAR, and ELE, as well as RCO, to investigate the effect of target species switching by explicitly standardising for target species effects. The second analysis [ELE 3(MIX)-trip] was done on all trips that targeted RCO, STA, BAR, TAR, and ELE at least once, then amalgamating all data to the level of a trip. This removed the differences between the TCEPR, TCER and CELR forms, but loses all targeting information.

The three sets of ELE 3 CPUE indices (ELE 3(RCO), ELE 3(MIX) and ELE 3(MIX)-trip) were very similar for the 1989–90 to 2010–11 years. The Working Group agreed in 2009 to drop the ELE 3-SN(SHK) and ELE 5-SN(SHK) (setnet with shark target species) indices because the setnet fisheries in these two QMAs have been substantially affected by management interventions (including measures to reduce the bycatch of Hector’s dolphins) and no longer appeared to be an appropriate index of ELE abundance in either QMA.

In 2014, the ELE 3(MIX) CPUE model was updated to include additional data from 2011–12 and 2012–13 (Langley 2014). The resulting CPUE indices were very similar to the previous analysis for the comparable period. The indices were updated again in 2016, extending the time-series to 2014–15. Standardised CPUE has fluctuated without trend since 2009–10 and the 2014–15 data point is near the interim target (see below) (Figure 4).

An analysis of recent CPUE data suggested that bottom trawl fishing operations may be attempting to avoid larger catches of elephant fish. During 2012–13 to 2014–15, there was a lower probability of successive larger catches of elephant fish. This may have negatively biased the CPUE indices from 2012–13 to 2014–15 (Langley 2016 - presentation).

B_{MSY} conceptual proxy: The Working Group proposed using the average of the ELE 3(MIX) series from 1998–99 to 2010–11 to represent a “ B_{MSY} conceptual proxy” for the ELE 3 Fishstock. This period was selected because of its relative stability following a period of continuous increase. However, the Working Group has concerns about the reliability of this as a proxy and suggested that it only be used on an interim basis.




Figure 4: Standardised CPUE indices for the ELE 3 bottom trawl fisheries [ELE 3(MIX)]. The horizontal grey line is the mean of ELE 3(MIX) from 1998–99 to 2010–11 (B_{MSY} conceptual proxy). The CPUE series has been normalised to a geometric mean of 1.0. Error bars show 95% confidence intervals.

ELEPHANT FISH (ELE)

Two standardised CPUE series for ELE 5 were prepared for 2012 with each series based on the bycatch of elephant fish in the bottom trawl fisheries defined by target species combinations (Starr & Kendrick 2013). One of these series [ELE 5 BT(MIX)] is analogous to the MIX series developed for ELE 3, with the series defined by six target species in all valid ELE 5 statistical areas. The second ELE 5 analysis [ELE 5 BT(MIX)-trip] was a trip- based analysis using the same target species selection method as described for ELE 3-BT(MIX)-trip series. The two sets of indices were very similar.

In 2014, the ELE 5-BT(MIX) CPUE model was updated to include data from 2011–12 to 2012–13 (Langley 2014). This model used the “daily effort” method to prepare the data, whereby every record was reduced to a day of fishing, with the predominant statistical area and target species for the day assigned to the record. This method was accepted by the WG as the best procedure to follow when reducing event-based forms to match earlier daily forms. The two most recent indices were lower than the peak CPUE from 2008–09 to 2010–11, although CPUE has been maintained at a relatively high level compared to the 1990s–early 2000s (Figure 5). The ELE 5-BT(MIX) model was again updated in 2017, with data current to the end of 2015–16. Although the fishery definition and data preparation methods were unchanged, a binomial presence/absence series was added because of a declining trend in the proportion of days with zero catch. The Plenary accepted a revised index which combined the binomial and lognormal series using the delta-lognormal method (Starr & Kendrick, in prep). This was done because the Inshore WGs have adopted the standard of combining positive catch and fishing success models when there is a trend in the proportion zero catch. As well, simulation work has indicated that calculating a combined index may reduce bias when reporting small catch amounts (Langley 2015). Recent indices estimated by this updated series are lower than the peak observed at the end of the 2010 decade, but these indices remain above the long-term average CPUE (Figure 5).

B_{MSY} conceptual proxy: The Plenary agreed in 2017 to use the mean combined ELE5-BT(MIX) CPUE for the period 2005–06 to 2015–16 as a “B_{MSY} conceptual proxy” for ELE 5. This period was selected because a plot of CPUE against catch (yield curve) appeared to have levelled out and is assumed to represent a stochastic equilibrium (Figure 6).




Figure 5: Plots of three ELE5-BT(MIX) CPUE series: a) positive catch (lognormal); b) presence/absence (binomial) and c) combined series using the delta-lognormal method.




Figure 6: Trace yield plot for ELE 5, showing CPUE and QMR/MHR landings plotted sequentially by fishing year.

• ELE 7

A preliminary CPUE analysis of the catch of elephant fish from the WCSI inshore trawl fishery was conducted in 2013 and updated in 2014 (Langley 2014). The analysis included all bottom trawl catch and effort data targeting either flatfish, red gurnard, red cod or elephant fish. These target trawl fisheries encompass almost all the trawl fishing effort within the depth range that encompasses most of the catch of elephant fish off the west coast of the South Island (5–80 m). The primary analysis was conducted based on catch and effort data from 1989–90 to 2012–13 aggregated in a format that was consistent with the CELR reporting format. The landed catch of elephant fish from each trip was apportioned to the effort records either based on the associated level of estimated catch or, where estimated catches were not recorded, in proportion to the number of trawls in each aggregated effort record.

The data set included a significant proportion of trip and effort records with no elephant fish catch, although the proportion of nil catch records decreased steadily over the study period. Thus, the overall CPUE for the fishery was modelled in two components: the binomial model of the proportion of positive catches and the lognormal model of the magnitude of the positive catch. The two components were combined to generate a time series of delta-lognormal CPUE indices. The sensitivity of the catch threshold used to define a positive catch (i.e. 0, 1 kg, 2 kg and 5 kg) was investigated. The resulting binomial and lognormal CPUE indices were sensitive to the applied catch threshold; however, the compensatory changes in the two sets of indices resulted in delta-lognormal indices that were relatively insensitive to the applied catch threshold.

The resulting CPUE indices fluctuated over the study period with a marked peak in CPUE in 1999–2000 and 2000–01 and low CPUE in 1997–98 and 2003–04 (Figure 7). The CPUE indices remained stable during 2007–08 to 2009–10, increased in 2010–11, increased markedly in 2011–12 and remained at the higher level in 2012–13. In 2014, the SINS WG concluded that the CPUE indices were unlikely to be a reliable index of stock abundance, primarily on the basis that the large inter-annual variations in the CPUE indices especially during the late 1990s and early 2000s were not consistent with the dynamics of the stock and may be attributable to changes in the operation of the WCSI trawl fishery at that time.

A separate delta-lognormal CPUE analysis was conducted for the location based TCER catch and effort

ELEPHANT FISH (ELE)

data from 2007–08 to 2012–13 (Langley 2014). The resulting CPUE models incorporated a number of additional explanatory variables available in the high resolution data format. The TCER delta-lognormal CPUE indices were broadly similar to the CELR format CPUE indices for the comparative period. The TCER indices exhibited a comparable increase in CPUE from 2009–10 to 2011–12, although the TCER indices were higher in 2007–08 to 2008–09 than the CELR format indices. In 2015, the TCER CPUE indices were updated to include the 2013–14 fishing year (Figure 7). The SINS WG concluded that the TCER CPUE indices represented the best available information for monitoring trends in ELE 7 stock abundance.




Figure 7. Standardised Delta-lognormal CPUE indices for the ELE 7 inshore WCSI trawl fishery for the entire time series configured in CELR data format and for indices derived from the location based TCER data set. Both sets of indices are normalised to the comparable time period (2007–08 to 2012–13). The error bars represent the 95% confidence interval.

Stock Assessment models

A preliminary stock assessment model was developed for ELE 3. Estimates of current and reference absolute biomass are not available for the other elephant fish stocks.

ELE 3

A stock assessment model was developed for ELE 3 in 2016 using the Stock Synthesis (3.24f) software to implement an age-structured population model. The data sets available for inclusion in the assessment model are, as follows.

- Annual reported catch of elephant fish (1931–2015). The historical catches were derived from Francis & Paul (2013). Additional unreported landed catches were included for the period prior to the introduction of the QMS. The level of unreported landed catch was assumed to represent a third of the reported catch. The magnitude of unreported landed catch was based on discussions with commercial operators in the ELE 3 fishery.
- A time-series of estimates of the magnitude of the discarded catch (unreported but not landed) of elephant fish (1931–2015). Based on the discussions with commercial operators it was assumed that the discarded (and unreported catch) represented 25% of total landed catch (reported and unreported combined). The discarded catch is comprised of smaller elephant fish, usually less than 50 cm FL.
- BT MIX CPUE indices 1989–90 to 2014–15 (26 observations).

ELEPHANT FISH (ELE)

- ECSI trawl survey pre-recruit (< 50 cm), recruited (50+ cm) and total biomass estimates from the time series of winter surveys, 30–400 m depth (11 observations).
- ECSI trawl survey length compositions (male and female); winter surveys, 30–400 m depth (11 observations).
- Aggregated length compositions (male and female) of the commercial trawl catch sampled by Scientific Observers during 2009–10.

Additional data are available from the summer ECSI trawl surveys. These data were not included in the analysis as it has previously been concluded that the summer survey series does not represent a reliable index of abundance for elephant fish. In recent years, the winter trawl survey has been extended to include the shallower areas of Canterbury Bight and Pegasus Bay (10–30 m), partly to improve the monitoring of the abundance of elephant fish. However, the time-series of surveys that includes this area is limited (four surveys).

Initial modelling results revealed that the scaled length compositions derived from the winter trawl surveys were highly variable (amongst surveys) and inconsistent with the other key input data sets. Further examination of the length composition data revealed that few elephant fish were caught and sampled during each survey and the scaled length compositions were typically dominated by the sampled catch from a limited number of trawls. The length and sex compositions of these larger catches were highly variable.

On that basis, it was concluded that the survey length compositions were unlikely to be representative of the length composition of the elephant fish population and these data were excluded from the final set of model options. Further, the estimates of trawl survey biomass for pre-recruit (<50 cm) fish are relatively imprecise (CVs 32–83%) and preliminary modelling indicated that these indices were not consistent with the other abundance indices (especially the CPUE indices). Thus, the pre-recruit trawl survey biomass indices were also excluded from the final set of model options.

Model configuration

The final assessment model was configured, as follows.

- Model period 1931–2015, terminal year represents 2014–15 fishing year.
- Age classes 0–19 and 20+ years, two sexes.
- Initial (1931) population age structure assumes equilibrium, unexploited conditions.
- Annual recruitment derived from Beverton and Holt stock-recruitment relationship; R_0 parameter estimated (uninformative beta prior) and steepness fixed at 0.6 (base model option), recruitment deviates from SRR estimated for 1989–2013 assuming a SigmaR of 0.6.
- Sexual maturity (female fish) at 70 cm (FL).
- Two commercial fisheries: discard and retained catch. The selectivity of the commercial catch is assumed to be equivalent for the two main fishing methods (BT and SN).
- Commercial length composition data from 2009–10 are partitioned at 50 cm to characterise the length composition of discard (<50 cm) and retained (50+ cm) commercial catches. Both length compositions are assigned a relatively high weighting (ESS 100) to ensure that the model approximates these observations.
- The length-based selectivity of discard commercial fishery is parameterised using a double normal selectivity function (equivalent for both sexes). Selectivity is effectively truncated at about 50 cm (FL).
- Two alternative length-based selectivity options were adopted for the retained commercial fishery with selectivity parameterised using either a logistic or double normal function. Selectivity was allowed to vary by sex.
- The CPUE indices are assumed to represent the relative abundance of the component of the population that is vulnerable to the retained commercial fishery. The CPUE indices were assigned a CV of 20%.
- The ECSI recruited (50+ cm) total biomass estimates were assigned the native CVs from individual surveys. The length-based selectivity of the survey was assumed to be knife edge at 50 cm (FL) with full selectivity for all the larger length intervals.

ELEPHANT FISH (ELE)

Model options that assumed a logistic selectivity function for the (retained) commercial fishery resulted in a poor fit to the (retained) commercial length composition for male and female fish (from 2009–10). These models consistently over-estimated the number of larger male (>68 cm FL) and female (>90 cm FL) elephant fish in the commercial catch.

The alternative model option with selectivity parameterised by a double normal function resulted in a substantial improvement in the fit to the commercial length compositions (relative to the logistic selectivity model). The double normal selectivity model estimated selectivity for male and female fish started to rapidly decline above 70 cm and 85 cm FL, respectively. The lower selectivity of larger female fish meant that approximately 40–50% of the mature female population (by weight) is estimated to be invulnerable to the commercial fishery and, consequently, not monitored by the CPUE indices.

Separate model runs were conducted for the two selectivity options, each with three assumed values of SRR steepness: a base level of 0.6 bracketed by values of 0.5 and 0.7. MCMCs were conducted for the six model options. However, the results of the MCMCs were not satisfactory for the model options with the lowest value of steepness and, consequently, only MCMC results for the 0.6 steepness options are reported.

Model results

The overall fit to the CPUE indices was acceptable for all model options. The CPUE indices exhibit a general increase with marked peaks in the early and late 2000s. The models account for these trends by estimating higher recruitments for 1996–1998, 2004, and 2009. As previously noted, the double normal selectivity parameterisation substantially improved the fit to the retained commercial length composition data (compared to logistic selectivity). There was also a marginal improvement in the fit to the CPUE indices with the double normal selectivity.

All model options also estimated an increase in stock abundance that was consistent with the overall increase in the ECSI trawl survey recruited biomass estimates between the 1990s and the more recent period, although the fit to the individual biomass estimates is poor. The quality of the fit is consistent with the relatively low precision of the biomass estimates and the likelihood that the survey vulnerability of elephant fish varies amongst survey years (as indicated by the variability in the length composition of the survey catches).

Two indicators of stock status were derived from the assessment models: current (2014–15) female spawning (=mature) biomass relative to unexploited spawning biomass (SB_{2015}/SB_0), and current spawning biomass relative to the spawning biomass in 1985 (SB_{2015}/SB_{1985}). The latter metric provides an indication of the extent of the stock recovery from the period when the stock was estimated to be at the lowest level.

The MPD results indicate that stock abundance has increased considerably from a low level (approx. 10–20% SB_0) in 1985. The double normal selectivity model runs represent a somewhat more optimistic estimate of the current stock status relative to both SB_0 and SB_{1985} . MPD estimates of stock status tended to be near the lower bound of the MCMC confidence intervals, indicating that the MPD estimates are likely to represent minimum biomass levels consistent with the catch history.

Table 6: Estimates of stock status for the range of commercial selectivity and SRR steepness options (MPD estimates). MCMC estimates (median value and 95% confidence interval) are also presented for the two selectivity options with SRR steepness of 0.60.

Selectivity	Steepness		SB_{2015}/SB_0	SB_{2015}/SB_{1985}
Double normal	0.6	MPD	0.390	2.99
		MCMC	0.471 (0.266–0.872)	2.86 (2.08–3.97)
	0.7	MPD	0.321	3.77
Logistic	0.6	MPD	0.279	2.50
		MCMC	0.386 (0.217–0.651)	2.63 (1.86–3.61)
	0.7	MPD	0.229	3.03

The results are also sensitive to the assumptions regarding SRR steepness. Higher values of steepness

correspond to lower estimates of SB_0 and a higher level of depletion by 1985, and while the relative level of recovery from 1985 is higher than for lower steepness options, the current level of stock biomass relative to SB_0 is lower.

The median estimates of SB_{2015}/SB_0 stock status from the MCMCs are more optimistic than the corresponding MPD results for the SRR steepness 0.60 model runs. The MCMC results also reveal that there is considerable uncertainty associated with the estimates of stock status, although the confidence intervals derived from the MCMCs suggest that current biomass is Likely to be above the default soft limit (20% SB_0) and About As Likely as Not to be at or above the default target biomass level (40% SB_0). However, the preliminary nature of the model precludes definitive statements about stock status.

These conclusions need to be tempered by the possibility that the models may be over-estimating recruitment in the more recent years. This may provide an explanation for the apparent over-estimation of the proportion of larger, older fish in the population in the late 2000s (that were not apparent in the commercial length composition). Conversely, the recent CPUE indices may be biased low (due to apparent avoidance behaviour) and consequently the model may under-estimate the current level of biomass.

Estimates of SB_{2015}/SB_0 stock status are also highly uncertain (and potentially biased) due to the assumptions associated with the estimation of historical, unexploited biomass.




Figure 8: Stock trajectories for the spawning biomass relative to SB_0 (upper panels) and SB_{1985} (lower panels) for logistic (left panels) and double normal (right panels) selectivity options with SRR steepness 0.6. The black line represents the median of the MCMCs (with 95% confidence interval) and the red line represents the MPD.

The Southern Inshore Working Group concluded that this preliminary model produced plausible biomass trajectories, but uncertainty about productivity and fits to commercial length data precluded acceptance of

ELEPHANT FISH (ELE)

the model as a reliable estimator of current stock status.

Yield estimates and projections

No other yield estimates are available.

4.4 Other factors

A data informed qualitative risk assessment was completed on all chondrichthyans (sharks, skates, rays and chimaeras) at the New Zealand scale in 2014 (Ford et al 2015). Elephant fish was ranked fourth highest in terms of risk of the eleven QMS chondrichthyan species. Data were described as existing and sound for the purposes of the assessment and consensus over this risk score was achieved by the expert panel. This risk assessment does not replace a stock assessment for this species but may influence research priorities across species.

5. STATUS OF THE STOCKS

- **ELE 1**

No estimates of current and reference biomass are available.

- **ELE 2**

It is not known if recent catch levels or the current TACC are sustainable. The state of the stock in relation to B_{MSY} is unknown.

- **ELE 3**


Stock Structure Assumptions

No information is available on the stock separation of elephant fish. The Fishstock ELE 3 is treated in this summary as a unit stock.

Stock Status	
Year of Most Recent Assessment	2016
Assessment Runs Presented	Update ELE 3 (MIX) CPUE series
Reference Points	InterimTarget: B_{MSY} -compatible proxy based on CPUE (average from 1998–99 to 2010–11 of the ELE 3(MIX) model as defined in Starr & Kendrick 2013) Soft Limit: 50% of target Hard Limit: 25% of target Overfishing threshold: F_{MSY} (assumed)
Status in relation to Target	About as Likely as Not (40–60%) to be at or above the target
Status in relation to Limits	Soft Limit: Unlikely (< 40%) to be below Hard Limit: Very Unlikely (< 10%) to be below
Status in relation to Overfishing	Overfishing is About as Likely as Not (40–60%) to be occurring

Historical Stock Status Trajectory and Current Status

CPUE, Catch and TACC Trajectories




Comparison of the mixed target species bottom trawl CPUE series (ELE 3(MIX)) with the trajectories of catch (ELE 3(QMR/MHR)) and TACCs from 1989–90 to 2014–15. The dashed lines represent the interim target and corresponding soft limit and hard limit.

Fishery and Stock Trends

Recent trend in Biomass or Proxy

The ELE 3(MIX) CPUE series, which is considered to be an index of stock abundance, showed a generally increasing trend from the beginning to reach a peak in 2007–08. CPUE indices have remained relatively stable below the peak level since 2009–10, remaining near the proposed target.

Recent trend in Fishing Intensity or Proxy



Fishing mortality proxy is Standardised Fishing Effort = Total catch/CPUE (normalised). Fishing mortality proxy has fluctuated about the average level and was at about the average in the most recent year.

ELEPHANT FISH (ELE)

Other Abundance Indices	Although there is high inter-annual variation, the winter ECSI trawl survey index shows a trend that is consistent with the ELE 3(MIX) CPUE index. Preliminary stock assessment modelling for ELE 3 estimates that the stock abundance has increased substantially from a low level in the 1980s. The assessment models indicate that current biomass levels are probably at or about the default target biomass levels.
Trends in Other Relevant Indicator or Variables	-

Projections and Prognosis

Stock Projections or Prognosis	Quantitative stock projections are unavailable.
Probability of Current Catch or TACC causing decline Biomass to remain below or to decline below Limits	Soft Limit: Unlikely (< 40%) Hard Limit: Very Unlikely (< 10%)
Probability of Current Catch or TACC causing Overfishing to continue or to commence	The TACC and current reported catches are About as Likely as Not (40–60%) to cause overfishing.

Assessment Methodology and Evaluation

Assessment Type	Level 2: Partial Quantitative Stock Assessment	
Assessment Method	Evaluation of agreed standardised CPUE indices which reflect changes in abundance.	
Assessment Dates	Latest assessment: 2016	Next assessment: Unknown
Overall assessment quality rank	1 – High Quality. The Southern Inshore Working Group agreed that the ELE 3(MIX) CPUE index was a credible measure of abundance.	
Main data inputs (rank)	- Catch and effort data	1 – High Quality
Data not used (rank)	- Compass Rose trawl survey data - Summer ECSI trawl survey data - Winter ECSI trawl survey data	3 – Low Quality: insufficient data 2 – Medium or Mixed Quality: variable catchability / selectivity between years 3 – Low Quality: Index compromised by area closures
Changes to Model Structure and Assumptions	None since 2012 assessment	
Major Sources of Uncertainty	- It is possible that fisher avoidance and discarding have biased (low) the CPUE trends reported for this fishery.	

• Qualifying Comments

- Elephant fish have shown good recovery since apparently being at low biomass levels in the mid-1980s.
- Preliminary stock assessment modelling results are consistent with assumed level of stock rebuilding, primarily reflecting the increase in the CPUE abundance indices. However, there are considerable uncertainties associated with key biological parameters (natural mortality and growth) and conflict amongst the main input data sets. The modelling results are not considered to be sufficiently reliable to estimate current stock status (relative to MSY levels) and potential yields for the stock. With respect to the conceptual B_{msy} proxy, the Plenary had concerns about the reliability of this as a proxy and suggested that it only be used on an interim basis.
- Historical catches may be poorly estimated. Both current and historical estimates of landings exclude fish discarded at sea and the quantum of discards is unknown. Management interventions since the stock was introduced into the QMS may have influenced the rate of discarding and therefore the reliability of CPUE as a measure of relative abundance.

• Fishery Interactions

Elephant fish in ELE 3 are taken as bycatch by bottom trawl fisheries targeting red cod, flatfish and barracouta. Targeting elephant fish in the bottom trawl fishery has increased to around 40% of the landings since 2004–05 when the deemed value regime changed. Around 15% of the ELE 3 landings are taken by setnet in a fishery targeted at a number of shark species, including rig, elephant fish, spiny dogfish and school shark. Both the trawl and setnet fisheries have been subject to management measures designed to reduce interactions with endemic Hector's dolphins. Bottom trawl fishers also have not trawled within one nautical mile of the coast (since 2001) in an effort to preserve ELE egg cases. This may have reduced juvenile and egg mortality in shallow water. Interactions with other species are currently being characterised.

• ELE 5


Stock Structure Assumptions

No information is available on the stock separation of elephant fish. The Fishstock ELE 5 is treated in this summary as a unit stock.


Stock Status	
Year of Most Recent Assessment	2017
Assessment Runs Presented	Standardised bottom trawl CPUE series based on mixed target species: combined delta-lognormal series
Reference Points	Target: B_{MSY} -compatible proxy based on mean ELE5-BT(MIX) standardised CPUE: 2005–06 to 2015–16 Soft Limit: 50% of Bmsy proxy Hard Limit: 25% of Bmsy proxy Overfishing threshold: Mean annual relative exploitation rate for the period: 2005–06 to 2015–16
Status in relation to Target	About as Likely as Not (40-60%) to be at or above Bmsy
Status in relation to Limits	Soft Limit: Unlikely (< 40%) to be below Hard Limit: Very Unlikely (< 10%) to be below
Status in relation to Overfishing	Overfishing is About as Likely as Not (40–60%) to be occurring

ELEPHANT FISH (ELE)

Historical Abundance and Catch Trajectories



Comparison of the ELE 5-BT(MIX) CPUE series with the TACC and QMR/MHR landings for ELE 5. The agreed BMSY proxy (geometric average: 2006–2016 ELE 5-BT(MIX) CPUE indices=2.051) is shown as a green line; the calculated Soft Limit ($=0.5 \times \text{BMSY}$ proxy) is shown as a purple line; the calculated Hard Limit ($=0.25 \times \text{BMSY}$ proxy) is shown as a grey line.



Relative fishing pressure for ELE 5 based on the ratio of QMR/MHR landings relative to the ELE5-BT(MIX) CPUE series which has been normalised so that its geometric mean=1.0. Horizontal green line is the geometric mean fishing pressure from 2006 to 2016.

Fishery and Stock Trends

Recent trend in Biomass or Proxy	The ELE 5 (MIX) CPUE series increased up to a peak in 2008–09, dropped sharply in 2011–12 and has fluctuated without trend close to the target since then.
Recent Trend in Fishing Mortality or Proxy	Fishing mortality proxy has remained relatively stable or declining over the last 10 years.
Other Abundance Indices	-
Trends in Other Relevant Indicator or Variables	-
Projections and Prognosis	
Stock Projections or Prognosis	Unknown
Probability of Current Catch and TACC causing biomass to remain below or to decline below Limits	Soft Limit: Unlikely (< 40%) Hard Limit: Very Unlikely (< 10%)

Probability of Current Catch or TACC causing Overfishing to continue or to commence	Current Catch: About as Likely as Not (40–60%) TACC: About as Likely as Not (40–60%)
---	---

Assessment Methodology and Evaluation		
Assessment Type	Level 2: Partial Quantitative Stock Assessment	
Assessment Method	Evaluation of agreed standardised CPUE indices	
Assessment Dates	Latest assessment: 2017	Next assessment: 2020
Overall assessment quality rank	1 – High Quality	
Main data inputs (rank)	- ELE 5 BT(MIX) CPUE series	1 – High Quality
Data not used (rank)	Length frequency data summarised from setnet logbooks compiled under the industry Adaptive Management Programme	3 – Low Quality: data sparse and outdated
Changes to Model Structure and Assumptions	Addition of a binomial index to produce a combined CPUE series	
Major Sources of Uncertainty	It is possible that discarding and management changes (including changes in deemed values) in this fishery has affected CPUE estimates.	

Qualifying Comments
Elephant fish have shown strong recovery since apparently being at low biomass levels in the mid-1980s. The historical catches may be poorly estimated. Both current and historical estimates of landings exclude fish discarded at sea and the quantum of discards is unknown. Confidence intervals for combined CPUE indices are not available.


Fishery Interactions
Elephant fish in ELE 5 are taken by bottom trawl in fisheries targeted at flatfish and stargazer. Targeting elephant fish in the bottom trawl fishery was low (average 14% from 1989–90 to 2015–16) but has increased to 19% of the landings since 2002–03. Around 12% of the ELE 5 landings are taken by setnet in a fishery targeted at rig and school shark. Incidental captures of seabirds and great white sharks occur, and there is a possibility of incidental capture of Hector's dolphins. However, both the trawl and setnet fisheries have been subject to management measures designed to reduce interactions with endemic Hector's dolphins. Interactions with other species are currently being characterised.

• ELE 7

Stock Status	
Year of Most Recent Assessment	2015
Assessment Runs Presented	ELE 7 standardised CPUE based mixed target species in the bottom trawl fishery
Reference Points	Target: Not established but B_{MSY} assumed Soft Limit: 20% B_0 Hard Limit: 10% B_0 Overfishing threshold: F_{MSY} (assumed)
Status in relation to Target	Unknown
Status in relation to Limits	Soft Limit: Unlikely (< 40%) Hard Limit: Very Unlikely (< 10%)
Status in relation to Overfishing	Overfishing is Unlikely (< 40%) to be occurring

ELEPHANT FISH (ELE)

Historical Stock Status Trajectory and Current Status



Standardised TCER CPUE index for ELE 7 (black dots), commercial landings (yellow line) and TACC (red line).

Fishery and Stock Trends

Recent Trend in Biomass or Proxy	CPUE indices indicate biomass increased considerably from 2009–10 to 2011–12, remained at the higher level in 2012–13 and declined in 2013–14.
Recent Trend in Fishing Intensity or Proxy	Catches declined from a high in 1998–99 to a low in 2003–04 but have risen to and fluctuated around the level of the TACC since 2006–07.
Other Abundance Indices	Trawl survey biomass trends for this stock are unreliably estimated by the West Coast South Island survey. However, recent biomass estimates have been relatively high compared to the long term average.
Trends in Other Relevant Indicators or Variables	-

Projections and Prognosis

Stock Projections or Prognosis	CPUE indices and catches for 2011–12 and 2012–13 were relatively high levels (series beginning 1989–90), lower in 2013–14. Recent trawl survey biomass estimates are also relatively high.
Probability of Current Catch or TACC causing Biomass to remain below or to decline below Limits	Soft Limit: Unlikely (< 40%) Hard Limit: Unlikely (< 40%)
Probability of Current Catch or TACC causing Overfishing to continue or to commence	Current catches and the current TACC are Unlikely (< 40%) to cause overfishing.

Assessment Methodology and Evaluation

Assessment Type	Level 2: Partial Quantitative Stock Assessment	
Assessment Method	Standardised CPUE index and relative biomass estimates from inshore WCSI trawl survey	
Assessment dates	Latest assessment: 2015	Next assessment: Unknown
Overall assessment quality rank	1 – High Quality	

ELEPHANT FISH (ELE)

ELEPHANT FISH (L)		
Main data inputs (rank)	<ul style="list-style-type: none">- Standardised CPUE (MIX) (from 2007–08)- Standardised CPUE (MIX) (pre 2007–08)- Catch and effort data derived from the Fisheries New Zealand compulsory catch reporting system	<p>1 – High Quality: The SINSWG had more confidence in this part of the CPUE index as a credible measure of abundance</p> <p>2 – Medium or Mixed Quality: less catch (data) and lack of spatial resolution</p> <p>1 – High Quality</p>
Data not used (rank)	<ul style="list-style-type: none">- Biomass estimates from inshore WCSI trawl survey	2 – Medium or Mixed Quality: low precision and high variability
Changes to Model Structure and Assumptions	-	
Major Sources of Uncertainty	<ul style="list-style-type: none">- It is possible that discarding and management changes in this fishery have biased the CPUE trends.- The CPUE indices are derived from a data set with a high proportion of zero catch records and the indices may be sensitive to the treatment of zero catch records (although this was not apparent from a limited number of sensitivity analyses conducted).	
Qualifying Comments		
The pre-QMS catches are not well reported. Both current and historical estimates of landings exclude fish discarded at sea and the quantum of discards is unknown.		

Fishery Interactions

Trawl target sets for ELE 7 tend to be in shallow water mostly around 25 m. Elephant fish are landed with rig, school shark and spiny dogfish in setnets and in bottom trawls as bycatch in flatfish and red cod target sets. Incidental captures of seabirds occur and there is a possibility of incidental capture of Hector's dolphins. Interactions with other species are currently being characterised.

7. FOR FURTHER INFORMATION

- Beentjes, M P; MacGibbon, D; Lyon, W S (2015) Inshore trawl survey of Canterbury Bight and Pegasus Bay, April–June 2014 (KAH1402). *New Zealand Fisheries Assessment Report 2015/14*. 136 p.
- Beentjes, M.P.; MacGibbon, D.; Parkinson, D. (2016). Inshore trawl survey of Canterbury Bight and Pegasus Bay, April–June 2016 (KAH1605). *New Zealand Fisheries Assessment Report 2016/61*. 135 p.
- Boyd, R O; Reilly, J L (2002) 1999/2000 national marine recreational fishing survey: harvest estimates. Draft New Zealand Fisheries Assessment Report. (Unpublished report held by Fisheries New Zealand, Wellington.)
- Bradford, E (1998) Harvest estimates from the 1996 national recreational fishing surveys. New Zealand Fisheries Assessment Research Document 1998/16. 27 p. (Unpublished report held by NIWA library, Wellington.)
- Coakley, A (1971) The biological and commercial aspects of the elephantfish. *Fisheries Technical Report* No: 76. 29 p.
- Coutin, P (1992) Sharks... and more sharks. *Australian fisheries* June 1992: 41–42.
- Ford, R B; Galland, A; Clark, M R; Crozier, P; Duffy, C A J; Dunn, M R; Francis, M P; Wells, R (2015) Qualitative (Level 1) Risk Assessment of the impact of commercial fishing on New Zealand Chondrichthyans. New Zealand Aquatic Environment and Biodiversity Report No. 157. 111 p.
- Francis, M P (1996) Productivity of elephantfish - has it increased? *Seafood New Zealand* February 96: 22–25.
- Francis, M P (1997) Spatial and temporal variation in the growth rate of elephantfish (*Callorhynchus milii*). *New Zealand Journal of Marine and Freshwater Research* 31: 9–23.
- Francis, R I C; Hurst, R J; Renwick, J A (2001) An evaluation of catchability assumptions in New Zealand stock assessments. *New Zealand Fisheries Assessment Report 2001/1*. 37 p.
- Francis, M P; Paul, L J (2013) New Zealand inshore finfish and shellfish commercial landings, 1931–82. *New Zealand Fisheries Assessment Report 2013/55*. 136 p.
- Gorman, T B S (1963) Biological and economic aspects of the elephantfish, *Callorhynchus milii* Bory, in Pegasus Bay and the Canterbury Bight. *Fisheries Technical Report* No: 8. 54 p.
- Langley, A D (2001) The analysis of ELE 3 catch and effort data from the RCO 3 target trawl fishery, 1989–90 to 1999–2000. *New Zealand Fisheries Assessment Report 2001/66*. 33 p.
- Langley, A D (2014) Updated CPUE analyses for selected South Island inshore finfish stocks. *New Zealand Fisheries Assessment Report 2014/40*.



ELEPHANT FISH (ELE)

- 116 p.
- Langley, A D (2015) Fishery characterisation and Catch-Per-Unit-Effort indices for John dory in JDO 1. *New Zealand Fisheries Assessment Report 2015/47*. 76 p.
- Lydon, G J; Middleton, D A J; Starr, P J (2006) Performance of the ELE 3 and ELE 5 Logbook Programmes. AMP-WG-06/18. (Unpublished manuscript available from the NZ Seafood Industry Council, Wellington.)
- MacGibbon, D J; Stevenson, M L (2013) Inshore trawl survey of the west coast South Island and Tasman and Golden Bays, March-April 2013 (KAH1305). *New Zealand Fisheries Assessment Report 2013/66*. 115 p.
- McClatchie, S; Lester, P (1994) Stock assessment of the elephantfish (*Callorhinchus milii*). New Zealand Fisheries Assessment Research Document 1994/6. 17 p. (Unpublished report held by NIWA library, Wellington.)
- Raj, L; Voller, R (1999) Characterisation of the south-east elephantfish fishery-1998. 55 p. (Unpublished report available at NIWA library, Wellington.)
- Seafood Industry Council (SeaFIC) (2000) Proposal to the Inshore Fishery Assessment Working Group. Placement of the ELE 3 into Adaptive Management Programme dated 23 March 2000 (presented to the Inshore Fishery Assessment Working Group 28 March 2000). (Unpublished report held by Fisheries New Zealand, Wellington.)
- Seafood Industry Council (SeaFIC) (2002) Report to the Inshore Fishery Assessment Working Group: Performance of the ELE 3 Adaptive Management Programme (dated 25 February 2002). (Unpublished report held by Fisheries New Zealand, Wellington.)
- Seafood Industry Council (SeaFIC) (2003a) 2003 performance report: ELE 3 Adaptive Management Programme. AMP-WG-2003/06. 3 p. (Unpublished report held by Fisheries New Zealand, Wellington.)
- Seafood Industry Council (SeaFIC) (2003b) Report to the Adaptive Management Fishery Assessment Working Group: Performance of the ELE 5 Adaptive Management Programme and request for an additional increase in ELE 5. AMP-WG-2003/07. 39 p. (Unpublished report held by Fisheries New Zealand, Wellington.)
- Seafood Industry Council (SeaFIC) (2005a) 2005 Report to the Adaptive Management Programme Fishery Assessment Working Group: Performance of the ELE 3 Adaptive Management Programme. AMP-WG-2005/16. (Unpublished report held by Fisheries New Zealand, Wellington.)
- Seafood Industry Council (SeaFIC) (2005b) 2005 Report to the Adaptive Management Programme Fishery Assessment Working Group: Performance of the ELE 5 Logbook Programme. AMP-WG-05/23. (Unpublished report held by Fisheries New Zealand, Wellington.)
- Southeast Finfish Management Company (SEFMC) (2002a) 2002 Report to the Inshore Fishery Assessment Working Group. Performance of the ELE 3 Adaptive Management Programme (dated 25 February 2002). (Unpublished report held by Fisheries New Zealand, Wellington.)
- Southeast Finfish Management Company (SEFMC) (2002b) 2002 Report to the Inshore Fishery Assessment Working Group. Performance of the ELE 5 Adaptive Management Programme (dated 25 February 2002). (Unpublished report held by Fisheries New Zealand, Wellington.)
- Southeast Finfish Management Company (SEFMC) (2003) 2003 Report to the Inshore Fishery Assessment Working Group. Performance of the ELE 5 Adaptive Management Programme and request for an increase in ELE 5 (dated 13 Nov 2003). (Unpublished report held by , Wellington.)
- Starr, P J; Kendrick, T H (2013) ELE 3&5 Fishery Characterisation and CPUE. *New Zealand Fisheries Assessment Report 2013/38*. 95 p.
- Starr, P.J.; Kendrick, T.H. (2017). ELE 5 Fishery Characterisation and CPUE Report. *New Zealand Fisheries Assessment Report 2017/50*. 63 p.
- Starr, P J; Kendrick, T H; Lydon, G J; Bentley, N (2007a) Report to the Adaptive Management Programme Fishery Assessment Working Group: Full-term review of the ELE 3 Adaptive Management Programme. AMP-WG-07/07. 104 p. (Unpublished report held by Fisheries New Zealand, Wellington.)
- Starr, P J; Kendrick, T H; Lydon, G J; Bentley, N (2007b) Report to the Adaptive Management Programme Fishery Assessment Working Group: Two-year review of the ELE 5 Adaptive Management Programme. AMP-WG-07/10. 89 p. (Unpublished report held by Fisheries New Zealand, Wellington.)
- Sullivan, K J (1977) Age and growth of the elephantfish *Callorhinchus milii* (Elasmobranchii: Callorhynchidae). *New Zealand Journal of Marine and Freshwater Research* 11: 745–753.
- Teirney, L D; Kilner, A R; Millar, R E; Bradford, E; Bell, J D (1997) Estimation of recreational catch from 1991/92 to 1993/94. New Zealand Fisheries Assessment Research Document 1997/15. 43 p. (Unpublished report held by NIWA library, Wellington.)
- Wynne-Jones, J; Gray, A; Hill, L; Heinemann, A (2014) National Panel Survey of Marine Recreational Fishers 2011–12: Harvest Estimates. *New Zealand Fisheries Assessment Report 2014/67*. 145 p.

FLATFISH (FLA)

(*Colistium nudipinnis*, *Peltorhamphus novaezealandiae*, *Colistium guntheri*, *Rhombosolea retiaria*, *Rhombosolea plebeia*, *Rhombosolea leporina*, *Rhombosolea tapirina*, *Pelotretis flavilatus*)

Patiki



1. FISHERY SUMMARY

1.1 Commercial fisheries

Flatfish Individual Transferable Quota (ITQ) provides for the landing of eight species of flatfish. These are: the yellow-belly flounder, *Rhombosolea leporine* (YBF); sand flounder, *Rhombosolea plebeian* (SFL); black flounder, *Rhombosolea retiaria* (BFL); greenback flounder, *Rhombosolea tapirina* (GFL); lemon sole, *Pelotretis flavilatus* (LSO); New Zealand sole, *Peltorhamphus novaezealandiae* (ESO); brill, *Colistium guntheri* (BRI); and turbot, *Colistium nudipinnis* (TUR). For management purposes landings of these species are combined.

Flatfish are shallow water species, taken mainly by target inshore trawl and Danish seine fleets around the South Island. Set and drag net fishing are important in the northern harbours and the Firth of Thames. Important fishing areas are:

Yellow-belly flounder	Firth of Thames, Kaipara and Manukau harbours;
Sand flounder	Hauraki Gulf, Tasman/Golden Bay, Bay of Plenty, Canterbury Bight and Te Wae Wae Bay;
Greenback flounder	Canterbury Bight, Southland;
Black flounder	Canterbury Bight;
Lemon sole	west coast South Island, Otago and Southland;
New Zealand sole	west coast South Island, Otago, Southland and Canterbury Bight;
Brill and turbot	west coast South Island.

TACCs were originally set at the level of the sum of the provisional ITQs for each fishery. Between 1983–84 and 1992–93 total flatfish landings fluctuated between 2750 t and 5160 t; from 1992–93 to 1997–98, landings were relatively consistent, between about 4500 t and 5000 t per year. Landings declined to 2963 t in 1999–00, the lowest recorded since 1986–87, then increased to a peak of 4051 t for the 2006–07 fishing year, and then declined to a new low of 2464 t in 2014–15. Total NZ FLA landings for 2016–17 were 2851 t. Historical estimated and recent reported flatfish landings and TACCs are shown in Tables 1 and 2, while Figure 1 shows the historical landings and TACC values for the main FLA QMAs. From 1 October 2007, a TAC and allowances were set for the first time in FLA 3. The FLA 3 TACC was reduced by 47% to 1430 t as well as implementing a management procedure that recommends an in-season increase in the commercial catch allowance if supported by early CPUE data (see Section 4.3 for a description of this procedure). All FLA fisheries have been put on to Schedule 2 of the Fisheries Act 1996. Schedule 2 allows that, for certain “highly variable”

FLATFISH (FLA)

stocks, the Total Annual Catch (TAC) can be increased within a fishing season. Increased commercial catch is provided for through the creation of additional ‘in-season’ ACE. The base TACC is not changed by this process and the “in-season” TAC reverts to the original level at the end of each season. The FLA 3 management procedure (Section 4.3) is an implementation of this form of management.

From 1 October 2008, a suite of regulations intended to protect Maui’s and Hector’s dolphins was implemented for all of New Zealand by the Minister of Fisheries. Commercial and recreational set netting were banned in most areas to 4 nautical miles offshore of the east coast of the South Island, extending from Cape Jackson in the Marlborough Sounds to Slope Point in the Catlins. Some exceptions were allowed, including an exemption for commercial and recreational set netting to only one nautical mile offshore around the Kaikoura Canyon, and permitting setnetting in most harbours, estuaries, river mouths, lagoons and inlets, except for the Avon-Heathcote Estuary, Lyttelton Harbour, Akaroa Harbour and Timaru Harbour. In addition, trawl gear within 2 nautical miles of shore was restricted to flatfish nets with defined low headline heights. The commercial minimum legal size for sand flounder is 23 cm, and for all other flatfish species is 25 cm.

Table 1: Reported landings (t) for the main QMAs from 1931 to 1982.

Year	FLA 1	FLA 2	FLA 3	FLA 7	Year	FLA 1	FLA 2	FLA 3	FLA 7
1931–32	767	290	219	265	1957	308	64	529	183
1932–33	958	219	61	276	1958	362	59	989	321
1933–34	698	277	181	346	1959	362	48	971	382
1934–35	708	203	83	195	1960	410	58	1257	361
1935–36	686	118	57	209	1961	386	102	665	273
1936–37	438	127	139	139	1962	383	156	584	228
1937–38	570	125	380	123	1963	352	106	627	228
1938–39	717	83	639	94	1964	499	134	879	350
1939–40	721	128	448	83	1965	599	109	917	518
1940–41	1004	180	494	101	1966	547	222	1141	496
1941–42	943	139	622	139	1967	646	231	1273	493
1942–43	591	192	594	154	1968	541	139	973	311
1943–44	669	89	606	172	1969	686	193	936	269
1944	441	104	783	78	1970	557	262	1027	471
1945	435	104	984	83	1971	407	149	1028	276
1946	392	168	1264	146	1972	475	114	548	166
1947	551	99	1685	198	1973	438	149	717	442
1948	433	93	1494	214	1974	503	147	637	748
1949	412	76	1473	202	1975	431	156	598	476
1950	284	31	1446	176	1976	548	132	802	929
1951	308	62	1178	135	1977	764	255	916	1165
1952	349	94	1117	166	1978	706	202	1730	1225
1953	349	149	1510	197	1979	742	287	1962	899
1954	376	112	1184	213	1980	906	219	1562	459
1955	377	125	913	248	1981	1082	760	1369	399
1956	308	106	772	190	1982	934	650	1214	468

1. The 1931–1943 years are April–March but from 1944 onwards are calendar years.
2. Data up to 1985 are from fishing returns; Data from 1986 to 1990 are from Quota Management Reports.
3. Data for the period 1931 to 1982 are based on reported landings by harbour and are likely to be underestimated as a result of under-reporting and discarding practices. Data includes both foreign and domestic landings.

Table 2: Reported landings (t) of flatfish by Fishstock from 1983–84 to present and actual TACCs (t) from 1986–87 to the present. QMS data from 1986–present. [Continued on next page.]

Fishstock FMA (s)	FLA 1 1 & 9		FLA 2 2 & 8		FLA 3 3, 4, 5 & 6		FLA 7 7		FLA 10 10		Total	
	Landings	TACC	Landings	TACC	Landings	TACC	Landings	TACC	Landings	TACC	Landings	TACC
1983–84*	1 215	-	378	-	1 564	-	1 486	-	0	-	5 160	-
1984–85*	1 050	-	285	-	1 803	-	951	-	0	-	4 467	-
1985–86*	722	-	261	-	1 537	-	385	-	0	-	3 215	-
1986–87	629	1 100	323	670	1 235	2 430	563	1 840	0	10	12 750	6 050
1987–88	688	1 145	374	677	2 010	2 535	1 000	1 899	0	10	14 072	6 266
1988–89	787	1 153	297	717	2 458	2 552	757	2 045	0	10	4 299	6 477
1989–90	791	1 184	308	723	1 637	2 585	745	2 066	0	10	3 482	6 568
1990–91	849	1 187	292	726	1 340	2 681	502	2 066	0	10	2 983	6 670
1991–92	940	1 187	288	726	1 229	2 681	745	2 066	0	10	3 202	6 670
1992–93	1 106	1 187	460	726	1 954	2 681	1 566	2 066	0	10	5 086	6 670

Table 2 [Continued]

Fishstock FMA (s)	FLA 1 1 & 9		FLA 2 2 & 8		FLA 3 3, 4, 5 & 6		FLA 7 7		FLA 10 10		Total	
	Landings	TACC	Landings	TACC	Landings	TACC	Landings	TACC	Landings	TACC	Landings	TACC
1993–94	1 136	1 187	435	726	1 926	2 681	1 108	2 066	0	10	4 605	6 670
1994–95	964	1 187	543	726	1 966	2 681	1 107	2 066	0	10	4 580	6 670
1995–96	628	1 187	481	726	2 298	2 681	1 163	2 066	1	10	4 571	6 670
1996–97	741	1 187	363	726	2 573	2 681	1 117	2 066	0	10	4 794	6 670
1997–98	728	1 187	559	726	2 351	2 681	1 020	2 066	0	10	4 657	6 670
1998–99	690	1 187	274	726	1 882	2 681	868	2 066	0	10	3 714	6 670
1999–00	751	1 187	212	726	1 583	2 681	417	2 066	0	10	2 963	6 670
2000–01	792	1 187	186	726	1 702	2 681	447	2 066	0	10	3 127	6 670
2001–02	596	1 187	177	726	1 693	2 681	614	2 066	0	10	3 080	6 670
2002–03	686	1 187	144	726	1 650	2 681	819	2 066	0	10	3 299	6 670
2003–04	784	1 187	218	726	1 286	2 681	918	2 066	0	10	3 206	6 670
2004–05	1 038	1 187	254	726	1 353	2 681	1 231	2 066	0	10	3 876	6 670
2005–06	964	1 187	296	726	1 177	2 681	1 283	2 066	0	10	3 720	6 670
2006–07	922	1 187	296	726	1 429	2 681	1 419	2 066	0	10	4 066	6 670
2007–08	703	1 187	243	726	1 365	1 430	1 313	2 066	0	10	3 624	5 419
2008–09	639	1 187	214	726	1 544	**1 780	1 020	2 066	0	10	3 417	5 419
2009–10	652	1 187	212	726	1 525	**1 763	884	2 066	0	10	3 273	5 835
2010–11	486	1 187	296	726	1 027	1 430	659	2 066	0	10	2 467	5 509
2011–12	445	1 187	262	726	1 507	1 430	646	2 066	0	10	2 861	5 419
2012–13	480	1 187	274	726	1 512	**1 727	526	2 066	0	10	2 792	5 716
2013–14	511	1 187	216	726	1 377	1 430	568	2 066	0	10	2 672	5 419
2014–15	426	1 187	166	726	1 231	1 430	640	2 066	0	10	2 464	5 419
2015–16	277	1 187	238	726	1 622	**1 650	656	2 066	0	10	2 792	5 639
2016–17	421	1 187	136	726	1 421	**2 065	873	2 066	0	10	2 851	6 054

* FSU data.

† Includes 11 t Turbot, area unknown but allocated to QMA 7.

§ Includes landings from unknown areas before 1986–87.

** Commercial catch allowance increased with additional ‘in-season’ ACE provided under S68 of FA1996

**# The increase in commercial catch under S68 of FA1996 was not approved until late August 2017

Fishers and processors are required to use a generic flatfish (FLA) code in the monthly harvest returns to report landed catches of flatfish species as well as in the landings section of the catch and effort forms. Although fishers are now instructed to use specific species codes when reporting estimated catches, they more often use the generic FLA code. Beentjes (2003) showed that, for all QMAs combined between 1989–90 and 2001–02, about half of the estimated catch of flatfish was recorded using the generic species code FLA, and the remainder was reported using a combination of 12 other species codes (Table 3). Flatfish species that comprised a large proportion of the total estimated catch over the 13 year period included ESO (16%), LSO (12%), SFL (12%) and YBF (6%). Species that are important contributors to catch in each QMA are FLA 1: YBF, SFL, GFL; FLA 2: ESO, SFL; FLA 3: ESO, LSO, SFL, BFL, BRI; FLA 7: GFL, SFL, TUR (codes described in the caption to Table 3). Starr & Kendrick (in prep) have recently shown that trips which report catches in FLA 3 by species rather than using the generic FLA code accounted for greater than 80% of the estimated catches in 2012–13 and 2013–14.

Table 3: Percent estimated flatfish catch by species and fishing year in FLA 3 for “splitter” trips, which are trips which landed FLA 3 but which did not use the FLA code in the estimated catch section of the catch/effort form. Codes are arranged in descending order of total estimated catch: lemon sole (LSO), New Zealand sole (ESO), sand flounder (SFL), black flounder (BFL), brill (BRI), yellow belly flounder (YBF), Turbot (TUR), greenback flounder (GFL) (Starr & Kendrick in prep). Also shown is the proportion by weight of estimated catch defined in the “splitter” category.

Year	LSO	ESO	SFL	BFL	BRI	YBF	FLO	TUR	GFL	Other "Splitters"
1990–91	14.7	32.1	22.2	18.1	5.2	4.5	0.0	1.3	1.9	0.0
1991–92	23.9	41.7	15.3	1.7	3.5	8.5	0.0	1.3	4.0	0.0
1992–93	23.6	42.9	20.3	0.4	3.2	4.5	0.0	0.4	4.8	0.0
1993–94	32.9	43.2	14.4	0.3	2.3	2.3	0.0	0.7	3.9	0.0
1994–95	34.8	35.4	16.3	3.5	2.0	2.8	0.0	1.1	3.6	0.5
1995–96	40.6	34.0	11.9	6.1	2.3	2.4	0.7	0.7	0.9	0.4
1996–97	38.2	36.8	14.6	2.4	2.0	1.2	2.4	0.7	1.6	0.1
1997–98	54.5	26.1	10.8	0.7	1.6	1.3	2.3	0.7	1.8	0.1
1998–99	57.2	22.4	8.9	1.3	2.7	2.0	2.4	1.6	1.4	0.1
1999–00	42.0	31.8	9.7	6.4	4.2	2.9	0.7	2.0	0.4	0.1
2000–01	36.4	37.3	9.7	3.5	3.2	2.9	1.1	1.9	0.2	3.8
2001–02	26.3	44.5	10.8	8.6	2.6	2.0	1.0	1.4	0.3	2.5
2002–03	33.0	40.2	11.2	2.2	4.1	4.3	1.3	1.8	0.2	1.7
2003–04	39.1	30.1	9.6	1.7	2.8	10.8	0.8	0.7	0.1	4.3
2004–05	33.9	27.0	12.7	13.4	2.9	3.6	1.1	1.2	0.3	3.9

FLATFISH (FLA)

Table 3 [Continued]

Year	LSO	ESO	SFL	BFL	BRI	YBF	FLO	TUR	GFL	Other "Splitters"
2005–06	46.3	25.0	12.1	5.3	2.9	3.0	2.1	0.9	1.1	1.3
2006–07	52.0	20.6	15.9	0.1	2.5	4.6	1.8	1.2	0.5	0.8
2007–08	65.4	18.2	7.3	0.0	3.3	0.7	1.3	1.1	1.9	0.7
2008–09	54.9	25.6	10.2	0.0	3.0	0.7	1.8	1.9	1.5	0.4
2009–10	59.9	19.3	11.4	0.3	3.1	1.0	1.4	1.8	1.0	0.8
2010–11	54.7	14.4	16.8	2.4	4.7	0.4	2.0	2.4	0.9	1.4
2011–12	51.0	18.6	15.0	4.2	3.4	0.6	3.4	2.5	0.3	1.0
2012–13	46.4	20.7	16.9	2.4	3.3	1.9	3.2	2.4	0.6	2.0
2013–14	39.2	20.7	21.9	3.2	3.4	4.4	2.5	2.4	1.2	1.2
Total	42.7	29.6	13.3	3.4	3.0	2.8	1.5	1.4	1.4	1.1




Figure 1: Historical landings and TACC for the four main FLA stocks. FLA 1 (Auckland), FLA 2 (Central).

1.2 Recreational fisheries

There are important recreational fisheries, mainly for the four flounder species, in most harbours, estuaries, coastal lakes and coastal inlets throughout New Zealand. The main methods are setnetting, drag netting (62.8% combined) and spearing (36.1%) (Wynne-Jones et al 2014). In the northern region, important areas include the west coast harbours, the lower Waikato, the Hauraki Gulf and the Firth of Thames. In the Bay of Plenty, Ohiwa and Tauranga Harbours are important. In the Challenger FMA, there is a moderate fishery in Tasman and Golden Bays and in areas of the Mahau-Kenepuru Sound and in Cloudy Bay. In the South-East and Southland FMAs, flatfish are taken in areas such as Lake Ellesmere, inlets around Banks Peninsula and the Otago Peninsula, the Oreti and Riverton estuaries, Bluff Harbour and the inlets and lagoons of the Chatham Islands (for further details see the 1995 Plenary Report).




Figure 1 [Continued]: Historical landings and TACC for the four main FLA stocks. FLA 3 (South East Coast, South East Chatham Rise, Sub-Antarctic, Southland).

1.2.1 Management controls

The main method used to manage recreational harvests of flatfish are minimum legal sizes (MLS) and daily bag limits. General spatial and method restrictions also apply, particularly to the use of set nets. The flatfish MLS for recreational fishers is 25 cm for all species except sand flounder for which the MLS is 23 cm. Fishers can take up to 20 flatfish as part of their combined daily bag limit in the Auckland, Central and Challenger Fishery Management Areas. Fishers can take up to 30 flatfish as part of their combined daily bag limit in the South-East, Kaikoura, Fiordland and Southland Fishery Management Areas.

1.2.2 Estimates of recreational harvest

There are two broad approaches to estimating recreational fisheries harvest: the use of onsite or access point methods where fishers are surveyed or counted at the point of fishing or access to their fishing activity; and, offsite methods where some form of post-event interview and/or diary are used to collect data from fishers.

The first estimates of recreational harvest for flatfish were calculated using an offsite regional telephone-diary survey approach. Estimates for 1996 came from a national telephone-diary survey (Bradford 1998). Another national telephone-diary survey was carried out in 2000 (Boyd & Reilly 2005). The harvest estimates provided by telephone-diary surveys between 1993 and 2001 are no longer considered reliable for various reasons. A Recreational Technical Working Group concluded that these harvest estimates should be used only with the following qualifications: a) they may be very inaccurate; b) the 1996 and earlier surveys contain a methodological error; and c) the 2000 and 2001 estimates are implausibly high for many important fisheries. In response to these problems and the cost and scale challenges associated with onsite methods, a National Panel Survey was conducted for the first time throughout the 2011–12 fishing year. The panel survey used face-to-face interviews of a random sample of 30 390 New Zealand households to recruit a panel of fishers and non-fishers for a full year. The panel members were contacted regularly about their fishing activities and harvest information collected in standardised phone interviews. Harvest estimates for blue mackerel are given in Table 4 (from Wynne-Jones et al 2014 and Hartill & Davey 2015).

A repeat of the National Panel Survey is being conducted over the 2017–18 October fishing year. Results are expected in early 2019.

FLATFISH (FLA)

Table 4: Estimated number and weight of flatfish, by Fishstock and survey, harvested by recreational fishers. Surveys were carried out in different years in the Fisheries regions: South in 1991–92, Central 1992–93, North 1993–94 (Teirney et al 1997) and nationally in 1996 (Bradford 1998) and 1999–00 (Boyd & Reilly 2005). (- Data not available). National panel survey conducted 01 October 2011 through 30 September 2012, used seasonally variable mean weights for flatfish of 345 to 426 g (Wynne-Jones et al 2014, Hartill & Davey 2015).

Fishstock	Survey	Number	CV	Harvest range (t)	Point estimate (t)
1991–92					
FLA 1	South	3 000	-	-	-
FLA 3	South	15 200	0.31	50–90	-
FLA 7	South	3 000	-	-	-
1992–93					
FLA 1	Central	6 100	-	-	-
FLA 2	Central	73 000	0.26	20–40	-
FLA 7	Central	37 100	0.59	10–30	-
1993–94					
FLA 1	North	520 000	0.19	225–275	-
FLA 2	North	3 000	-	0–5	-
1996					
FLA 1	National	308 000	0.11	95–125	110
FLA 2	National	67 000	0.19	13–35	24
FLA 3	National	113 000	0.14	30–50	40
FLA 7	National	44 000	0.18	10–20	16
1999–00					
FLA 1	National	702 000	0.25	203–336	-
FLA 2	National	380 000	0.49	82–238	-
FLA 3	National	395 000	0.33	128–252	-
FLA 7	National	114 000	0.53	23–73	-
2011–12					
FLA 1	Panel	64 999	0.37	-	27.2
FLA 2	Panel	12 885	0.31	-	5.4
FLA 3	Panel	53 475	0.31	-	21.7
FLA 7	Panel	12 259	0.37	-	4.7
All areas combined	Panel	143 619	0.21	-	58.9

1.3 Customary non-commercial fisheries

Quantitative information on the current level of customary non-commercial catch is not available.

1.4 Illegal catch

There is no quantitative information on the current level of illegal catch available.

1.5 Other sources of mortality

The extent of unrecorded fishing mortality is unknown.

2. BIOLOGY

Some New Zealand flatfish species are fast-growing and short-lived, generally only surviving to 3–4 years of age, with very few reaching 5–6 years, others such as brill and turbot are longer lived, reaching a maximum age of 21 years and 16 years, respectively (Stevens et al 2001). However, these estimates have yet to be fully validated. Size limits (set at 25 cm for most species) are generally at or above the size at which the fish reach maturity and confer adequate protection to the juveniles.

Sutton et al (2010) undertook an age and growth analysis of greenback flounder. That analysis showed that growth is rapid throughout the lifespan of greenback flounder. Females reached a slightly greater maximum length than males, but the difference was not significant at the 95% level of confidence. Over 90% of sampled fish were 2 or 3 years of age, with maximum ages of 5 and 10 years being obtained for male and female fish respectively. This difference in maximum age resulted in estimated natural mortalities using Hoenig's (1983) regression method, of 0.85 for males and 0.42 for females. It is suggested that 0.85 is the most appropriate estimate at this stage as only 1% of all fish exceeded 5 years. However, it was also noted that a complete sample of the larger fish was not obtained and as a result these estimates should be considered preliminary. Growth rings were not validated.

Flatfish are shallow-water species, generally found in waters less than 50 m depth. Juveniles congregate in sheltered inshore waters, e.g., estuarine areas, shallow mudflats and sandflats, where they remain for up to two years. Juvenile survival is highly variable. Flatfish move offshore for first spawning at 2–3 years of age during winter and spring. Adult mortality is high, with many flatfish spawning only once and few spawning more than two or three times. However, fecundity is high, e.g., from 0.2 million eggs to over 1 million eggs in sand flounders.

Available biological parameters relevant to stock assessment are shown in Table 5. The estimated parameters in sections 1 and 3 of the table apply only to sand flounder in Canterbury and brill and turbot in west coast South island - growth patterns are likely to be different for these species in other areas and for other species of flatfish.

Table 5: Estimates of biological parameters for flatfish.

Fishstock	Estimate				Source		
<u>1. Natural mortality (<i>M</i>)</u>							
Brill - West coast South Island (FLA 7)	0.20				Stevens et al (2001)		
Turbot - West coast South island (FLA 7)	0.26				Stevens et al (2001)		
Sand flounder - Canterbury (FLA 3)	1.1–1.3				Colman (1978)		
Lemon sole - West coast South island (FLA 7)	0.62–0.96				Gowing et al (unpub.)		
<u>2. Weight = a(length)^b (Weight in g, length in cm total length).</u>							
	Females		Males				
	a	b	a	b			
Brill (FLA 7)	0.01443	2.9749	0.02470	2.8080	Hickman & Tait (unpub.)		
Turbot (FLA 7)	0.00436	3.3188	0.00571	3.1389	Hickman & Tait (unpub.)		
Sand flounder (FLA 1)	0.03846	2.6584	-	-	McGregor et al (unpub.)		
Yellow-belly flounder (FLA 1)	0.07189	2.5117	0.00354	3.3268	McGregor et al (unpub.)		
New Zealand sole (FLA 3)	0.03578	2.6753	0.007608	3.0728	McGregor et al (unpub.)		
<u>3. von Bertalanffy growth parameters</u>							
	Females			Males			
	<i>L</i> _∞	<i>k</i>	<i>t</i> ₀	<i>L</i> _∞	<i>k</i>	<i>t</i> ₀	
Brill							
West coast South Island (FLA 7)	43.8	0.10	−15.87	38.4	0.37	38.4	Stevens et al (2001)
Turbot							
West coast South island (FLA 7)	57.1	0.39	0.30	49.2	0.34	49.2	Stevens et al (2001)
Sand flounder							
Canterbury (FLA 3)	59.9	0.23	−0.083	37.4	0.781	37.4	Mundy (1968), Colman (1978)
Lemon sole							
West coast South island (FLA 7)	26.1	1.29	−0.088	25.6	1.85	25.6	Gowing et al (unpub.)
Greenback flounder (FLA 5)	55.82	0.26	−1.06	52.21	0.25	−1.32	Sutton et al (2010)

3. STOCKS AND AREAS

There is evidence of many localised stocks of flatfish. However, the inter-relationships of adjacent populations have not been well studied. The best information is available from studies of the variation in morphological characteristics of sand flounders and from the results of tagging studies, conducted mainly on sand and yellow-belly flounders. Variation in morphological characteristics indicate that sand flounder stocks off the east and south coasts of the South Island are clearly different from stocks in central New Zealand waters and from those off the west coast of the South Island. There also appear to be differences between west coast sand flounders and those in Tasman Bay, and between sand flounders on either side of the Auckland-Northland peninsula. Tagging experiments show that sand flounders, and other species of flounder, can move substantial distances off the east and south coasts of the South Island. However, fish tagged in Tasman Bay or the Hauraki Gulf have never been recaptured very far from their point of release.

Thus, although the sand flounders off the east and south of the South Island appear to be a single, continuous population, fish in enclosed waters may be effectively isolated from neighbouring populations and should be considered as separate stocks. Examples of such stocks are those in Tasman Bay and the Hauraki Gulf and possibly areas such as Hawke Bay and the Bay of Plenty.

FLATFISH (FLA)

There are no new data which would alter the stock boundaries used in previous assessment documents.

4. STOCK ASSESSMENT

4.1 Estimates of fishery parameters and abundance

FLA 1

Four standardised CPUE series have been used to track FLA 1 abundance (Kendrick & Bentley 2011; Kendrick & Bentley 2012), which were last updated in 2015 (Kendrick & Bentley in prep.):

1. FLA+YBF in Manukau Harbour (Statistical Area 043);
2. FLA+YBF in Kaipara Harbour (Statistical Area 044);
3. YBF in Hauraki Gulf (Statistical Areas 005, 006, and 007);
4. SFL in Hauraki Gulf (Statistical Areas 005, 006, and 007).

These series were updated in 2018 with an additional three years of data (Starr & Kendrick in prep.), based on declared estimated catches as well as using a procedure (termed “F2”) which scales estimated catches to landings using a “vessel correction factor” by multiplying estimated catches with the ratio of landings to estimated catches for a vessel in a fishing year. A comparison of the two series showed no material difference in output between the two procedures, even though the F2 procedure truncates the data set in order to avoid excessively large and small ratios. Starr & Kendrick (in prep.) also summed all flatfish estimated catches for the Manukau and Kaipara Harbour analyses to create a TOT category. This was done because estimated catches of other flatfish species are negligible in these harbours (Table 6) and a comparison with 2015 series showed no difference in the overlapping years. The Northern Inshore Working Group accepted series 1, 2 and 3 (above) as reflecting abundance. However, the SFL series in the Hauraki Gulf was rejected by the NINSWG because it was noted that the reporting of SFL in the estimated catches fell away in the early to mid-2000s which was also a period when the SFL CPUE dropped while, at the same time, there was little change in the species-specific reporting of YBF. This trend in the reporting pattern for SFL makes the associated CPUE series unreliable, resulting in a recommendation that the SFL series be replaced with a TOT series (which sums all flatfish species catch).

Less than half of the estimated FLA 1 flatfish catch in each year is identified by species (Table 6), but most of the flatfish caught in FLA 1 West are likely to be yellow-belly flounder under the assumption that the flatfish reported using the generic “FLA” code are YBF. This assumption is supported by the fact that the preferred muddy bottom habitat of yellow-belly flounder dominates the west coast harbours. Over 80% of the west coast catch is taken from Kaipara and Manukau Harbours (Table 6). Standardised CPUE trends were derived for these two areas using TOT (sum of all flatfish estimated catches) or the F2 procedure applied to the TOT estimated catches (upper panels, Figure 2). In spite of fluctuations, both the Manukau and Kaipara series show a long-term declining trend and are currently 68% and 65% below the respective peaks in the early to mid-1990s (upper panels, Figure 2). Work by NIWA (McKenzie et al 2013) in the Manukau Harbour has linked the decrease in local CPUE with an increase in eutrophication, suggesting that there may be factors other than fishing contributing to the decline.

Table 6: Total FLA 1 estimated catches by declared flatfish species, summed over the period 1989–90 to 2016–17.

	Manukau	Kaipara	Lower Waikato	Northwest	FLA 1 West	East Northland	Hauraki Gulf	Bay of Plenty	FLA 1 East	Total FLA 1
FLA	1 876.2	2 682.4	543.9	523.1	5 625.6	565.3	3 097.1	264.3	3 926.7	9 552.3
YBF	127.4	1 661.8	96.6	163.5	2 049.3	401.9	2 510.9	133.4	3 046.2	5 095.5
SFL	3.9	44.0	18.5	8.7	75.1	58.2	1 198.5	308.0	1 564.7	1 639.8
ESO	0.0	0.0	10.8	16.1	26.9	1.1	5.4	204.7	211.2	238.1
GFL	0.0	0.1	7.5	0.2	7.8	0.0	202.6	12.7	215.3	223.1
LSO	0.0	0.0	1.2	2.4	3.6	0.5	1.0	76.8	78.3	81.9
BRI	0.0	0.0	7.4	2.6	10.0	0.1	0.1	19.5	19.7	29.7
BFL	0.0	0.0	0.1	0.2	0.3	0.3	26.3	2.3	28.9	29.2
TUR	0.0	0.0	4.3	4.4	8.7	0.1	0.3	1.2	1.6	10.3
Total	2 007.5	4 388.2	690.4	721.3	7 807.4	1 027.6	7 042.2	1 022.7	9 092.5	16 899.9




Figure 2: Standardised CPUE indices for yellowbelly flounder from models of catch rate in successful set net trips in Manukau Harbour, Kaipara Harbour (YBF assumed) and in the Hauraki Gulf (YBF reported). Also shown is the series for total FLA in Hauraki Gulf. All models based on estimated catches scaled by a “vessel correction factor” (F2 procedure).

Seventy-seven percent of the flatfish catch from FLA 1 East, including a substantial and variable proportion of sand flounder, is taken in the Hauraki Gulf, particularly from the Firth of Thames (Area 007). Separate indices were calculated for sand and yellowbelly flounder in Statistical Areas 005 to 007, and the portion of FLA catch not identified by species was excluded. However, the SFL series was not accepted by the NINSWG in 2018 (see above for rationale) and a FLA(TOT) series was prepared instead. The Hauraki Gulf yellow belly CPUE index peaked in 2006–07 and then declined steadily to 2015–16. However, there was a sharp upturn in the YBF series in 2016–17, with the final index returning to above the series mean (lower left panel, Figure 2). A total FLA series for the Hauraki Gulf was created to replace the rejected sand flounder index in the same region (lower right panel, Figure 2). This series shows an overall declining trend except for a three-year increase from 2002 to 2005 and a single strong increase in the final 2017 fishing year, which brings the series above the long-term average.

FLA 2

In 2017, Schofield et al (2018a) provided standardised CPUE for FLA 2 (Figure 3) based on the flatfish target fishery in Statistical Areas 013 and 014. Estimated catches were allocated to daily aggregated effort using methodology described in Langley (2014) to improve the comparability between the data collected from two different statutory reporting forms (CELR and TCER). A core fleet of 15 vessels that had completed at least five trips per year in at least seven years was identified. The model, using a gamma error distribution, adjusted for changes in duration, month and vessel, and accounted for 33% of the variance in catch. Area was not included in the model as the change in reporting forms appears to have influenced the catch split between areas 013 and 014.

The NINS WG noted that most of the records in the aggregated data had catches of flatfish and that a binomial index was flat. As a result the positive catch index was retained as the key monitoring series. The CPUE series exhibits moderate fluctuations around the long term mean, with no overall trend up or down and appears currently to be in an increasing phase.

FLATFISH (FLA)

Characterisation using the estimated catch data suggests that the FLA 2 catch comprises mainly sand flounder (SFL) and New Zealand sole (ESO). CPUE indices for ESO and SFL were provided by Schofield et al (2018a) for 2008 to 2016 using the tow by tow data from vessels consistently estimating catches by flatfish species. Trends were apparent in the probability of catch, so combined (binomial and positive catch modelled with a gamma distribution) indices were produced. There is reasonable consistency between the species specific indices and the overall FLA 2 index (Figure 3), noting that – as the FLA 2 fishery is small - the datasets for the individual species are small and the indices variable.

These indices were updated in 2018 (Schofield et al 2018b) to include data to 30 September 2017.

Establishing B_{MSY} compatible reference points

In 2014, the Working Group adopted mean CPUE from the bottom trawl flatfish target series for the period 1989–90 to 2012–13 as a B_{MSY} -compatible proxy for FLA 2. The Working Group accepted the default Harvest Strategy Standard definitions that the Soft and Hard Limits would be one half and one quarter the target, respectively.




Figure 3: Standardised CPUE indices in FLA 2 for BT targeting all species of flatfish, (aggregated to combine data across form types, BT_flats(day)), and shorter combined series for sand flounder (BT_sfl(tow)) and New Zealand sole (BT_eso(tow)) based on tow by tow resolution data (Schofield et al 2018b).

FLA 3

CPUE trends

As in 2010 (Kendrick & Bentley in prep), CPUE trends for the three principal FLA 3 species (New Zealand sole [ESO], sand flounder [SFL] and lemon sole [LSO]) and an aggregated catch landed to FLA [TOT], based on bottom trawl catch and effort data, were estimated. The species-specific data were based on “splitter” trips, defined as trips which landed FLA 3 but which did not use the FLA code in the estimated catch section of the catch/effort form. Alternative definitions of “splitters” based on vessel performance were also investigated, but CPUE trends were found to be similar to those

derived from the “trip splitter” algorithm. The latter was selected because it retained the greatest amount of catch, particular in the early years of the series.

The CPUE data were prepared by matching the landing data for a trip with the effort data from the same trip that had been amalgamated to represent a day of fishing. The procedure assigns the modal statistical area and modal target species (defined as the observation with the greatest effort) to the trip/date record. All estimated catches for the day were summed and the five top species with the greatest catch were assigned to the date. This “daily-effort stratum” preparation method was followed so that the event-based data forms that are presently being used in these fisheries can be matched as well as possible with the earlier daily forms to create a continuous CPUE series. Each analysis was confined to a set of core vessels which had participated consistently in the fishery for a reasonably long period (ESO, LSO and SFL: 5 trips for at least 5 years; TOT: 10 trips for at least 5 years). The explanatory variables offered to each model included fishing year (forced), month, vessel, statistical area, number tows and duration of fishing.

These trends were used to evaluate the relative status of these species and to predict in-season abundance of FLA based on early harvest returns for the fishery. There are similarities in the fluctuations of the four standardised CPUE indices (Figure 4), with all indices increasing in the early 1990s and peaking at some point in the five years between 1989–90 and 1993–94. All indices then have a trough in the early- to mid-2000s, followed by an increase for LSO and SFL and a decrease for ESO. The FLA, ESO and SFL indices show the greatest similarity in their fluctuations. The LSO index had its peak in the 1990s; i.e. later than the other indices, and increased sooner than the other species in the mid-2000s (Figure 4). The SFL index has continued to increase up to 2013–14 while the other three indices have dropped from peaks reached in 2009–10.




Figure 4: Comparison of standardised bottom trawl lognormal CPUE indices in FLA 3 for FLA (all flatfish species combined) LSO (lemon sole), ESO (New Zealand sole) and SFL (sand flounder). Note that only the FLA index is available for the 1989–90 fishing year because very little species composition data are available for that year (Starr & Kendrick, in prep).

ECSI trawl survey biomass estimates for LSO

Lemon sole biomass indices in the core strata (30–400 m) for the East Coast South Island trawl survey (Table 7) show no trend (Figure 5). Coefficients of variation are moderate to low, ranging from 18 to 33% (mean 24%). The additional biomass captured in the 10–30 m depth range accounted for only 4% and 1% of the biomass in the core plus shallow strata (10–400 m) for 2007 and 2012, respectively, indicating that the existing core strata time series in 30–400 m are the most important, but that shallow strata should also be monitored. A comparison of the two sets of LSO biomass indices shows that both series fluctuate without trend, with considerable variability (Figure 6). However, the correspondence between the two sets of indices is weak ($\rho = -0.294$; $R^2 = 9\%$).




Figure 5: Lemon sole total biomass and 95 % confidence intervals for all ECSI winter surveys in core strata (30–400 m), and core plus shallow strata (10–400 m) in 2007, 2012 and 2014.




Figure 6: Lemon sole total biomass and 95% confidence intervals for the all ECSI winter surveys in core strata (30–400 m) plotted against the LSO bottom trawl CPUE series.

Table 7: Relative biomass indices (t) and coefficients of variation (CV) for lemon sole for the east coast South Island (ECSI) - winter survey area.

Region	Fishstock	Year	Trip number	Total Biomass estimate (t)	CV (%)
ECSI (winter)	FLA 3: LSO				30–400 m
		1991	KAH9105	89	27
		1992	KAH9205	57	18
		1994	KAH9406	77	21
		1996	KAH9606	49	33
		2007	KAH0705	74	26
		2008	KAH0806	116	25
		2009	KAH0905	55	27
		2012	KAH1207	65	18
		2014	KAH1402	107	27

In-season Management Procedure

In 2007 concerns were expressed about the sustainability of FLA 3 catches and the TACC was reduced from 2 681 t to 1 430 t from 1 October 2007. In the 2008–09 fishing year anecdotal information indicated an increase in abundance of lemon and New Zealand sole in the FLA 3 QMA above a level that fishers were able to utilise within the available TACC. It was considered that there was opportunity for increased utilisation that would not adversely impact on the long term sustainability of the FLA 3 stock complex and for 2008–09 ‘in-season’ commercial allowances were set at 1 780 t based on the 15 year average of commercial FLA3 catches.

In 2010 an ‘in-season’ Management Procedure (MP) was developed and has been used to inform in-season adjustments to the FLA 3 TACC since 2010–11 (Kendrick & Bentley in prep.), being updated and revised in 2015 (Starr et al in prep.). This MP used the relationship between annual standardised CPUE for all FLA 3 species (shown as FLA in Figure 4) and the total annual FLA 3 landings to estimate an average exploitation rate which is then used to recommend a level of full-season catch based on an early estimate of standardised CPUE. Only the period 1989–90 to 2006–07 was used to estimate the average exploitation rate because this was the period before the TACC was reduced which allowed the fishery to operate at an unconstrained level. A partial year in-season estimate of standardised CPUE is used as a proxy for the final annual index, with the recommended catch defined by the slope of the regression line (Figure 7) multiplied by the CPUE proxy estimate (Figure 8 shows the outcome of this procedure).

The previous FLA 3 MP, adopted in 2010, approximated the standardisation procedure by applying fixed coefficients to a data set specified by a static core vessel definition. This approach deteriorated over time as vessels dropped out of the core vessel fleet, thus reducing the available data set. The revised 2015 MP is based on a re-estimated standardisation procedure using a data set specified annually by a dynamic core vessel definition, allowing new vessels to enter the data set as they meet the minimum eligibility criteria. The 2015 MP was validated through a retrospective analysis which used the data available up to end of the previous year and the partial data in the final year to determine how the model performed across years (Figure 8). In most years, the MP performance was satisfactory after only two months of data were accumulated. The poor performance of the model in some years (e.g., 2012) persisted across all four early months, indicating that collecting additional data in those years would not have improved the recommendation (relative to the end of year recommendation).

FLATFISH (FLA)




Figure 7: [left panel] Relationship between annual FLA 3 CPUE (=FLA in Figure 4) and total annual FLA 3 QMR/MHR landings from 1989–90 to 2006–07 (calculated for the 2018 inseason MP); [right panel]: residuals from the left panel regression.




Figure 8: Operation of the 2015 FLA 3 MP in 2018, showing the relationship of the fitted catch estimates to the observed MHR/QMR landings and the annual recommended catches from 2008 onward based on the estimated standardised CPUE up to the end of November.

Table 8 shows the results of the operation of the FLA 3 in-season MP since the inception of the Schedule 2 programme. Five TACC in-season increases have been recommended since 2010 based on the operation of the MP (2009–10, 2010–11, 2012–13, 2015–16 and 2016–17; Table 8). However, MPI approval of the 2016–17 increase was delayed until late August, resulting in limited opportunity to take advantage of the increase in commercial catch allowance.

Table 8: Results of the operation of the FLA 3 MP by prediction year. NA: not available.

Prediction Year	Fishing Year	CPUE Prediction	CPUE Total year ¹	Recom- mended commercial allowance	Approved commercial allowance (t) ²	Annual catch (t)	Date of Approval ²	Reference
2010*	2009–10	64.98 (kg/tow)	75.82	1 846	1 763	1 525	18 June 2010	Bentley (2010)
2011*	2010–11	59.83 (kg/tow)	58.76	1 520	1 430	1 027	–	Bentley (2011)
2012	2011–12	58.45 (kg/tow)	57.56	1 495	–	1 507	–	Bentley (2012)
2013*	2012–13	67.97 (kg/tow)	69.70	1 727	1 727	1 512	17 May 2013	Brouwer (2013)
2014	2013–14	NA	54.80	NA	–	1 377	–	NA
2015	2014–15	53.20 (kg/tow)	NA	1 362	1 352	1 231	–	Bentley (2015)
2016*	2015–16	0.984	1.048	1 650	1 650	1 622	15 July 2016	Starr et al (2016)
2017*	2016–17	1.215	0.978	2 065	2 065	1 421	23 Aug 2017	Starr & Kendrick (2017)
2018	2017–18	0.870	–	1 461	NA	NA	NA	Starr & Kendrick (2018)

¹ calculated in the year following² information provided by MPI

* MP operation that resulted in a commercial catch allowance increase recommendation

Establishing B_{MSY} compatible reference points

The Working Group accepted mean CPUE from the bottom trawl flatfish target series for the period 1989–90 to 2006–07 as a B_{MSY} -compatible proxy for FLA and 1990–01 to 2006–07 for LSO, SFL and ESO. These periods were chosen as catches were not constrained by the TACC. 1989–90 to 2006–07 was also the period used to determine average exploitation rate for the in season adjustment Management Procedure. The Working Group accepted the default Harvest Strategy Standard definitions that the Soft and Hard Limits would be one half and one quarter the target, respectively.

4.2 Other Factors

The flatfish complex is comprised of eight species although typically only a few are dominant in any one QMA and some are not found in all areas. For management purposes all species are combined to form a unit fishery. The proportion that each species contributes to the catch is expected to vary annually. It is not possible to estimate MCY for each species and stock individually.

Because the adult populations of most species generally consist of only one or two year classes at any time, the size of the populations depends heavily on the strength of the recruiting year class and is therefore thought to be highly variable. Brill and turbot are notable exceptions with the adult population consisting of a number of year classes. Early work revealed that although yellow belly flounder are short-lived, inter-annual abundance in FLA 1 was not highly variable, suggesting that some factor, e.g., size of estuarine nursery area, could be smoothing the impact of random environmental effects on egg and larval survival. Work by NIWA (McKenzie et al 2013) in the Manukau harbour has linked the decrease in local CPUE with an increase in eutrophication, suggesting that there may be factors other than fishing contributing to the decline.

Flatfish TACCs were originally set at high levels so as to provide fishers with the flexibility to take advantage of the perceived variability associated with annual flatfish abundance. This approach has been modified with an in-season increase procedure for FLA 3.

4.2 Research needs

- Conduct CPUE analyses for brill and turbot, which are two of the longest-lived flatfish species and as such may be more susceptible to overfishing and depletion, particularly if they are caught in conjunction with other more productive species.


5. STATUS OF THE STOCKS


Estimates of current and reference biomass are not available.

- Yellow-belly flounder in FLA 1**


Stock Structure Assumptions

Based on tagging studies, yellow-belly flounder appear to comprise localised populations, especially in enclosed areas such as harbours and bays.





CPUE and total annual estimated catches for YBF in Kaipara Harbour. Also shown is the fishing intensity (catch/CPUE), standardised relative to the geometric mean. Fishing year designated by second year of the pair.



CPUE and total annual estimated catches for YBF in the Hauraki Gulf. Also shown is the fishing intensity (catch/CPUE), standardised relative to the geometric mean. Fishing year designated by second year of the pair.

FLATFISH (FLA)

Fishery and Stock Trends	
Recent Trend in Biomass or Proxy	In spite of fluctuations, both the Manukau and Kaipara series show a long-term declining trend. The Hauraki Gulf yellow belly CPUE index has fluctuated, peaking in 2006–07 at the highest point in the series and then declining steadily to 2015–16. However, there was a strong upturn in the final year of the series, with the 2016–17 index returning to above the series mean.
Recent Trend in Fishing Intensity or Proxy	Recent fishing intensity is relatively low in both of the west coast harbours while it sits near the series mean in the Hauraki Gulf series.
Other Abundance Indices	-
Trends in Other Relevant Indicators or Variables	-
Projections and Prognosis	
Stock Projections or Prognosis	Unknown
Probability of Current Catch or TACC causing Biomass to remain below or to decline below Limits	Soft Limit: Unknown Hard Limit: Unknown
Probability of Current Catch or TACC causing Overfishing to continue or to commence	Unknown

Assessment Methodology and Evaluation		
Assessment Type	Level 2 – Partial Quantitative Stock Assessment	
Assessment Method	Standardised CPUE based on positive catches	
Assessment Dates	Latest assessment: 2018	Next assessment: 2021
Overall assessment quality rank	1 – High Quality	
Main data inputs (rank)	Catch and effort data	1 – High Quality
Data not used (rank)	-	
Changes to Model Structure and Assumptions	-	
Major Sources of Uncertainty	- Uncertainty in the stock structure and relationship between CPUE and biomass	

Qualifying Comments
Work by NIWA (McKenzie et al 2013) in the Manukau harbour has linked the decrease in local CPUE with an increase in eutrophication, suggesting that there may be factors other than fishing contributing to the decline.
The lack of species specific reporting for FLA stocks is limiting the ability to assess these stocks, as is the possible reduction in carrying capacity for Manukau and Kaipara Harbours.


Fishery Interactions
Main bycatch is sand flounder, especially on the east coast. FLA 1 species are mostly targeted with setnets in harbours. Interactions with other species are currently being characterised.

- **Total FLA in Hauraki Gulf**

Because the Hauraki Gulf sand flounder CPUE series was rejected by the Northern Inshore Working Group, a total FLA CPUE analysis is substituted, which will be predominantly comprised of mixed sand flounder and yellow belly flounder.

Stock Status	
Year of Most Recent Assessment	2018
Assessment Runs Presented	Standardised CPUE for Hauraki Gulf
Reference Points	Target(s): Not established but B_{MSY} assumed Soft Limit: 20% B_0 Hard Limit: 10% B_0 Overfishing threshold: -
Status in relation to Target	Unknown
Status in relation to Limits	Unknown
Status in relation to Overfishing	Unknown

Historical Stock Status Trajectory and Current Status



CPUE and total annual estimated catches for FLA(TOT) in the Hauraki Gulf. Also shown is the fishing intensity (catch/CPUE), standardised relative to the geometric mean. Fishing year designated by second year of the pair.

Fishery and Stock Trends	
Recent Trend in Biomass or Proxy	The FLA(TOT) series shows an overall declining trend except for a three-year increase from 2002 to 2005 and a single strong increase in the final 2017 fishing year, which brings the series above the long-term average.
Recent Trend in Fishing Intensity or Proxy	Fishing intensity appears to be dropping after peaking in 2005
Other Abundance Indices	-
Trends in Other Relevant Indicators or Variables	-
Projections and Prognosis	
Stock Projections or Prognosis	Unknown

FLATFISH (FLA)

Probability of Current Catch or TACC causing Biomass to remain below or to decline below Limits	Soft Limit: Unknown Hard Limit: Unknown
Probability of Current Catch or TACC causing Overfishing to continue or to commence	Unknown

Assessment Methodology		
Assessment Type	Level 2 - Partial Quantitative Stock Assessment	
Assessment Method	Standardised CPUE based on positive catches	
Assessment Dates	Latest assessment: 2018	Next assessment: 2021
Overall assessment quality rank	1 – High Quality	
Main data inputs (rank)	Catch and effort data	1 – High Quality
Data not used (rank)	-	
Changes to Model Structure and Assumptions	-	
Major Sources of Uncertainty	- Uncertainty in the catch of sand flounder	

Qualifying Comments
The lack of species specific reporting for FLA stocks limits the ability to assess these stocks.

Fishery Interactions
Main QMS bycatch species is yellow belly flounder, especially on the east coast. FLA 1 species are mostly targeted with setnets in harbours. Interactions with other species are currently being characterised.


• FLA 2

Stock Structure Assumptions


Sand flounder off the East Coast (FMA2) of North Island appear to be a single continuous population. The stock structure of New Zealand sole (ESO) is unknown.

Stock Status	
Year of Most Recent Assessment	2018
Assessment Runs Presented	Standardised CPUE for all flatfish combined in FLA 2
Reference Points	Target: B_{MSY} -compatible proxy based on the mean CPUE 1989–90 to 2012–13 for the bottom trawl flatfish target series Soft Limit: 50% of target Hard Limit: 25% of target Overfishing threshold: F_{MSY}
Status in relation to Target	About as Likely as Not (40–60%) to be at or above the target
Status in relation to Limits	Soft Limit: Unlikely (< 40%) to be below Hard Limit: Very Unlikely (< 10%) to be below
Status in relation to Overfishing	Overfishing is Very Unlikely (< 10%) to be occurring

Historical Stock Status Trajectory and Current Status



Annual landings and standardised CPUE index based on positive catches for BT_FLA, (all flatfish species combined) at day resolution (Schofield et al 2018b). Fishing years are labelled according to the second calendar year e.g. 1990 = 1989–90. Horizontal lines are the target and the soft and hard limits.



Annual relative exploitation rate for flatfish in FLA 2.

Fishery and Stock Trends

Recent Trend in Biomass or Proxy	Relative abundance has fluctuated without trend since 1989–90 and is currently just below the target.
Recent Trend in Fishing Intensity or Proxy	Fishing intensity has trended down since the mid-1990s and is currently below the reference period (1990–2013) average
Other Abundance Indices	Tow based CPUE analysis for SFL and ESO from 2007–08 to 2016–17 data are reasonably consistent with the aggregated data index for combined species, although the decrease in abundance from 2016 to 2017 is more evident in ESO than SFL
Trends in Other Relevant Indicators or Variables	-

FLATFISH (FLA)

Projections and Prognosis	
Stock Projections or Prognosis	Stock is likely to continue to fluctuate around current levels
Probability of Current Catch or TACC causing Biomass to remain below or to decline below Limits	Soft Limit: Unknown for TACC; Unlikely (< 40%) for current catch Hard Limit: Unknown for TACC; Unlikely (< 40%) for current catch
Probability of Current Catch or TACC causing Overfishing to continue or to commence	Unknown for TACC; Unlikely (< 40%) for current catch

Assessment Methodology		
Assessment Type	Level 2 – Partial Quantitative Stock Assessment	
Assessment Method	Standardised CPUE based on positive catches	
Assessment Dates	Latest assessment: 2018	Next assessment: 2019
Overall assessment quality rank	1 – High Quality	
Main data inputs (rank)	- Catch and effort data	1 – High Quality
Data not used (rank)	N/A	
Changes to Model Structure and Assumptions	-	
Major Sources of Uncertainty	-	

Qualifying Comments
-

Fishery Interactions
The fishery is mainly confined to the inshore domestic trawl fleet except for a small incidental bycatch of soles, brill and turbot by offshore trawlers. The main fisheries landing flatfish as bycatch in FLA 2 target gurnard, snapper and trevally. Interactions with other species are currently being characterised.


- FLA 3 (all species combined)**

Stock Structure Assumptions


New Zealand sole and lemon sole appear to be a continuous population extending from Canterbury Bight to Foveaux Strait. Sand flounder off the East and South Coasts of South Island show localised concentrations that roughly correspond to the existing statistical areas. The stock relationships among these localised concentrations are unknown.

Stock Status	
Year of Most Recent Assessment	2018
Assessment Runs Presented	Standardised lognormal bottom trawl CPUE for all flatfish combined in FLA 3
Reference Points	Interim Target: B_{MSY} proxy based on the mean standardised lognormal CPUE from 1989–90 to 2006–07 (the final year of unconstrained catches) Soft Limit: 50% B_{MSY} proxy Hard Limit: 25% B_{MSY} proxy Overfishing threshold: F_{MSY} proxy
Status in relation to Target	About as Likely as Not (40–60%) to be at or above the target
Status in relation to Limits	Soft Limit: Unlikely (< 40%) to be below Hard Limit: Very Unlikely (< 10%) to be below
Status in relation to Overfishing	Unlikely (< 40%) that overfishing is occurring

Historical Stock Status Trajectory and Current Status



Standardised CPUE indices based on positive catches for all flatfish species combined, showing the agreed B_{MSY} proxy (green dashed line: average 1989–90 to 2006–07 CPUE index) and the associated Soft (purple dashed line) and Hard (grey dashed line) Limits (Starr & Kendrick in prep). Also shown are the QMR/MHR declared FLA 3 landings and the annual FLA 3 commercial catch allowance. Fishing year designated by second year of the pair.



Fishing intensity (catch/CPUE), standardised relative to the geometric mean, plot over time for FLA 3 (combined species). Also shown are the trajectory of total QMR/MHR catches (t) and the mean fishing intensity from 1989–90 to 2006–07 (green line). Fishing year designated by second year of the pair.

Fishery and Stock Trends

Recent Trend in Biomass or Proxy	CPUE has fluctuated over the long-term near the 25-year
----------------------------------	---

FLATFISH (FLA)

	mean.
Recent Trend in Fishing Intensity or Proxy	Fishing intensity has dropped since the reduction of the TACC in 2007–08 and the introduction of in-season variation to commercial catch allowance and remains below the F_{MSY} proxy.
Other Abundance Indices	-
Trends in Other Relevant Indicators or Variables	-

Projections and Prognosis

Stock Projections or Prognosis	Stock managed with annual in-season adjustment procedure: expected to vary in abundance around the long-term mean
Probability of Current Catch or TACC causing Biomass to remain below or to decline below Limits	Soft Limit: Unlikely (< 40%) for current catch and TACC Hard Limit: Very Unlikely (< 10%) for current catch and TACC
Probability of Current Catch or TACC causing Overfishing to continue or to commence	Unlikely (< 40%) for current catch and TACC

Assessment Methodology

Assessment Type	Level 2 – Partial Quantitative Stock Assessment	
Assessment Method	Standardised CPUE based on positive catches	
Assessment Dates	Latest assessment: 2018	Next assessment: 2019
Overall assessment quality rank	1 – High Quality	
Main data inputs (rank)	- Catch and effort data	1 – High Quality
Data not used (rank)	N/A	
Changes to Model Structure and Assumptions	-	
Major Sources of Uncertainty	<ul style="list-style-type: none"> - Mixed species complex managed without explicitly considering each species - Uncertainty in stock structure assumptions - The decline in fishing intensity in recent years is inconsistent with the increases for individual stock components 	

Qualifying Comments

The lack of historical species specific reporting for FLA stocks limits the ability to assess the long-term trends in these stocks; there is evidence that reporting by flatfish species has substantially improved in FLA 3 in 2012–13 and 2013–14.
--

Fishery Interactions

The fishery is mainly confined to the inshore domestic trawl fleet except for a small incidental bycatch of soles, brill and turbot by offshore trawlers. The main target species landing flatfish as bycatch in FLA 3 are red cod, barracouta, stargazer, gurnard, tarakihi and elephant fish. Interactions with other species are currently being characterised.
--


- FLA 3: New Zealand (ESO) sole

Stock Structure Assumptions

New Zealand sole appear to be a continuous population extending from Canterbury Bight to Foveaux Strait.


Stock Status	
Year of Most Recent Assessment	2015
Assessment Runs Presented	Standardised combined delta-lognormal bottom trawl CPUE for ESO in FLA 3, based on trips which landed FLA 3 but which did not use the FLA species code
Reference Points	Interim Target: B_{MSY} proxy based on mean standardised CPUE from 1990–91 to 2006–07 (the final year of unconstrained catches) Soft Limit: 50% B_{MSY} proxy Hard Limit: 25% B_{MSY} proxy Overfishing threshold: F_{MSY} proxy based on mean relative exploitation rate for the period 1989–90 to 2006–07
Status in relation to Target	Unlikely (< 40%) to be at or above target
Status in relation to Limits	Soft Limit: About as Likely as Not (40–60%) to be below Hard Limit: Unlikely (< 40%) to be below
Status in relation to Overfishing	Likely (> 60%) that overfishing is occurring

Historical Stock Status Trajectory and Current Status



Standardised CPUE indices based on combined delta-lognormal CPUE series for New Zealand sole (ESO), showing the agreed B_{MSY} proxy (green dashed line: average 1990–91 to 2006–07 CPUE index) and the associated Soft (purple dashed line) and Hard (grey dashed line) Limits (Starr & Kendrick in prep). Also shown is the ESO estimated catch by trips that landed FLA 3 but which did not use the FLA code. Fishing year designated by second year of the pair.

FLATFISH (FLA)



Fishery and Stock Trends

Recent Trend in Biomass or Proxy	CPUE has declined from a peak reached in 2001–02 and has been near the Soft Limit since 2010–11.
Recent Trend in Fishing Intensity or Proxy	Fishing intensity has increased since 2010–11 to more than 50% above the mean level.
Other Abundance Indices	-
Trends in Other Relevant Indicators or Variables	-

Projections and Prognosis

Stock Projections or Prognosis	-
Probability of Current Catch or TACC causing Biomass to remain below or to decline below Limits	Soft Limit: About as Likely as Not (40–60%) for current catch Hard Limit: Unlikely (< 40%) for current catch
Probability of Current Catch or TACC causing Overfishing to continue or to commence	Likely (> 60%) for current catch

Assessment Methodology and Evaluation

Assessment Type	Level 2 – Partial Quantitative Stock Assessment	
Assessment Method	Standardised CPUE based on positive catches	
Assessment Dates	Latest assessment: 2015	Next assessment: 2020
Overall assessment quality rank	1 – High Quality	
Main data inputs (rank)	- Catch and effort data	1 – High Quality
Data not used (rank)	N/A	
Changes to Model Structure and Assumptions	-	
Major Sources of Uncertainty	- uncertainty in stock structure assumptions	

Qualifying Comments

The lack of historic species specific reporting for FLA stocks limits the ability to assess the long-term trends in these stocks; there is evidence that reporting by flatfish species has substantially improved in FLA 3 in 2012–13 and 2013–14.

Fishery Interactions

The fishery is mainly confined to the inshore domestic trawl fleet except for a small incidental bycatch of soles, brill and turbot by offshore trawlers. The main target species landing flatfish as bycatch in FLA 3 are red cod, barracouta, stargazer, gurnard, tarakihi and elephant fish. Interactions with other species are currently being characterised.


- **FLA 3: Lemon (LSO) sole**

Stock Structure Assumptions


Lemon sole appear to be a continuous population extending from Canterbury Bight to Foveaux Strait.

Stock Status	
Year of Most Recent Assessment	2015
Assessment Runs Presented	Standardised combined delta-lognormal bottom trawl CPUE for LSO in FLA 3, based on trips which landed FLA 3 but which did not use the FLA species code
Reference Points	Interim Target: B_{MSY} proxy based on mean standardised CPUE from 1990–91 to 2006–07 (the final year of unconstrained catches) Soft Limit: 50% B_{MSY} proxy Hard Limit: 25% B_{MSY} proxy Overfishing threshold: F_{MSY} proxy based on mean relative exploitation rate for the period 1989–90 to 2006–07
Status in relation to Target	About as Likely as Not (40–60%) to be at or above target
Status in relation to Limits	Soft Limit: Unlikely (< 40%) to be below Hard Limit: Very Unlikely (< 10%) to be below
Status in relation to Overfishing	Likely (> 60%) that overfishing is occurring


Historical Stock Status Trajectory and Current Status



Standardised CPUE indices based on combined delta-lognormal CPUE series for Lemon sole (LSO), showing the agreed B_{MSY} proxy (green dashed line: average 1990–91 to 2006–07 CPUE index) and the associated Soft (purple dashed line) and Hard (grey dashed line) Limits (Starr & Kendrick in prep). Also shown is the LSO estimated catch by trips that landed FLA 3 but which did not use the FLA code. Fishing year designated by second year of the pair.



Fishing intensity (catch/CPUE, standardised relative to the geometric mean) plot over time for Lemon sole (LSO) in FLA 3. Also shown are the trajectory of LSO estimated catches by trips that landed FLA 3 but which did not use the FLA code and the mean fishing intensity from 1990–91 to 2006–07 (green line). Fishing year designated by second year of the pair.



Each relative series scaled so that the geometric mean=1.0 from 1991 to 1994, 1996, 2007 to 2009, 2012, 2014

Standardised CPUE indices based on combined delta-lognormal CPUE series for Lemon sole (ESO), shown with the 10 trawl survey LSO biomass indices from the Kaharoa ECSI winter trawl survey. Fishing year designated by second year of the pair.

Fishery and Stock Trends	
Recent Trend in Biomass or Proxy	CPUE reached a nadir in 2003–04, but then climbed to a high level in 2007–08 and has since declined to the long-term mean level.
Recent Trend in Fishing Intensity or Proxy	Fishing intensity has fluctuated, mostly above the F_{MSY} proxy since 1994–95, and in 2013–14 was nearly 40% above this level.
Other Abundance Indices	Relative abundance from the ECSI trawl survey has fluctuated without trend since 1991.
Trends in Other Relevant Indicators or Variables	-

Projections and Prognosis	
Stock Projections or Prognosis	-
Probability of Current Catch or TACC causing Biomass to remain below or to decline below Limits	Soft Limit: Unlikely (< 40%) Hard Limit: Very Unlikely (< 10%)
Probability of Current Catch or TACC causing Overfishing to continue or to commence	Likely (> 60%)

Assessment Methodology and Evaluation		
Assessment Type	Level 2 – Partial Quantitative Stock Assessment	
Assessment Method	Standardised CPUE based on positive catches	
Assessment Dates	Latest assessment: 2015	Next assessment: 2020
Overall assessment quality rank	1 – High Quality	
Main data inputs (rank)	- Catch and effort data	1 – High Quality
Data not used (rank)	N/A	
Changes to Model Structure and Assumptions	-	
Major Sources of Uncertainty	- uncertainty in stock structure assumptions	
Qualifying Comments		
The lack of historic species specific reporting for FLA stocks limits the ability to assess the long-term trends in these stocks; there is evidence that that reporting by flatfish species has substantially improved in FLA 3 in 2012–13 and 2013–14.		

Fishery Interactions
The fishery is mainly confined to the inshore domestic trawl fleet except for a small incidental bycatch of soles, brill and turbot by offshore trawlers. The main target species landing flatfish as bycatch in FLA 3 are red cod, barracouta, stargazer, gurnard, tarakihi and elephant fish. Interactions with protected species are believed to be low. Incidental captures of seabirds occur.

- **FLA 3: Sand Flounder (SFL)**

Stock Structure Assumptions


Sand flounder off the East and South Coasts of South Island show localised concentrations that roughly correspond to the existing statistical areas. The stock relationships among these localised concentrations are unknown.

Stock Status	
Year of Most Recent Assessment	2015
Assessment Runs Presented	Standardised combined delta-lognormal bottom trawl CPUE for SFL in FLA 3, based on trips which landed FLA 3 but which did not use the FLA species code
Reference Points	Interim Target: B_{MSY} proxy based on mean standardised CPUE from 1990–91 to 2006–07 (the final year of unconstrained


FLATFISH (FLA)

	catches) Soft Limit: 50% B_{MSY} proxy Hard Limit: 25% B_{MSY} proxy Overfishing threshold: F_{MSY} proxy based on mean relative exploitation rate for the period 1989–90 to 2006–07
Status in relation to Target	Very Likely (> 90%) to be at or above target
Status in relation to Limits	Soft Limit: Very Unlikely (< 10%) to be below Hard Limit: Very Unlikely (< 10%) to be below
Status in relation to Overfishing	About as Likely as Not (40–60%) that overfishing is occurring

Historical Stock Status Trajectory and Current Status



Standardised CPUE indices based on combined delta-lognormal CPUE series for Sand flounder (SFL), showing the agreed B_{MSY} proxy (green dashed line: average 1990–91 to 2006–07 CPUE index) and the associated Soft (purple dashed line) and Hard (grey dashed line) Limits (Starr & Kendrick in prep). Also shown is the SFL estimated catch by trips that landed FLA 3 but which did not use the FLA code. Fishing year designated by second year of the pair.



Fishing intensity (catch/CPUE, standardised relative to the geometric mean) plot over time for Sand flounder (SFL) in FLA 3. Also shown are the trajectory of SFL estimated catches by trips that landed FLA 3 but which did not use the FLA code and the mean fishing intensity from 1990–91 to 2006–07 (green line). Fishing year designated by second year of the pair.

Fishery and Stock Trends	
Recent Trend in Biomass or Proxy	CPUE has been climbing steadily from a nadir in 2003–04.
Recent Trend in Fishing Intensity or Proxy	Fishing intensity dropped to relatively low levels in the late 2000s, and has since climbed back to the level of the F_{MSY} proxy
Other Abundance Indices	-
Trends in Other Relevant Indicators or Variables	-

Projections and Prognosis	
Stock Projections or Prognosis	-
Probability of Current Catch or TACC causing Biomass to remain below or to decline below Limits	Soft Limit: Very Unlikely (< 10%) for current catch Hard Limit: Very Unlikely (< 10%) for current catch
Probability of Current Catch or TACC causing Overfishing to continue or to commence	Unknown

Assessment Methodology and Evaluation		
Assessment Type	Level 2 – Partial Quantitative Stock Assessment	
Assessment Method	Standardised CPUE based on positive catches	
Assessment Dates	Latest assessment: 2015	Next assessment: 2020
Overall assessment quality rank	1 – High Quality	
Main data inputs (rank)	- Catch and effort data	1 – High Quality
Data not used (rank)	N/A	
Changes to Model Structure and Assumptions	-	
Major Sources of Uncertainty	- uncertainty in stock structure assumptions	

Qualifying Comments
The lack of historic species specific reporting for FLA stocks limits the ability to assess the long-term trends in these stocks; there is evidence that reporting by flatfish species has substantially improved in FLA 3 in 2012–13 and 2013–14.


Fishery Interactions
The fishery is mainly confined to the inshore domestic trawl fleet except for a small incidental bycatch of soles, brill and turbot by offshore trawlers. The main target species landing flatfish as bycatch in FLA 3 are red cod, barracouta, stargazer, gurnard, tarakihi and elephant fish. Interactions with other species are currently being characterised.

6. FOR FURTHER INFORMATION

- Beentjes, M P (2003) Review of flatfish catch data and species composition. *New Zealand Fisheries Assessment Report 2003/17*. 22 p.
- Bentley, N. (2010). INS 2008-02: Approaches for determining in season TAC increases for 2nd Schedule stocks Draft FAR for FLA 3. Document 2010/unknown. 23 p. (Unpublished report held by Fisheries New Zealand, Wellington.)
- Bentley, N. (2011). Start-of-season CPUE and in-season adjustment of TACC for FLA 3 in 2010/2011. Document 2011/unknown. 21 p. (Unpublished report held by Fisheries New Zealand, Wellington.)
- Bentley, N. (2012). Start-of-season CPUE and in-season adjustment of TACC for FLA 3 in 2011/2012. Document 2012/unknown. 12 p. (Unpublished report held by Fisheries New Zealand, Wellington.)
- Bentley, N. (2015). Start-of-season CPUE and in-season adjustment of TACC for FLA 3 in 2014/2015. SINSWG 2015-05A. 15 p. (Unpublished report held by Fisheries New Zealand, Wellington.)
- Boyd, R O; Reilly, J L (2005) 1999/2000 national marine recreational fishing survey: harvest estimates. Draft New Zealand Fisheries Research Report. (Unpublished report held by Fisheries New Zealand, Wellington.)
- Bradford, E (1998) Harvest estimates from the 1996 national recreational fishing surveys. New Zealand Fisheries Assessment Research Document 1998/16. 27 p. (Unpublished report held by NIWA library, Wellington.)
- Brouwer, S. (2013). Start-of-season CPUE and in-season adjustment of TACC for FLA 3 in 2012/2013. SINSWG 2013-10. 14 p. (Unpublished report held by Fisheries New Zealand, Wellington.)
- Coburn, R P; Beentjes, M P (2005) Abundance estimates for flatfish in FLA 1 from standardized catch per unit effort analysis of the set net fisheries, 1989–90 to 2003–04. *New Zealand Fisheries Assessment Report 2005/57*. 46 p.
- Colman, J A (1972) Size at first maturity of two species of flounders in the Hauraki Gulf, New Zealand. *New Zealand Journal of Marine and Freshwater Research* 6(3): 240–245.
- Colman, J A (1973) Spawning and fecundity of two flounder species in the Hauraki Gulf, New Zealand. *New Zealand Journal of Marine and Freshwater Research* 7(1 & 2): 21–43.
- Colman, J A (1974) Movements of flounders in the Hauraki Gulf, New Zealand. *New Zealand Journal of Marine and Freshwater Research* 8(1): 79–93.
- Colman, J A (1978) Tagging experiments on the sand flounder, *Rhombosolea plebeia* (Richardson), in Canterbury, New Zealand, 1964 to 1966. *Fisheries Research Bulletin* 18: 42 p.
- Colman, J A (1985) Flatfish. In: Colman, J A; McKoy, J L; Baird, G G (Comps. and Eds.). Background papers for the 1985 Total Allowable Catch recommendations 74–78. (Unpublished report, held in NIWA library, Wellington.)
- Hartill, B (2004) Characterisation of the commercial flatfish, grey mullet, and rig fisheries in the Kaipara Harbour. *New Zealand Fisheries Assessment Report 2004/1*. 23 p.
- Hartill, B; Davey, N (2015) Mean weight estimates for recreational fisheries in 2011–12. *New Zealand Fisheries Assessment Report 2015/25*.
- Hoenig, J M (1983) Empirical use of longevity data to estimate mortality rates. *Fishery Bulletin* 81: 898–903.
- Kirk, P D (1988) Flatfish. New Zealand Fisheries Assessment Research Document 1988/13. 16 p. (Unpublished document held by NIWA library, Wellington.)
- Kendrick, T H; Bentley, N (2011). Fishery characterisation and setnet catch-per-unit-effort indices for flatfish in FLA 1; 1989–90 to 2007–08. *New Zealand Fisheries Assessment Report 2011/3*. 74p.
- Kendrick, T; Bentley, N (2012) Fishery characterisation and setnet catch-per-unit-effort indices for flatfish in FLA 1; 1989–90 to 2010–11. New Zealand. *New Zealand Fisheries Assessment Report 2012/32*. 88 p.
- Kendrick T.H., Bentley, N (2015). Report to the Northern Inshore Fishery Assessment Working Group: Fishery characterisation and setnet catch-per-unit-effort indices for flatfish in FLA 1; 1989–90 to 2013–14. Document 2015/23, 81 p. (Unpublished document held by Fisheries New Zealand, Wellington, N.Z.)
- Langley, A.D. (2014). Updated CPUE analyses for selected South Island inshore finfish stocks. *New Zealand Fisheries Assessment Report 2014/40*. 116 p.
- McGregor, G; Coakley, M; Warren, E (Unpublished). Review of the conversion factor measurement for flatfish (gutted). Draft Working report. (Unpublished Working Group paper held by Fisheries New Zealand, Wellington.)
- McKenzie, J R; Parsons, D M; Bian, R (2013) Can juvenile yellow belly and sand flounder abundance indices and environmental variables predict adult abundance in the Manukau and Mahurangi Harbours? *New Zealand Fisheries Assessment Report 2013/10*. 31 p.
- Mundy, A R (1968) A study of the biology of the sand flounder, *Rhombosolea plebeia* (Richardson), off the Canterbury coast. (Unpublished Ph.D. thesis lodged in University of Canterbury library, Christchurch, New Zealand.)
- Schofield, M.I.; Langley, A.D.; Middleton, D.A.J. (2018a). Fisheries characterisation and catch-per unit-effort analyses FLA 2. New Zealand Fisheries Assessment Report 2018/04. 61 p
- Schofield, M.I.; Langley, A.D.; Middleton, D.A.J. (2018b). Catch-per unit-effort (CPUE) update for FMA 2 flatfish (FLA 2). Report for Fisheries Inshore New Zealand. <http://www.mpi.govt.nz/news-and-resources/publications/>
- Starr, P.J.; Kendrick, T.H. (2017). Prediction of 2016/17 FLA 3 CPUE and Operation Of FLA 3 Management Procedure–February Update. SINSWG-2017/7. 5 p. (Unpublished document held by Fisheries New Zealand, Wellington N.Z.)
- Starr, P.J.; Kendrick, T.H. (2018). Prediction of 2017/18 FLA 3 CPUE and Operation of FLA 3 Management Procedure. SINSWG-2018/06. 23 p. (Unpublished document held by Fisheries New Zealand, Wellington N.Z.)
- Starr, P.J.; Kendrick, T.H. (in prep.). FLA 1 Fishery Characterisation and CPUE. New Zealand Fisheries Assessment Report 2018/XX. xxx p.
- Starr, P J; Kendrick, T H; Bentley, N; Langley, A D (in prep.). Development of a TACC in-season increase procedure for FLA 3. Draft New Zealand Fisheries Assessment Report.
- Starr, P.J.; Kendrick, T.H.; Bentley, N. (2016). Prediction of 2016/17 FLA 3 CPUE and Operation of FLA 3 Management Procedure. SINSWG-2016/5A. 22 p. (Unpublished document held by Fisheries New Zealand, Wellington N.Z.)
- Stevens, D W; Francis, M P; Shearer, P J; McPhee, R P; Hickman, R W; Tait, M (2001) Age and growth of brill (*Colistium guntheri*) and turbot (*C. nudipinnis*) from the west coast South Island. Final research report for Ministry of Fisheries research project FLA2000/01. 35p. (Unpublished report held by Fisheries New Zealand, Wellington.)
- Sutton, C P; MacGibbon, D J; Stevens, D W (2010) Age and growth of greenback flounder (*Rhombosolea tapirina*) from southern New Zealand. *New Zealand Fisheries Assessment Report 2010/48*. 15 p
- Teirney, L D; Kilner, A R; Millar, R E; Bradford, E; Bell, J D (1997) Estimation of recreational catch from 1991/92 to 1993/94 New Zealand. Fisheries Assessment Research Document 1997/15. 43 p. (Unpublished report held by NIWA library, Wellington.)
- Wynne-Jones, J; Gray, A; Hill, L; Heinemann, A (2014) National Panel Survey of Marine Recreational Fishers 2011–12: Harvest Estimates. *New Zealand Fisheries Assessment Report 2014/67*.

FRESHWATER EELS (SFE, LFE, ANG)

(*Anguilla australis*, *Anguilla dieffenbachii*, *Anguilla reinhardtii*)



1. FISHERY SUMMARY

1.1 Commercial fisheries

The freshwater eel fishery is distributed throughout accessible freshwaters (lakes, rivers, streams, farm ponds, tarns) and some estuarine and coastal waters of New Zealand, including the Chatham Islands. The contemporary commercial fishery dates from the mid-1960s when markets were established in Europe and Asia.

The New Zealand eel fishery is based on the two temperate species of freshwater eels occurring in New Zealand, the shortfin eel *Anguilla australis* and the longfin eel *A. dieffenbachii*. A third species of freshwater eel, the Australasian longfin (*A. reinhardtii*), identified in 1996, has been confirmed from North Island landings. The proportion of this species in landings is unknown but is thought to be small. Virtually all eels (98%) are caught with fyke nets. Eel catches are greatly influenced by water temperature, flood events (increased catches) and drought conditions (reduced catches). Catches decline in winter months (May to September), particularly in the South Island where fishing ceases.

The South Island eel fishery was introduced into the Quota Management System (QMS) on 1 October 2000 with shortfin and longfin species combined into six fish stocks (codes ANG 11 to ANG 16). The Chatham Island fishery was introduced into the QMS on 1 October 2003 with two fish stocks (shortfins and longfins separated into SFE 17 and LFE 17, respectively). The North Island eel fishery was introduced into the QMS on 1 October 2004 with eight fish stocks (four longfin stocks LFE 20–23 and four shortfin stocks SFE 20–23). On 1 October 2017 the former South Island ANG QMAs were split into corresponding longfin (LFE 11–16) and shortfin (SFE 11–16) QMAs, each with its own TACC. The Australasian longfin eel is combined as part of the shortfin eel stocks in the Chatham and North Islands, as this species has productivity characteristics closer to shortfins than longfins, and because the catch is not sufficient to justify its own separate stocks. The occasional catch of Australasian longfins is mainly confined to the upper North Island.

The fishing year for all stocks extends from 1 October to 30 September except for ANG 13 (Te Waihora/Lake Ellesmere) which has a fishing year from 1 February to 31 January (since 2002). Currently, there exist minimum and maximum commercial size limits for both longfins and shortfins

FRESHWATER EELS (SFE, LFE, ANG)

(220 g and 4 kg, respectively) throughout New Zealand. North Island quota owners agreed in August 2012 to use 31 mm escapement tubes (equivalent to South Island regulation). The minimum legal diameter for escape tubes on the North Island was increased to 31 mm in October 2013. Quota owners from both islands formally agreed in 1995–96 not to land migratory female longfin eels. In the South Island the eel industry agreed to voluntary incremental increases in the diameter of escape tubes in fyke nets which increased from 25 mm to 26 mm in 1990–91, to 27 mm in 1993–94, to 28.5 mm in 1994–95, and finally to 31 mm in 1997–98, which effectively increases the minimum size limit of both main species to about 300 g. Since about 2006 there has been a voluntary code of practise to return all longfin eels caught in Te Waihora; catches of these longfins are recorded on Eel Catch Effort Returns (ECERs), but not on the Eel Catch Landing Returns (ECLRs).

In early 2005 the Mohaka, Motu and much of the Whanganui River catchments were closed to commercial fishing and there are a number of smaller areas elsewhere that have been reserved as customary fisheries (see Section 1.3). In addition, all Public Conservation lands managed by the Department of Conservation require at a minimum a concession to be commercially fished and in most cases are closed to commercial fishing. In the Waikato-Tainui rohe (region), fisheries bylaws were introduced in March 2014 to limit the minimum harvest size to 300 g for SFE and 400 g for LFE. Amongst other things, these bylaws also introduced an upper limit of 2 kg for both species (to prevent the taking of longfin females that are in a migratory state) and added seasonal closures in some reaches.

Commercial catch data are available from 1965 and originate from different sources. Catch data prior to 1988 are for calendar years, whereas those from 1988 onwards are for fishing years (Table 1, Figure 1). Licensed Fish Receiver Returns (LFRRs), Quota Management Reports (QMRs), and Monthly Harvest Returns (MHRs) provide the most accurate data on landings over the period 1988–89 to 2015–16 for the whole of New Zealand.

Table 1: Eel catch data (t) from for calendar years 1965 to 1988 and fishing years 1988–89 to 2016–17 based on MAF Fisheries Statistics Unit (FSU) and Licensed Fish Receiver Returns (LFRR), Quota Management Reports (QMR), and Monthly Harvest Returns (MHR).

Year	Landings	Year	Landings	Year	Landings	Year	Landings
1965	30	1980	1 395	1994–95	1 438	2009–10	560
1966	50	1981	1 043	1995–96	1 429	2010–11	626
1967	140	1982	872	1996–97	1 342	2011–12	755
1968	320	1983	1 206	1997–98	1 210	2012–13	717
1969	450	1984	1 401	1998–99	1 219	2013–14	678
1970	880	1985	1 505	1999–00	1 133	2014–15	547
1971	1 450	1986	1 166	2000–01	1 071	2015–16	455
1972	2 077	1987	1 114	2001–02	978	2016–17	511
1973	1 310	1988	1 281	2002–03	808		
1974	860	1988–89	1 315	2003–04	729		
1975	1 185	1989–90	1 356	2004–05	708		
1976	1 501	1990–91	1 590	2005–06	771		
1977	906	1991–92	1 585	2006–07	718		
1978	1 583	1992–93	1 466	2007–08	660		
1979	1 640	1993–94	1 255	2008–09	518		

MAF data, 1965–1982; FSU, 1983 to 1989–90; CELR, 1990–91 to 1999–00; ECLR 2000–01 to 2003–04; MHR 2004–05–present.




Figure 1: Total eel landings from 1965 to 2016–17, as well as separate shortfin and longfin landings from 1989–90 to 2016–17. Prior to 1988–89, the data points represent estimates for the period prior to the introduction of Eel Catch Landing Return (ECLR) forms, and were generated by pro-rating the unidentified eel catch by the LFE:SFE ratio (see below).

There was a rapid increase in commercial catches during the late 1960s, with catches rising to a peak of 2077 t in 1972. Landings were relatively stable from 1983 to 2000, a period when access to the fishery was restricted, although overall catch limits were not in place. In 2000–01 landings dropped to 1070 t, and these were further reduced during 2001–02 to 2004–05 as eel stocks were progressively introduced into the Quota Management System (QMS). While landings since 2007–08 were further affected by the reduction in TACCs for both species in the North Island on 1 Oct. 2007, eel landings have remained below the TACCs as a result of reduced international market demand and ACE shelving by some iwi, and since 2007–08 have ranged between 487 and 642 tonnes. For the period 1991–92 to 2013–14, the North Island provided on average 61% of the total New Zealand eel catch (Table 2).

In 2016, South Island eel stocks (ANG 11–16) were separated into individual shortfin (SFE 11–16) and longfin (LFE 11–16) stocks. The new stocks utilise the same geographical areas as the preexisting stocks (ANG 11–16), but were separated to allow species specific management of the individual eel species. After the stocks were separated new catch limits and allowances were set. For the SFE stocks the new TACs were based on the highest historical catch, apart from SFE 13, which received a 10% increase as the CPUE index was well above the target. For LFE stocks, the TAC was reduced to a point that effectively eliminated commercial targeting (a TAC close to zero) for four of the six stocks (LFE 11, 12, 13 and 14). For the remaining two LFE stocks (LFE 15 and 16), TACs allow continued commercial utilisation, but at significantly reduced levels. The separated stocks and their associated catch limits and allowances came into force on 1 October 2016 for SFE/LFE 11, 12, 14, 15 and 16 and 1 Feb 2017 for SFE/LFE 2017.

Table 2: North and South Island eel catch (t) compiled from data from individual processors 1991–92 to 1999–00 and LFRR/QMR/MHR 2000–01 to 2015–16. Numbers in parentheses represent the percentage contribution from the North Island fishery.

Fishing year	North Island	South Island	Total individual processors	LFRR/QMR/MHR (excluding Chatham Islands)	Total NZ
1991–92	989	631	1 621 (61%)		–
1992–93	865	597	1 462 (59%)		–
1993–94	744	589	1 334 (56%)		–
1994–95	1 004	510	1 515 (66%)		–
1995–96	962	459	1 481 (65%)		–
1996–97	830	418	1 249 (66%)		–
1997–98	795	358	1 153 (69%)		–
1998–99	804	381	1 185 (68%)		–
1999–00	723	396	1 119 (65%)		–
2000–01	768	303	–		1 071 (72%)
2001–02	644	319	–		962 (67%)

FRESHWATER EELS (SFE, LFE, ANG)

Table 2: [Continued]

Fishing year	North Island	South Island	Total individual processors	LFRR/QMR/MHR (excluding Chatham Islands)	Total NZ
2002–03	507	296	—	—	803 (63%)
2003–04	454	282	—	—	737 (62%)
2004–05	426	285	—	—	712 (60%)
2005–06	497	285	—	—	781 (64%)
2006–07	440	278	—	—	718 (61%)
2007–08	372	288	—	—	660 (56%)
2008–09	303	215	—	—	517 (59%)
2009–10	318	242	—	—	560 (57%)
2010–11	330	296	—	—	626 (53%)
2011–12	418	337	—	—	755 (55%)
2012–13	364	353	—	—	717 (51%)
2013–14	367	311	—	—	678 (54%)
2014–15	306	241	—	—	547 (56%)
2015–16	254	201	—	—	455 (56%)

Table 3: Total NZ eel landings (t) by species and fishing year. Numbers in bold represent data collected following the introduction of the ECLR forms, whereas all others are pro-rated as described above. Numbers in parentheses represent the longfin proportion of total landings.

Fishing year	Shortfin (SFE)	Longfin (LFE)	Total landings
1989–90	617	453	1 069 (42%)
1990–91	808	616	1 424 (43%)
1991–92	941	612	1 553 (39%)
1992–93	872	741	1 613 (46%)
1993–94	692	588	1 279 (46%)
1994–95	909	588	1 497 (39%)
1995–96	977	518	1 495 (35%)
1996–97	841	465	1 307 (36%)
1997–98	881	442	1 323 (33%)
1998–99	824	434	1 258 (34%)
1999–00	741	413	1 154 (36%)
2000–01	698	388	1 086 (36%)
2001–02	660	360	1 020 (35%)
2002–03	560	279	839 (33%)
2003–04	510	216	726 (30%)
2004–05	460	254	713 (36%)
2005–06	553	226	774 (29%)
2006–07	520	210	730 (29%)
2007–08	470	196	666 (29%)
2008–09	424	95	519 (18%)
2009–10	441	114	555 (20%)
2010–11	440	159	599 (26%)
2011–12	515	237	752 (32%)
2012–13	491	230	721 (32%)
2013–14	475	201	676 (30%)
2014–15	434	116	550 (21%)
2015–16	378	89	467 (19%)

Prior to the 2000–01 fishing year, three species codes were used to record species landed, SFE (shortfin), LFE (longfin) and EEU (eels unidentified). A high proportion of eels (46% in 1990–91) were identified as EEU between the fishing years 1989–90 and 1998–99. Pro-rating the EEU catch by the ratio of LFE : SFE by fishing year provides a history of landings by species (Table 3), although it should be noted that pro-rated catches prior to 1999–00 are influenced by the high proportion of EEU from some eel statistical areas (e.g., Waikato) and therefore may not provide an accurate species breakdown. The introduction of the new Eel Catch Landing Return (ECLR) form in 2001–02 improved the species composition information, as the EEU code was not included. There was a gradual decline in the proportion of longfin eels in landings, from over 40% in 1989–90 to about 30% in 2007–08, followed by a marked drop to 18% in 2008–09 (Table 3). The proportion of longfins in the catch then gradually

increased and was about 30% of the total in 2013–14. Several factors have contributed to the pattern in the proportion of longfin eels, including: declining abundance in the early part of the series; reduced quotas; the closure of some catchments to commercial fishing; and declining/fluctuating market demand.

The species proportion of the landings varies by geographical area. From analyses of landings to eel processing factories and estimated catch from ECLRs, longfins are the dominant species in most areas of the South Island, except for a few discrete locations such as lakes Te Waihora (Ellesmere) and Brunner, and the Waipori Lakes, where shortfins dominate landings. Shortfins are dominant in North Island landings. The shortfin eel catches are mostly comprised of pre-migratory female feeding eels, with the exception of Te Waihora (Lake Ellesmere), where significant quantities of seaward migrating male shortfin eels (under 220 g) are taken during the period of February to March.

Table 4: TACCs and commercial landings (t) for South Island eel stocks (based on ECLR data).

Fishing Year	ANG11		ANG12		ANG13		ANG14		ANG15		ANG16		Total landings
	TACC	Landings	TACC	Landings	TACC	Landings	TACC	Landings	TACC	Landings	TACC	Landings	
Shortfin Eel (SFE)													
2000–01	40	4.5	43	4.4	122	102.2	35	6.1	118	19.4	63	9.8	146.6
2001–02	40	18.9	43	5.7	122	63.6*	35	10.1	118	20.2	63	20.2	83.8
2002–03	40	19.2	43	5.9	122	95.4	35	9.9	118	11.7	63	4.5	146.7
2003–04	40	8.7	43	4.8	122	118.2	35	7.5	118	13.0	63	9.4	161.8
2004–05	40	2.7	43	1.4	122	121.3	35	5.7	118	1.5	63	9.6	156.0
2005–06	40	9.0	43	4.3	122	119.9	35	7.4	118	12.0	63	11.2	164.0
2006–07	40	10.9	43	6.3	122	121.5	35	4.4	118	15.4	63	16.5	175.2
2007–08	40	8.5	43	1.2	122	119.7	35	5.8	118	21.2	63	11.5	167.9
2008–09	40	4.7	43	<1	122	123.0	35	1.8	118	16.6	63	19.7	166.0
2009–10	40	3.8	43	5.8	122	97.3	35	3.9	118	29.1	63	30.3	170.2
2010–11	40	10.0	43	6.9	122	89.3	35	3.7	118	19.4	63	19.9	149.2
2011–12	40	8.8	43	10.8	122	113.3	35	7.3	118	21.4	63	13.1	174.8
2012–13	40	7.6	43	19.9	122	125.0	35	2.6	118	16.7	63	22.8	194.6
2013–14	40	3.4	43	16.5	122	119.3	35	2.5	118	11.7	63	16.8	170.2
2014–15	40	2.8	43	13.6	122	112.1	35	1.3	118	14.4	63	11.8	156.0
2015–16	40	<1	43	0	122	109.9	35	<1	118	22.7	63	10.2	144.4
New FMA	SFE11		SFE 12		ANG 13		SFE 14		SFE 15		SFE 16		Total
2016–17	19	0	20	0.2	134.12	132.8	10	0	29	20.7	30	12.97	166.7
Longfin Eel (LFE)													
2000–01	40	10.6	43	22.6	122	2.1	35	12.6	118	63.6	63	28.4	140.1
2001–02	40	16.4	43	15.6	122	1.0*	35	6.0	118	80.5	63	30.2	150.1
2002–03	40	10.6	43	10.1	122	1.4	35	10.0	118	73.0	63	27.2	132.6
2003–04	40	2.8	43	2.7	122	<1	35	10.2	118	64.7	63	21.2	102.9
2004–05	40	2.8	43	3.4	122	<1	35	2.3	118	79.6	63	34.4	123.7
2005–06	40	6.0	43	9.8	122	<1	35	6.4	118	61.1	63	21.1	105.5
2006–07	40	4.4	43	1.7	122	<1	35	7.0	118	65.0	63	32.8	112.1
2007–08	40	11.9	43	6.5	122	<1	35	7.4	118	73.0	63	23.1	122.9
2008–09	40	1.4	43	<1	122	0	35	2.3	118	33.7	63	13.2	51.0
2009–10	40	8.0	43	<1	122	<1	35	3.2	118	40.0	63	15.3	68.0
2010–11	40	13.1	43	6.1	122	<1	35	6.7	118	73.9	63	14.1	114.9
2011–12	40	11.2	43	11.0	122	2.0	35	18.4	118	85.4	63	27.6	155.7
2012–13	40	15.6	43	7.6	122	<1	35	22.3	118	88.6	63	30.4	164.5
2013–14	40	14.0	43	6.1	122	<1	35	10.7	118	77.9	63	29.3	138.5
2014–15	40	2.5	43	3.7	122	0	35	2.1	118	56.3	63	15.3	79.9
2015–16	40	<1	43	0	122	0	35	4.5	118	43.0	63	10.5	59.0
New FMA	LFE11		LFE 12		ANG 13		LFE 14		LFE 15		LFE 16		Total
2016–17	1	0	1	<1	1	0	1	0	52	33.4	25	14.1	47.5

*For the transition from a 1 October to 1 February fishing year, an interim TACC of 78 t was set for the period 1 October 2001 to 31 January 2002. From January 2002 the Te Waihora (Lake Ellesmere) fishing year was 1 February to 31 January. Fishing year for all other areas is 1 October to 30 September.

FRESHWATER EELS (SFE, LFE, ANG)

Table 5: TACCs and commercial landings (t) for Chatham Island (SFE 17) and North Island shortfin stocks from 2003–04 to 2016–17 (based on ECLR data).

Fishing Year	SFE 17		SFE 20		SFE 21		SFE 22		SFE 23		Total landings
	TACC	Landings	TACC	Landings	TACC	Landings	TACC	Landings	TACC	Landings	
2003–04	10	0.7	-	-	-	-	-	-	-	-	0.7
2004–05	10	1.3	149	78.4	163	123.0	108	80.5	37	15.0	298.1
2005–06	10	2.7	149	93.3	163	144.3	108	106.9	37	31.5	378.6
2006–07	10	0.0	149	107.8	163	113.5	108	91.3	37	30.2	342.8
2007–08	10	0.0	86	76.0	134	125.3	94	82.5	23	15.8	299.5
2008–09	10	0.0	86	66.8	134	110.0	94	70.9	23	10.3	258.0
2009–10	10	0.0	86	60.2	134	124.1	94	68.5	23	17.5	270.3
2010–11	10	0.0	86	85.5	134	133.9	94	58.8	23	16.1	294.3
2011–12	10	0.0	86	85.6	134	140.9	94	95.7	23	18.8	341.0
2012–13	10	0.0	86	78.8	134	124.3	94	82.0	23	14.7	299.8
2013–14	10	0.0	86	71.6	134	139.2	94	82.1	23	14.5	307.4
2014–15	10	0.0	86	63.8	134	122.8	94	73.3	23	13.7	273.6
2015–16	10	0.0	86	53.8	134	119.1	94	49.2	23	10.4	232.5
2016–17	10	0.0	86	46.2	134	123.4	94	81.3	23	13.0	263.9

The Total Allowable Commercial Catch (TACC) and reported commercial landings by species for the South Island eel stocks are shown in Table 4 from 2000–01 (when eels were first introduced into the QMS) to 2016–17. The annual landings are based on data recorded on ECLR forms, as the MHR forms report QMA catches for the two species combined.

The TACCs and commercial landings for the Chatham Island and North Island shortfin and longfin eel stocks are shown in Tables 5 and 6. The Chatham Island and North Island fisheries were first introduced into the QMS in 2003–04 and 2004–05, respectively. Note that from 1 October 2007 the TACCs were markedly reduced for all North Island shortfin and longfin stocks.

Table 6: TACCs and commercial landings (t) for Chatham Island (LFE 17) and North Island longfin stocks from 2003–04 to 2016–17 (based on ECLR data).

Fishing Year	LFE 17		LFE 20		LFE 21		LFE 22		LFE 23		Total landings
	TACC	Landings	TACC	Landings	TACC	Landings	TACC	Landings	TACC	Landings	
2003–04	1	< 1	-	-	-	-	-	-	-	-	0.2
2004–05	1	< 1	47	27.4	64	53.5	41	23.9	41	24.5	129.3
2005–06	1	< 1	47	23.7	64	41.2	41	31.6	41	24.2	120.8
2006–07	1	0	47	27.2	64	29.8	41	25.9	41	14.5	97.4
2007–08	1	0	19	17.5	32	31.0	21	17.7	9	6.5	72.8
2008–09	1	0	19	11.5	32	22.7	21	7.7	9	2.5	44.4
2009–10	1	< 1	19	9.6	32	21.6	21	10.6	9	5.8	47.6
2010–11	1	< 1	19	10.2	32	13.7	21	5.7	9	6.2	35.8
2011–12	1	< 1	19	19.9	32	32.0	21	18.6	9	6.7	77.3
2012–13	1	< 1	19	18.3	32	25.1	21	15.1	9	5.6	64.1
2013–14	1	0	19	14.7	32	25.9	21	14.7	9	4.4	59.7
2014–15	1	0	19	10.1	32	9.9	21	12.0	9	3.3	35.3
2015–16	1	< 1	19	6.5	32	9.4	21	4.1	9	1.5	21.5
2016–17	1	0	19	8.0	32	13.9	21	7.4	9	3.9	33.2

1.2 Recreational fisheries

In October 1994, a recreational individual daily bag limit of six eels was introduced throughout New Zealand. There is no quantitative information on the recreational harvest of freshwater eels. The recreational fishery for eels includes any eels taken by people fishing under the amateur fishing regulations and includes any harvest by Maori not taken under customary provisions. The extent of the recreational fishery is not known although the harvest by Maori might be significant.

1.3 Customary non-commercial fisheries

Eels are an important food source for use in customary Maori practices. Maori developed effective methods of harvesting, and hold a good understanding of the habits and life history of eels. Fishing methods included ahuriri (eel weirs), hinaki (eel pots) and other methods of capture. Maori exercised conservation and management methods, which included seeding areas with juvenile eels and imposing restrictions on harvest times and methods. The customary fishery declined after the 1900s but in many areas Maori retain strong traditional ties to eels and their harvest.

In the South Island, Lake Forsyth (Waiwera) and its tributaries have been set aside exclusively for Ngai Tahu. Other areas, such as the lower Pelorus River, Taumutu (Te Waihora), Wainono Lagoon and its catchment, the Waihao catchment, the Rangitata Lagoon and the Ahuriri Arm of Lake Benmore, have been set aside as non-commercial areas for customary fisheries. Mātaitai Reserves covering freshwater have been established in the South Island on the Maitara River, Okarito Lagoon, Waihao River (including Wainono Lagoon and parts of Waituna Stream and Hook River), Lake Forsyth and the Waikawa River. Commercial fishing is generally prohibited in mātaitai reserves. In the North Island, commercial fishing has been prohibited from the Taharoa lakes, Whakaki Lagoon, Lake Poukawa and the Pencarrow lakes (Kohangapiripiri and Kohangatera) and associated catchments.

Table 7: TACs, and customary non-commercial and recreational allowances (t) for South Island eel stocks. Note that an allowance for other sources of fishing-related mortality has not been set.

	LFE 11 Nelson/ Marlborough	LFE 12 North Canterbury	LFE 13 Te Waihora Lake Ellesmere	LFE 14 South Canterbury	LFE 15 Otago/ Southland	LFE 16 West Coast
2016 TAC	3	3	3	3	66.54	32.41
TACC	1	1	1	1	52	25
Customary Non-Commercial Allowance	1	1	1	1	13.27	6.41
Recreational Allowance	1	1	1	1	1.27	1
	SFE 11	SFE 12	SFE 13	SFE 14	SFE 15	SFE 16
2016 TAC	24.87	26.1	171.94	13.57	37.42	38.69
TACC	19	20	134.12	10	29	30
Customary Non-Commercial Allowance	4.87	5.1	34.38	2.57	7.42	7.69
Recreational Allowance	1	1	3.44	1	1	1

Table 8: TACs, and customary non-commercial, recreational, and other fishing-related mortality allowances (t) for the Chatham Island and North Island shortfin stocks.

	SFE 17	SFE 20	SFE 21	SFE 22	SFE 23
TAC	15	148	181	121	36
Customary Non-Commercial Allowance	3	30	24	14	6
Recreational Allowance	1	28	19	11	5
Other fishing-related mortality	1	4	4	2	2

Customary non-commercial fishers desire eels of a greater size, i.e. over 750 mm and 1 kg. Currently, there appears to be a substantially lower number of larger eels in the main stems of some major river catchments throughout New Zealand, which may limit customary fishing. Consequently the access to eels for customary non-commercial purposes has declined over recent decades in many areas. There is no overall assessment of the extent of the current or past customary non-commercial take. For the introduction of the South Island eel fishery into the QMS, an allowance was made for customary non-commercial harvest. It was set at 20% of the TAC for each QMA, equating to 107 t (Table 7). For the introduction of the North Island fishery into the QMS, the customary non-commercial allowance was set at 74 t for shortfins and 46 t for longfins (Tables 8 and 9). For the Chatham Islands, the customary non-commercial allowance was 3 t for shortfin and 1 t for longfin eels (Tables 8 and 9).

Eels may be harvested for customary non-commercial purposes under an authorisation issued under fisheries regulations. Such authorisations are used where harvesting is undertaken beyond the recreational rules. The majority of the South Island customary harvest comes from QMAs ANG 12 (North Canterbury) and ANG 13 (Te Waihora/Lake Ellesmere). Customary regulations were only extended to freshwaters of the Chatham and North Islands in November 2008.

Table 9: TACs, and customary non-commercial, recreational, and other mortality allowances (t) for the Chatham Island and North Island longfin eel fisheries.

	LFE 17	LFE 20	LFE 21	LFE 22	LFE 23
TAC	3	39	60	34	34
Customary Non-Commercial Allowance	1	10	16	6	14
Recreational Allowance	1	8	10	5	9
Other fishing-related mortality	0	2	2	2	2

1.4 Illegal catch

No reliable estimates of illegal catch are available. There is some evidence of fishers exceeding the amateur bag limit, and some historical incidences of commercial fishers operating outside of the reporting regime, but overall the extent of any current illegal take is not considered to be significant.

1.5 Other sources of mortality

Although there is no information on the level of fishing-related mortality associated with the eel fishery (i.e., how many eels die while in the nets), it is not considered to be significant given that the fishing methods used are passive and catch eels in a live state.

Eels are subject to significant sources of mortality from non-fishing activities, although this has not been quantified. Direct mortality occurs through the mechanical clearance of drainage channels, and damage by hydro-electric turbines and flood control pumping (Beentjes et al 2005). Survival of eels through hydroelectric turbines is affected by eel length, turbine type and turbine rotation speed. The mortality of larger eels (specifically longfin females), is estimated to be 100%. Given the large number of eels in hydro lakes, this source of mortality could be significant and reduce spawner escapement from New Zealand. Mitigation activities such as trap and transfer of downstream migrants, installation of downstream bypasses and spillway opening during runs, is expected to have reduced this impact at those sites where such measures have been implemented. In addition to these direct sources of mortality, eel populations are likely to have been significantly reduced since European settlement from the 1840s by wetland drainage (wetland areas have been reduced by up to 90% in some areas), and on-going habitat modification brought about by irrigation, channelisation of rivers and streams and the reduction in littoral habitat. On-going drain maintenance activities by mechanical means to remove weeds may cause direct mortality to eels through physical damage or by stranding and subsequent desiccation.

2. BIOLOGY

Species and general life history

There are 16 species of freshwater eel worldwide, with the majority of species occurring in the Indo-Pacific region. New Zealand freshwater eels are regarded as temperate species, similar to the Northern Hemisphere temperate species, the European eel *A. anguilla*, the North American eel *A. rostrata*, and the Japanese eel *A. japonica*. Freshwater eels have a life history unique among fishes that inhabit New Zealand waters. All *Anguilla* species are facultative catadromous, living predominantly in freshwater and undertaking a spawning migration to an oceanic spawning ground. They spawn once and then die (i.e., are semelparous). The major part of the life cycle is spent in freshwater or estuarine/coastal habitat. Spawning of New Zealand species is presumed to take place in the southwest Pacific. Progeny undertake a long oceanic migration to freshwater where they grow to maturity before migrating to the oceanic spawning grounds. The average larval life is 6 months for shortfins and 8 months for longfins.

The longfin eel is endemic to New Zealand and is thought to spawn east of Tonga. The shortfin eel is also found in South Australia, Tasmania, and New Caledonia; spawning is thought to occur northeast of Samoa. Larvae (leptocephali) are transported to New Zealand largely passively on oceanic surface currents, and the metamorphosed juveniles (glass eels) enter freshwater from August to November. The subsequent upstream migration of elvers (pigmented juvenile eels) in summer distributes eels throughout the freshwater habitat. The two species occur in abundance throughout New Zealand and have overlapping habitat preferences with shortfins predominating in lowland lakes and slow moving soft bottom rivers and streams, while longfins prefer fast flowing stony rivers and are dominant in high country lakes.

Growth

Age and growth of New Zealand freshwater eels was reviewed by Horn (1996). Growth in freshwater is highly variable and dependent on food availability, water temperature and eel density. Eels, particularly longfins, are generally long lived. Maximum recorded age is 60 years for shortfins and 106 years for longfins. Ageing has been validated (e.g. Chisnall & Kalish, 1993). Growth rates determined from the commercial catch sampling programme (1995–97) indicate that in both the North and South Islands, growth rates are highly variable within and between catchments. Shortfins often grow considerably faster than longfins from the same location, although in the North Island longfins grow faster than shortfins in some areas (e.g. parts of the Waikato catchment). South Island shortfins take, on average, 12.8 years (range 8.1–24.4 years) to reach 220 grams (minimum legal size), compared with 17.5 years (range 12.2–28.7 years) for longfins, while in the North Island the equivalent times are 5.8 years (3–14.1 years) and 8.7 years (range 4.6–14.9 years) respectively. Australasian longfin growth is generally greater than that of New Zealand longfins, and closer to that of shortfins.

Growth rates (in length) are usually linear. Sexing immature eels is difficult, but from length at age data for migratory eels, there appears to be little difference in growth rate between the sexes. Sex determination in eels appears to be influenced by environmental factors and by eel density, with female eels being more dominant at lower densities. Age at migration may vary considerably between areas depending on growth rate. Males of both species mature and migrate at a smaller size than females. Migration appears to be dependent on attaining a certain length/weight combination and condition. The range in recorded age and length at migration for shortfin males is 5–22 years and 40–48 cm, and for females 9–41 years and 64–80 cm. For longfinned eels the range in recorded age and length at migration is 11–34 years and 48–74 cm for males, and 27–61 years and 75–158 cm for females. However because of the variable growth rates, eels of both sexes and species may migrate at younger or older ages.

Recruitment

The most sensitive measure of recruitment is monitoring of glass eels, the stage of arrival from the sea. In the Northern Hemisphere where glass eel fisheries exist, catch records provide a long term time series that is used to monitor eel recruitment. In the absence of such fisheries in New Zealand, MPI took the unique opportunity that exists to monitor the relative abundance of elvers arriving at large in-stream barriers, where established elver trap and transfer programmes operate. Provided that the data are collected in a consistent manner every year, these data can be used to provide an index of eel recruitment into New Zealand's freshwaters.

Although New Zealand has a small dataset of elver catch data compared to Asian, European and North American recruitment records, including the 2014–15 season, there are now up to 20 years of reliable and accurate elver catch information for some sites (Martin et al 2016). These records show that the magnitude of the elver catch varies markedly between sites and that there are large variations in catches between seasons at all the sites (Table 10a). Whilst the majority of this variability is likely to be caused by natural oceanic and climatic influences, some is due to changes in fishing effort, technological advances and recording procedures. Consequently, a number of existing records need to be excluded from recruitment trend analyses.

Because of the variability between sites and years, elver catch records were normalised following the method of Durif et al (2008), and a “normal” catch index was calculated for each species, season, and location. The normalised catch index (X_{ij}) is calculated as follows:

$$X_{i,j} = (x_{i,j} - \mu_j) / \sigma_j$$

Where:

$x_{i,j}$ = elver catch for a season

μ_j = mean elver catch at a site for all seasons

σ_j = standard deviation of elver catch at a site for all seasons.

Although several of the sites show that catches peaked during the 2007–08 and 2008–09 migration seasons this is not consistent across all sites and also varies slightly between shortfins and longfins. A trend of increasing catches at Piripaua, however, stand out at present (Figure 2a).

Variation in the distance of dam sites from the sea and possibly differences in migration rates and growth rate between rivers has resulted in some variability in the size (age) structure of elvers captured at the monitored sites. Consequently the median ages of elvers at key sites were determined from examination of otoliths extracted from elvers captured during the 2013–14 season (Table 10b). The median ages were then used to standardise the normalised catch index so that it reflected the relative recruitment of glass eels (0 yrs old) into each catchment.

The standardised recruitment indices indicate that there was a recruitment peak for both shortfins and longfins in the Waikato, Mokau, Patea and Grey rivers around 2006–2007 (Figure 2b). A recruitment peak also occurred at the same time on the Rangitaiki River which, unlike the other four rivers, is on the East Coast.

The Waikato and Northern Wairoa rivers and possibly the Patea River on the West Coast and the Rangitaiki and Wairoa rivers on the East Coast of the North Island all show an increased recruitment of shortfins around 2011 and 2012. In the South Island the Grey River on the West Coast and the Waitaki River on the East Coast also showed increased recruitment of shortfins in 2012 (Figure 2b). Because of the time it takes for longfins to reach these two South Island dams it is still too early to know if longfin recruitment also increased in 2011 and 2012.

The Wairoa and Waiau rivers do not follow the general patterns shown by other sites. Issues with inconsistent fishing effort in the past most likely have disguised the actual recruitment trend for the Waiau River (Figure 2b).

Since the early 1990s there have been four peaks of the average recruitment index for shortfins (1996, 2001, 2006 and 2013) and longfins (1996, 2000, 2006 and 2012) (Figure 2b). The length of time between these peaks varies from four to seven years, indicating a short-term cycle that appears to be influencing recruitment of both species.

Eel larvae are thought to not only actively swim but also use sea currents to reach the New Zealand continental shelf. Examination of regional differences in glass eel mean size and condition indicated an arrival pattern from the north in an anti-clockwise dispersal pattern around New Zealand (Chisnall et al 2002).

There is evidence from duration of runs and catch-effort data that glass eel runs may now be smaller in the Waikato River than in the 1970s (Jellyman et al 2009). However, studies on the variability and temporal abundance of glass eels over a seven year period from 1995 to 2002 at five sites showed no decline in recruitment for either species (Jellyman & Sykes 2004). At these same sites the density of shortfin glass eels exceeded that of longfins for any one year but the annual trends for both species were generally similar (Jellyman et al 2002).

There is some evidence of annual variation influenced by the El Nino Southern Oscillation (ENSO), with the arrival route of glass eels from the northwest being stronger during the La Nina phase and stronger from the northeast during the El Nino phase (Chisnall et al 2002). This may also explain the recruitment pattern seen in the elver trap and transfer programmes (Martin et al 2014). A greater understanding of sea currents, notably along the coastline, and their effects on recruitment patterns, together with longer catch records, particularly from the east coast (e.g., Waitaki and Roxburgh dams), may further elucidate recruitment trends and drivers.

Spawning

As eels are harvested before spawning, the escapement of sufficient numbers of eels to maintain a spawning population is essential to maintain recruitment. For shortfin eels the wider geographic distribution for this species (Australia, New Zealand, southwest Pacific) means that spawning escapement occurs from a range of locations throughout its range. In contrast, the more limited distribution of longfin eels (New Zealand and offshore islands) means that the spawning escapement must occur from New Zealand freshwaters and offshore islands.

FRESHWATER EELS (SFE, LFE, ANG)




Figure 2a: Normal catch index for longfin (LFE) and shortfin (SFE) elvers at monitored sites from 1995-96 to 2014-15. (Notes: incomplete records for season have been omitted; 0 = mean index for entire monitoring period for each site; few shortfins recorded at Mararoa Weir). Mararoa has inconsistent fishing effort so the trend shown may reflect increased trapping efficiency rather than increased recruitment.

FRESHWATER EELS (SFE, LFE, ANG)




Figure 2b: Normal recruitment indices for longfin (LFE) and shortfin (SFE) elvers at the main monitored sites from 1995–96 to 2014–15 (0 = mean catch for entire monitoring period for each site). Mararoa has inconsistent fishing effort so the trend shown may reflect increased trapping efficiency rather than increased recruitment.

Table 10a: Estimated numbers (1000s) of all elvers and, in brackets, longfins only; trapped at key elver trap and transfer monitoring sites by season (Dec–April) 1992–93 to 2013–14. Shaded cells indicate seasons when the records are considered unsuitable for trend analysis (monitoring disruption, flood damage etc.). N/A = no species composition. (From Martin et al 2016 and NIWA unpublished records.).

Year	Wairua	Karapiro	Matahina	Wairere	Patea	Piripaua	Arnold	Waitaki	Roxburgh	Mararoa
1992–93		92 (31)	> 32 (>2)							
1993–94		518 (176)	> 215 (NA)							
1994–95		282 (96)	> 39 (NA)							
1995–96		1 155 (333)	> 144 (NA)							
1996–97		1 220 (246)	14 (4)			2.1 (1)			0.3	
1997–98		2 040 (510)	615 (136)			7.3 (NA)			11	
1998–99		1 097 (341)	1 002 (NA)			3.1 (0.4)			7.4	43 (43)
1999–00		892 (94)	2 001 (NA)	166 (NA)	461 (NA)	2.6 (<0.1)				90 (90)
2000–01		782 (155)	2 054 (NA)	191 (NA)	495 (NA)	6 (0.2)				28 (28)
2001–02		1 596 (246)	619 (27)	130 (NA)	754 (48)	4.1 (0.4)			1	NA
2002–03		1 942 (176)	1 484 (124)	289 (22)	380 (8)	10.2 (0.2)		<0.1 (<0.1)	0.1	36 (36)
2003–04		2 131 (200)	945 (64)	330 (NA)	391 (1)	4.9 (0.2)		4.6 (4.6)	1.4	98 (98)
2004–05		1 333 (132)	1 117 (15)	155 (13)	450 (NA)	8.1 (0.5)	27 (7)	1.5 (1.5)		64 (64)
2005–06		2 178 (483)	1 193 (228)	163 (28)	562 (87)	2.8 (0.1)	14 (8)	4.7 (4.7)		46 (46)
2006–07		1 296 (179)	485 (159)	294 (25)	896 (53)	4.2 (0.3)	107 (52)	3.3 (3.3)		118 (118)
2007–08		2 728 (701)	3 378 (928)	204 (57)	857 (98)	5.7 (1.1)	186 (78)	4.1 (4.1)		133 (133)
2008–09		2 288 (298)	4 307 (517)	216 (16)	480 (82)	9.5 (2.2)	183 (87)	4.7 (3.5)		81 (81)
2009–10		1 708 (232)	1 002 (78)	146 (7)	309 (20)	10.3 (2.9)	20 (5)	2.4 (2.1)		71 (71)
2010–11		1 434 (175)	1 841 (84)	227 (NA)	247 (20)	11.8 (2.5)	114 (49)	2.9 (2.4)		198 (198)
2011–12	3 178 (11)	1 003 (36)	641 (15)	119 (0.5)	72 (6.8)	15.6 (3.1)	76 (26)	7 (5.8)	NA (NA)	266 (266)
2012–13	5 488 (98)	1 771 (139)	2 421 (317)	182 (NA)	74 (16)	33 (5.2)	90 (36)	8.9 (7.1)	14 (14)	128 (128)
2013–14	2 780 (16.2)	1 843 (160)	2 068 (220)	193.1 (NA)	193.2 (23.5)	68.7 (7.9)	65.3 (29.4)	0.2 (0.1)	0.8 (0.8)	150.4 (150.4)
2014–15	3 010 (118)	1 604 (160)	4 736 (275)	241.9 (NA)	260.6 (23.1)	61.2 (4.7)	152.5 (65)	6.0 (4.6)	1.3 (1.3)	135.6 (135.5)

Table 10b: Summary of elver weights, lengths and estimated ages at sites where individual weights and lengths of 100 SFE and 100 LFE (if available) were measured monthly during 2013–14 (from Martin et al 2016).

Location	Species	n	Length (mm)			Weight (g)			Estimated age ^a
			Mean	Median	Range	Mean	Median	Range	
Wairua Falls	LFE	7	60	59	66–55	0.24	0.22	0.35–0.17	– ^b
	SFE	1 318	63	61	130–48	0.26	0.22	1.67–0.07	0
Karapiro	LFE	140	106	104	157–75	1.60	1.3	5.2–0.5	1
	SFE	295	93	91	153–74	0.9	0.8	3.9–0.4	1
Matahina	LFE	272	111	110	152–86	1.53	1.4	4.0–0.6	1
	SFE	750	97	96	133–75	0.96	0.9	2.9–0.4	1
Piripaua	LFE	166	115	112	188–90	1.7	1.5	8.7–0.8	1
	SFE	497	101	100	142–85	1.1	1.1	3.4–0.5	1
Patea	LFE	124	80	79	124–59	0.62	0.56	2.57–0.18	0
	SFE	1 247	74	73	121–57	0.46	0.43	1.95–0.16	0
Arnold	LFE	400	130	126	202–101	2.1	1.8	8.9–0.7	2
	SFE	418	111	108	175–90	1.1	1.0	4.3–0.5	1
Waitaki	LFE	53	196	200	260–118	10.0	8.65	22.1–1.7	4
	SFE	103	132	130	203–102	2.25	1.98	11.3–0.9	2
Roxburgh	LFE	16	159	163	210–120	4.38	4.34	7.5–2.3	– ^b
Mararoa Weir	LFE	1 591	152	137	240–92	4.9	3.0	18.92–0.7	2
	SFE	15	108	104	150–92	1.34	0.99	3.8–0.6	– ^b

^a Fresh water age based on median lengths of elver at each site and nation-wide age vs length regression.

^b Insufficient number of elvers measured to accurately determine age distribution.

3. STOCKS AND AREAS

The lifecycle of each species has not been completely resolved but evidence supports the proposition of a single (panmictic) stock for each species. Biochemical evidence suggests that shortfins found in both New Zealand and Australia form a single biological stock. Longfins are endemic to New Zealand and are assumed to be a single biological stock.

Within a catchment, post-elver eels generally undergo limited movement until their seaward spawning migration. Therefore once glass eels have entered a catchment, each catchment effectively contains a separate population of each eel species. The quota management areas mostly reflect a combination of these catchment areas.

Shortfin and longfin eels have different biological characteristics in terms of diet, growth, maximum size, age of maturity, reproductive capacity, and behavioural ecology. These differences affect the productivity of each species, and the level of yield that may be sustainable on a longer term basis, as well as their interactions with other species. In order that catch levels for each species are sustainable in the longer term, and the level of removals does not adversely affect the productivity of each species, it is appropriate that the level of removals of each species is effectively managed.

4. STOCK ASSESSMENT

There is no formal stock assessment available for freshwater eels. Fu et al (2012) developed a length-structured longfin population model that generated New Zealand-wide estimates of the pre-exploitation female spawning stock biomass (approximately 1 700 t) as well as the pre-exploitation biomass of legal-sized eels (16 000 t in all fished areas and 6 000 t in protected areas). By contrast, the model estimated current female spawning stock biomass to be approximately 55% of pre-exploitation levels, whereas the current biomass of legal-sized eels ranged from 20% to 90% of the pre-exploitation level for the fished areas. However, the Working Group did not accept the assessment and noted that further analyses were necessary to investigate the models underlying assumptions; given that the results were strongly driven by estimates of longfin commercial catches from individual eel statistical areas as well as GIS-based estimates of recruitment.

4.1 Size/age composition of commercial catch

Catch sampling programmes sampled commercial eel landings throughout New Zealand over three consecutive years between 1995–96 and 1997–98, and then in 1999–2000 and 2003–04 (Beentjes 2005, Speed et al 2001). Sampling provided information on the length and age structure, and sex composition of the commercially caught eel populations throughout the country, and indicated a high degree of variability within and among catchments.

Monitoring commercial eel fisheries programme

The commercial eel monitoring programme collects processor recorded catch data for each species by size-grade (market determined; two to three grades) and catch location (eel statistical sub-area; catchment based), from virtually all commercial landings throughout New Zealand. This programme began in 2003–04 in the North Island and 2010–11 in the South Island (Beentjes 2013, 2016) with twelve years of North Island data and five years of South Island data collected by the end of 2014–15. This programme is ongoing with collection of data from 2015–16 to 2017–18 in progress.

North Island – North Island commercial eel catch is highly aggregated with nearly one-third of the shortfin catch caught from just 3 of the 65 subareas (AA4, Dargaville; AD12, Lake Waikare and Port Waikato; and AC1, Hauraki plains west). Similarly, one third of North Island longfin was caught from just four subareas (AA4, Dargaville; AD10, Waipa River; AD12, Lake Waikare, Port Waikato; and AL1, Lake Wairarapa). North Island shortfin annual catch over 12 years showed no consistent trend in annual catch weight or in the distribution of these catches in the three size grades. The longfin fishery is more prone to market demand fluctuations than shortfin because it is a less desirable species of eel. North Island longfin annual catch over the 12 years has fluctuated with an overall trend of declining catch. Factors that may have influenced annual longfin catches, overall and within size grades, include port price, the 58% TACC reductions for North Island longfin stocks implemented in the 2007–08 fishing year, market fluctuations, and limited quota being offered as ACE in some years. The number of subareas for which shortfin and longfin catch was landed has been declining indicating a contraction in the spatial distribution of fishing effort over time. Despite this the catch of both species in the key subareas over the 12 years shows no apparent trends.

South Island – South Island commercial eel catch is highly aggregated especially shortfin where nearly three-quarters of the catch originates from just two of the 58 subareas (Te Waihora, AS1 and AS2; and Lake Brunner, AX4). Longfin in the South Island is less aggregated than shortfin, but half of the catch originated from just seven subareas (AW11, Maitai River coast; AW9, Oreti River coast; AW3, Oreti River inland down to Bog Burn; AV10, Clutha River coast; AP2, Wairau River; AU5, Waitaki River; and AX2, Buller River). There are no trends in catch by size grade for either species over the five year time series. Catch of longfin has been stable in the key subareas, but more variable for the subareas with smaller catches. Shortfin catches by subareas were generally similar each year, except that AS1 (lake) and AS2 (migration area) catches tend to display opposite trends because the Te Waihora quota (ANG 13) can be filled from either the lake or the migration area.

4.2 Catch-per-unit-effort analyses

Each species of eel is considered to be a New Zealand wide stock, with common species-specific spawning grounds within the Fiji Basin. However, once recruited to a river system, eels do not move between catchments, so eels within each catchment may be regarded as separate sub-populations for management purposes. Maintaining sub-populations within each QMA at or above (sub-area proxies for) B_{MSY} , will ensure that the entire (national) stock of each species is maintained at that level. To develop sub-area proxies, standardised catch-per-unit-effort (CPUE) analyses have been conducted for the commercial shortfin and longfin eel fisheries by Eel Statistical Area (ESA; Table 11 and Figure 3) from 1990–91 to 2011–12 for all North Island ESAs and from 1990–91 to 2012–13 for all South Island ESAs (Tables 12 to 13 and Figures 4–7). These CPUE series monitor the relative abundance of each eel species within the area fished commercially within each ESA.

North Island CPUE

The North Island CPUE analyses undertaken in 2016 and 2017, using data up to 2014–15 included, for the first time, a binomial analyses on the valid zero catches, as well as the routine GLM analyses of positive catch. In addition, reconstructed target species was included as an explanatory variable, as were water quality variables. The variable ‘catcher_ID’ was not included because it has only been recorded since 2001–02 on the new ECE returns (Beentjes & McKenzie 2017); however, the data were linked by permit holder and client name (see below). Target species was recorded in CELR forms, but not in ECER forms. Target species was reconstructed for all records from recorded CELR target species and species proportions using a simple optimisation to evaluate the best proportion to use (Cohen’s kappa coefficient). Target species was reconstructed for all records, including those from CELR data. In some cases, target species was defined on the basis of a minimum catch composition of 80%. Higher values tended to assign too many records to the category ‘either’, when kappa was above 80%. Target species often explained the most variance in the positive catch GLM, especially for longfin for which the trends in CPUE changed more than shortfin compared to previous analyses when target was not offered to the model. Target species could not be offered to the binomial model because, by definition, a target of longfin or shortfin cannot result in zero catch in the models and consequently the May 2017 plenary rejected the binomial model.

Prior to the introduction of North Island eel stocks into the QMS in 2004–05, some fishers had fished for existing permit holders during the permit moratorium and following introduction of eels into the QMS began fishing under their own permit numbers (Beentjes & Dunn 2010). If these fishers had fished for someone else pre-QMS and if they were the only fisher that had landed catch under a pre-QMS *Client_name*, and that client did not land catch pre- and post-QMS, they were linked in the analyses. There were 16 linkages made.

The transition between CELR and ECER in 2001–02 is unlikely to have biased trends in relative abundance (CPUE) as the way in which catches of longfin and shortfin eels were estimated and effort data was recorded remained unchanged across form types, with both forms providing estimated catch of shortfin and longfin eels, the number of nets set per night, and the statistical area where eels were caught.

In general CPUE for North Island shortfin, with the exception of Northland (ESA AA) where CPUE steadily increased throughout the time series, either initially declined or there were no trends, followed by strong increases, beginning from around 2002 (Table 12, Figure 4) (Beentjes & McKenzie 2017).

For longfin there were generally fewer data than for shortfin for most areas and indices were often more variable, associated with wider confidence intervals, or could not be estimated for all years (Table 13, Figure 5). The addition of reconstructed target species as an explanatory variable had a much greater impact on longfin indices than shortfin indices (Beentjes & McKenzie 2017). The apparent trends for longfin have therefore changed considerably since the last analyses (which used data up to 2011–12; Beentjes & Dunn 2013b). For ESAs with the largest data sets, trends were as follows: Northland (AA) - very slight downward trend over the time series; Auckland (AB) - a slight decline to 2005, but stable thereafter; Hauraki (AC) - steep decline to 2000–01, and then without trend/stable to 2014–15; Waikato (AD) - moderate decline to 1998, and then a gradual increase to around the level of the former peak by 2014–15 (Table 13, Figure 5). For the other ESAs, which were data poor, CPUE increased after an initial decline (AE, AG, AH, AJ, AK, AL), but gradually declined since about 2012 for AJ, AK, and AL.

Several factors may have resulted in conservative estimates of North Island longfin eel CPUE, especially after 2005–06:

1. The unrecorded return of small and medium sized longfin eels to the water. This became more prevalent after the substantial reduction in NI longfin quotas in 2007–08, as many fishers do not have ACE to cover all of their catch (larger longfins are more valuable than small and medium specimens). Industry were previously unaware that eels of legal size (220 g–4 kg) that are released are supposed to be recorded on ECL returns under the destination X code which was only available as a legitimate code on ECL forms since 2007–08. Further, at the Eel Working Group Meeting in April 2017 it was established that

some fishers are incorrectly recording only their retained legal sized eels on the ECE returns and thus the estimated catch used in CPUE analyses will be biased downward as will the CPUE in recent years. North Island destination X catch was only 3% of the landed catch in 2014–15. The way in which individual fishers report discarded legal catch needs further investigation.

2. The introduction of a maximum size of 4 kg in 2007–08. Longfins over 4 kg could be legally landed before this date. There is currently no legal requirement to record the catch of eels over 4 kg.
3. Avoidance of longfin habitat post 2006–07 in some statistical areas as there is currently insufficient available ACE to allow targeting of longfin eels. The QMA most affected is LFE 23 (current TACC is 9 t) where, since 2007–08 up to half the ACE has not been made available for lease. Of the available longfin ACE, almost all is leased to a fisher operating in the Taranaki statistical area (AJ) of this QMA, leaving very little for the Wanganui-Rangitikei statistical area. The fisher in the latter statistical area consequently targets shortfin eels in farm dams, dune lakes and the lower reaches of some rivers; thereby avoiding high longfin eel catch rates in the Rangitikei River.
4. Voluntary uptake of larger escape tubes (31 mm) from 2010–11 (regulated in 2012–13) may have resulted in a stepped drop in CPUE.

Table 11: New Zealand Eel Statistical Areas (ESAs). Areas were given a numeric designation prior to Oct. 2001, at which point letter codes were assigned.

ESA	Letter code	Numeric code
Northland	AA	1
Auckland	AB	2
Hauraki	AC	3
Waikato	AD	4
Bay of Plenty	AE	5
Poverty Bay	AF	6
Hawke Bay	AG	7
Rangitikei-Wanganui	AH	8
Taranaki	AJ	9
Manawatu	AK	10
Wairarapa	AL	11
Wellington	AM	12
Nelson	AN	13
Marlborough	AP	14
South Marlborough	AQ	14
Westland	AX	15
North Canterbury	AR	16
South Canterbury	AT	17
Waitaki	AU	18
Otago	AV	19
Southland	AW	20
Te Waihora (outside-migration area)	AS1	21
Te Waihora migration area	AS2	21
Chatham Islands	AZ	22
Stewart Island	AY	23

South Island CPUE

The Eel Working Group (EELWG-2012-05) made the decision to split South Island CPUE analyses into pre- and post-QMS time series with post-QMS CPUE analyses only required for areas with sufficient data and fishers (ESAs: Westland AX, Otago AV, Southland AW). This was done because many fishers fishing under existing permits pre QMS obtained their own quota and entered the fishery as “new” entrants when the QMS was introduced. Fishing coefficients for existing permit holders were therefore likely to have changed considerably after the QMS was introduced. It is not possible to separate catches in the pre-QMS data into individual fisher catch and effort, as was done in the North Island analysis, as the CELR forms used up to 2001–02 included only a field for permit holder, with no way of identifying individual operators. This problem was solved in 2001–02 with the introduction of the new ECER form by adding a field which identified the fisher (i.e., “catcher”) filling out the form.




Figure 3: New Zealand Eel Statistical Areas (ESAs).

Table 12: South Island CPUE indices for shortfin eels by Eel Statistical Area (ESA). Separate indices are presented for pre-QMS (1991–2000) and post-QMS (2001–2010). Fishing years are referred to by the second year (e.g., 1990–91 is referred to as 1991). – insufficient data. (See Table 11 for ESA area names). (Data from Beentjes & Dunn 2015).

QMS status	Year	Shortfin (South Island ESAs)								
		AN	AP_AQ	AR	AT	AU	AV	AW	AX	AS1
Pre-QMS	1991	–	2.36	1.13	2.09	1.7	1.51	1.3	0.96	–
	1992	–	1.94	1.09	1.07	1.46	1.2	1.03	0.61	–
	1993	1.24	1.59	0.94	0.84	0.69	1.05	0.99	1.07	–
	1994	–	1.34	1.01	1.01	1.06	1.03	1.33	0.95	–
	1995	1.16	1.14	0.81	0.79	0.84	0.92	1.01	0.9	–
	1996	0.89	0.65	0.98	0.97	1.31	0.87	0.88	0.85	–
	1997	0.41	0.55	0.97	0.85	0.85	0.9	0.79	0.75	–
	1998	0.97	0.38	1	1.07	1.1	0.84	0.89	1.31	–
	1999	1.37	0.73	1.13	0.67	0.61	0.83	0.9	1.52	–
	2000	1.43	0.91	0.99	1.13	0.88	1.02	1.01	1.48	–
	2001	–	–	–	–	–	–	–	–	–
Post-QMS	2002	–	–	–	–	–	0.86	0.68	0.81	0.37
	2003	–	–	–	–	–	0.86	0.61	0.73	0.42
	2004	–	–	–	–	–	0.76	0.91	0.87	0.51
	2005	–	–	–	–	–	1.05	1.03	0.99	0.58
	2006	–	–	–	–	–	0.89	0.83	0.87	0.79
	2007	–	–	–	–	–	1.21	1.07	0.99	1.17
	2008	–	–	–	–	–	0.8	1.29	0.89	1.28
	2009	–	–	–	–	–	1.26	0.8	1.49	1.31
	2010	–	–	–	–	–	1.27	1.23	1.16	1.17
	2011	–	–	–	–	–	1.34	1.35	1.16	2.34
	2012	–	–	–	–	–	1.12	1.26	1.11	2.29
	2013	–	–	–	–	–	0.81	1.34	1.16	2.23

Table 13: South Island CPUE indices for longfin eels by Eel Statistical Area (ESA). Separate indices are presented for pre-QMS (1991–2000) and post QMS (2001–2010). Fishing years are referred to by the second year (e.g., 1990–91 is referred to as 1991). - insufficient data; –, no analysis. (See Table 11 for ESA area names). Data from Beentjes & Dunn (2015).

QMS status	Year	Longfin (South Island ESAs)							
		AN	AP_AQ	AR	AT	AU	AV	AW	AX
Pre-QMS	1991	2.29	1.72	1.29	1.89	1.19	1.35	1.46	1.09
	1992	1.15	1.18	0.87	0.74	0.95	1.2	1.13	0.95
	1993	0.8	1.21	1.00	0.78	0.82	1.14	1.13	0.76
	1994	1.06	1.43	1.06	1.05	0.78	1.27	1.22	0.89
	1995	0.85	1.17	0.75	0.88	0.69	0.93	0.99	1.1
	1996	0.81	1.19	1.21	0.78	1.22	0.8	1	0.99
	1997	0.66	0.68	1.09	0.96	1.11	0.86	0.92	0.94
	1998	0.72	0.77	0.75	0.99	0.97	0.87	0.79	0.97
	1999	1.1	0.83	1.02	0.85	1.34	0.85	0.68	1.11
	2000	1.23	0.47	1.10	1.59	1.14	0.91	0.91	1.29
	2001	–	–	–	–	–	–	–	–
	2002	–	–	–	–	–	0.91	1	0.8
	2003	–	–	–	–	–	0.84	1.09	0.79
	2004	–	–	–	–	–	0.92	0.85	0.93
	2005	–	–	–	–	–	1.11	1.1	0.94
	2006	–	–	–	–	–	0.95	1.05	0.96
	2007	–	–	–	–	–	1.05	0.82	1.01
	2008	–	–	–	–	–	0.98	0.92	0.95
	2009	–	–	–	–	–	1.12	0.92	1.06
	2010	–	–	–	–	–	0.94	0.86	1.28
	2011						1.32	1.23	1.23
	2012						0.96	1.15	1.01
	2013						0.99	1.12	1.16

This problem was less severe in the North Island because NI eels were introduced to the QMS after the new ECER forms had been developed, making it possible to link catcher and permit holders before and after the introduction to the QMS. The most recent South Island CPUE analyses, up to 2012–13, included new predictor variables including: target species, water quality data (e.g., nitrogen, phosphates, clarity, temperature), and catcher (Beentjes & Dunn 2015). Catcher was only available for the post-QMS analyses. The first year in the post-QMS standardised CPUE time series is 2001–02 when catcher was first recorded on the new ECERs.

Westland (AX) – Shortfin pre-QMS CPUE fluctuated without trend from 1990–91 to 1996–97 and then increased sharply to 1999–2000. Post-QMS shortfin CPUE increased steadily from 2001–02 to 2012–13. Longfin pre-QMS CPUE declined from 1990–91 to 1992–93, and then increased steadily to 1999–2000. Post-QMS longfin CPUE increased steadily from 2001–02 to 2012–13 (Tables 12 and 13, Figure 6).

Otago (AV) – Shortfin pre-QMS CPUE declined steadily to 1998–99, then increased sharply to 1999–2000. Post-QMS shortfin CPUE increased steadily from 2001–02 to 2010–11, and then declined. Longfin pre-QMS CPUE declined steadily from 1990–91 to 1995–96 and was stable from then to 1999–2000. Post-QMS longfin CPUE was variable but overall increased slightly from 2001–02 to 2012–13 (Tables 12 and 13, Figure 6).

Southland (AW) – Shortfin pre-QMS CPUE declined slowly from 1990–91 to 1996–97 and then gradually increased to 1999–2000. Post-QMS shortfin CPUE was variable but generally increased steadily from 2001–02 to 2012–13. Longfin pre-QMS CPUE declined steadily from 1990–91 to 1999–2000. Post-QMS longfin CPUE was variable and showed a gradual decline from 2001–02 to 2009–10, and then a substantial increase to 2012–13 (Tables 12 and 13, Figure 6).

Te Waihora

CPUE analyses for Te Waihora were only carried out for AS1 feeder shortfin (the lake, outside the migration area) from 2000–01, coinciding with the introduction of the reporting codes (AS1 and AS2), to 2012–13. The most recent analyses included new predictor variables: lake level, status of lake opening (i.e., open or closed), catcher (Beentjes & Dunn 2015). The standardised CPUE time series begins in 2001–02, when the new ECER form was introduced and catcher was first recorded. CPUE of feeder shortfin eels in Te Waihora increased six fold from 2001–02 to 2010–11 and was reasonably stable from 2010–11 to 2012–13 (Figure 7).




Figure 4: Trends in North Island shortfin CPUE indices for all North Island ESAs from 1990–91 to 2014–15, except Poverty Bay (AF) and Wellington (AM) where there was insufficient data. Vertical dotted line indicates the introduction to the QMS in 2004–05 (from Beentjes & McKenzie 2017).





Figure 5: Trends in North Island longfin CPUE indices for all North Island ESAs from 1990–91 to 2014–15, except Poverty Bay (AF) and Wellington (AM) where there was insufficient data. Vertical dotted line indicates the introduction to the QMS in 2004–05. (From Beentjes & McKenzie 2017).

It is very likely that the fishery has experienced a progressive improvement in yield per recruit as the minimum legal size was incrementally increased from 140 g in 1993–94 to 220 g in 2001–02. Analyses of eel size composition in the lake in the 1990s compared to that in recent years demonstrates that the size of commercially caught eels has substantially increased over time, supporting the concept of an improved yield per recruit (Figure 8; Beentjes & Dunn 2014).


4.3 Biomass estimates

Estimates of current and reference biomass for any eel fish stock are not available. Recent estimates of approximately 12 000 t have been made for longfin eels (Graynoth et al 2008, Graynoth & Booker 2009), but these are based on limited data on density, growth and sex composition of longfin eel populations in various habitat types, including lakes and medium to large rivers.

Otago (AV)



Southland (AW)



Westland (AX)




Figure 6: Trends in South Island shortfin and longfin CPUE indices for key ESAs: Otago (AV), Southland (AW), and Westland (AX). Separate indices are presented for pre-QMS (1991–2000) and post-QMS (2002–2013). (From Beentjes & Dunn 2015).

Te Waihora (AS1)




Figure 7: Te Waihora shortfin CPUE indices for AS1 (outside migration area) from 2001–02 to 2012–13. (From Beentjes & Dunn 2015).




Figure 8: Size grade proportions of shortfin eels harvested from Te Waihora AS1 (lake) from eel processors Levin Eel Trading Ltd in 2009–10 to 2011–12, and Mossburn Enterprises Ltd in 2010–11 and 2011–12. The equivalent size grades have been estimated from the length of eels taken during commercial catch sampling of the commercial catch in 1995–96 to 1997–98 (from Beentjes & Dunn 2014).

4.4 Yield estimates and projections

In the absence of accurate current biomass estimates, this could not be estimated. Biological parameters relevant to the stock assessment are given in Table 14.

Table 14: Estimates of biological parameters.

Fishstock	Estimate	Source
1. Natural mortality (M)		
Unexploited shortfins (Lake Pounui)	$M = 0.038$	Jellyman (unpub. Data)
Unexploited longfins (Lake Pounui)	$M = 0.036$	Jellyman (unpub. Data)
Unexploited longfins (Lake Rotoiti)	$M = 0.02$	Jellyman (1995)
2. Weight (g) of shortfin and longfin eels at 500 mm total length		
	Mean weight	Range
Shortfins Lake Pounui	263	210–305
Shortfins Waihora	250	210–303
Longfins Lake Pounui	307	250–380

4.5 Other factors

Yield-per-recruit

Yield-per-recruit (YPR) models have been run on Te Waihora (Lake Ellesmere) and Lake Pounui data to test the impact of increases in size limit. Results indicated that an increase in minimum size should result in a small gain in YPR for shortfins in Te Waihora and longfins in Lake Pounui, but a decrease for shortfins in Lake Pounui.

A practical demonstration of the benefits of an increase in size limit has been reported from the Waikato area, where a voluntary increase in minimum size from 150 to 220 g in 1987 resulted in decreased CPUE for up to 18 months, but an increase thereafter.

Spawning escapement

A key component to ensuring the sustainability of eels is to maintain spawner escapement. As a sustainability measure, the Mohaka, Motu and much of the Whanganui River catchments were closed to commercial fishing in early 2005 to aid spawning escapement. The importance of adequate spawner escapement for eels is evident from the three northern hemisphere (*A. anguilla*, *A. rostrata* and *A. japonica*) species, which are all extensively fished at all stages of their estuarine/freshwater life stage and are subject to a variety of anthropogenic impacts similar to the situation in New Zealand. There has been a substantial decline in recruitment for all three northern hemisphere species since the mid-1970s with less than 1% of juvenile resources estimated to be remaining for major populations in 2003 (Quebec Declaration of Concern 2003). More recently, Dekker & Casselman (2014) concluded that “the recent recruitment increase of some [northern hemisphere] stocks, and the relative stability of others, indicate that after many decades of continued decline depleted eel stocks around the world have the potential to recover”.

Longfin habitat

It was estimated, based on GIS modelling in the early 2000s (Graynoth et al 2008), that 5% of longfin eel habitat throughout New Zealand is in water closed to fishing where there is protected egress to the sea to ensure spawning escapement. A further 10% of longfin habitat was estimated to be in areas closed to fishing in upstream areas but where the spawning migration could be subject to exploitation in downstream areas (migratory eels are not normally taken by commercial fishers). An additional 17% of longfin habitat was in small streams that are rarely or not commercially fished. Therefore, about 30% of longfin habitat in the North Island and 34% in the South Island was either in a reserve or in rarely/non-fished areas (Graynoth et al 2008). However, the estimate of the proportion of longfin habitat in streams rarely or not commercially fished was based on poor assumptions and was consequently vastly underestimated.

In 2015, commercial longfin eel fishing effort throughout New Zealand was mapped using GIS methods, providing the first detailed and high resolution representation of where and how often fishers set their nets in New Zealand rivers, lakes and harbours. The data used in the study came from face to face interviews with 53 commercial longfin fishers from throughout New Zealand and covered the five year period from 2009–10 to 2013–14. From these data, estimates were made of the proportion of

longfin habitat that is currently fished (Beentjes et al 2016). The total current longfin habitat in rivers was derived from ‘probability of longfin capture’ models. About one quarter (27.2%) of the New Zealand longfin river and lake habitat, currently accessible to longfin eels, was commercially fished (32.5% in the South Island, and 22.5% in the North Island) (Table 15). The proportion of virgin/original longfin habitat affected by anthropogenic activity (impeded access by dams and other structures, habitat degradation, and commercial fishing) is estimated at 42% (= Max. impacted abundance) (Table 15). Forty percent of the current habitat available to longfin eels in New Zealand is estimated to be within DOC Public Conservation Land, and just over half of this is in natural lakes (Beentjes et al 2016). Generally DOC will not issue concessions for commercial eel fishing in Public Conservation Land, except for short fin eels in Lake Brunner.

Table 15: Estimates of total current longfin habitat fished, virgin habitat fished, and maximum impacted abundance from all rivers and lakes by QMA, eel statistical area, and overall for South Island, North Island and New Zealand. Current lake habitat includes that from natural lakes over 0.9 km², and rivers where longfin eels have unimpeded access to, and egress to the sea. Maximum impacted abundance is the proportion of virgin habitat affected by anthropogenic activities including loss to dams, impeded access, commercial fishing, and habitat loss. Max, maximum. QMA, Quota Management Area. (Table from Beentjes et al 2016).

Island	QMA	Eel Area	Statistical	Percent (%)		
				Current habitat fished	Virgin habitat fished	Max. impacted abundance
North Island	LFE 20	AA		36.1	34.7	40.2
North Island	LFE 20	AB		34.9	33.8	38.2
North Island	LFE 21	AC		50.0	47.6	55.0
North Island	LFE 21	AD		43.2	34.4	55.7
North Island	LFE 21	AE		17.4	16.2	23.9
North Island	LFE 21	AF		8.6	8.2	13.6
North Island	LFE 22	AG		17.3	16.0	24.7
North Island	LFE 23	AH		24.8	23.6	29.9
North Island	LFE 23	AJ		17.0	15.9	23.6
North Island	LFE 22	AK		36.0	34.5	40.6
North Island	LFE 22	AL		4.2	4.1	5.0
North Island	LFE 22	AM		2.4	2.2	7.4
South Island	ANG 11	AN		11.5	11.1	15.5
South Island	ANG 11	AP		42.1	40.1	47.1
South Island	ANG 12	AQ		7.9	7.6	12.4
South Island	ANG 12	AR		58.1	55.9	61.7
South Island	ANG 13	AS		0.0	0.0	0.4
South Island	ANG 14	AT		38.6	37.3	42.1
South Island	ANG 14	AU		52.2	12.4	85.9
South Island	ANG 15	AV		46.2	12.5	82.8
South Island	ANG 15	AW		32.2	24.2	40.7
South Island	ANG 16	AX		30.2	29.0	34.0
North Island	All	All		22.5	20.9	29.0
South Island	All	All		32.5	21.8	52.6
New Zealand	All	All		27.2	21.4	42.1

Sex ratio

The shortfin fishery is based on the exploitation of immature female eels, as most shortfin male eels migrate before reaching the minimum size of 220 g. The exception to this is Te Waihora where migratory male shortfin eels are also harvested. The longfin fishery is based on immature male and female eels.

A study on the Aparima River in Southland in 2001–02 found that female longfins were rare in the catchment. Only five of 738 eels sexed were females (McCleave & Jellyman 2004). This is in contrast to a predominance of larger female longfins in southern rivers established by earlier research in the 1940s and 1950s, prior to commercial fishing. The sex ratio in other southern catchments, determined from analysis of commercial landings, also show a predominance of males. In contrast some other catchments (Waitaki River, some northern South Island rivers) showed approximately equal sex ratios. The predominance of males in the size range below the minimum legal size of 220 g cannot be attributed

directly to the effects of fishing. Because the sexual differentiation of eels can be influenced by environmental factors, it is possible that changing environmental factors are responsible for the greater proportion of male eels in these southern rivers (Davey & Jellyman 2005).

Enhancement

The transfer of elvers and juvenile eels has been established as a viable method of enhancing eel populations and increasing productivity in areas where recruitment has been limited. Elver transfer operations are conducted in summer months when elvers reach river obstacles (e.g., the Karapiro Dam on the Waikato River; see Table 10a) on their upriver migration. Nationally some 10 million elvers are now regularly caught and transferred upstream of dams each year.

To mitigate the impact of hydro turbines on migrating eels, a catch and release programme for large longfin females has been conducted from Lake Aniwhenua with release below the Matahina Dam since 1995. An extensive capture and release programme has also been conducted from Lake Manapōuri to below the Mararoa Weir on the Waiau River, Southland by Meridian Energy since 1998. Limited numbers of longfin migrants are also transferred to below the Waitaki Dam by local Runanga. Adult eel bypasses have been installed at the Wairere Falls and Mokauiti power stations in the Mokau River catchment since 2002 and controlled spillway openings have been undertaken at Patea Dam during rain events in autumn (when eels are predicted to migrate downstream) since the late 1990s. Additional eel protection infrastructure are currently being installed at Patea Dam and ongoing studies, including downstream bypass trials are in progress at Karapiro Dam (Waikato), Lake Whakamarino (Waikaremoana Power Scheme) and Wairua (Titoki) Power Station. So far, the effectiveness of none of these varied mitigation activities has been fully assessed.

Several projects have been undertaken to evaluate the enhancement of depleted customary fisheries through the transfer of juvenile eels. In 1997, over 2000 juvenile shortfin eels (100–200 g) were caught from Te Waihora (Lake Ellesmere), tagged and transferred to Cooper's Lagoon a few kilometres away (Jellyman & Beentjes 1998, Beentjes & Jellyman 2002). Only ten tagged eels, all females, were recovered in 2001. It is likely that a large number of eels migrated to sea as males following the transfer. Another project in 1998 transferred 7600 (21% tagged) mostly shortfin eels weighing less than 220 g from Lake Waahi in the Waikato catchment to the Taharoa Lakes near Kawhia (Chisnall 2000). No tagged eels were recovered when the lakes were surveyed in 2001. It is considered that a large number of shortfin eels migrated from the lake as males following the transfer. The conclusion from these two transfers is that transplanted shortfin eels need to be females, requiring that eels larger than 220 g and above the maximum size of migration for shortfin males need to be selected for transfer.

In 1998 approximately 10 000 juvenile longfin eels were caught in the lower Clutha River and transferred to Lake Hawea, of which 2010 (about 20%) were tagged (Beentjes 1998). In 2001, of 216 recaptured eels, 42 (19.4%) had tags (i.e. very little tag loss) (Beentjes & Jellyman 2003). The transferred eels showed accelerated growth and the mean annual growth in length was almost double that of eels from the original transfer site and all recaptures were females. A further sample of Lake Hawea in 2008 showed that of 399 longfin eel recaptures, 79 had tags (19.2%), indicating continued good tag retention (Beentjes & Jellyman 2011). Growth rate from the 2008 tag-recaptures was significantly greater than at release, but less than in 2001 and all recaptures were females.

Trends in the commercial catches from areas upstream of hydro dams on the Waikato, Rangitaiki and Patea rivers indicate that elver trap and transfer operations has improved or at least maintained the eel populations upstream of barriers (Beentjes & Dunn, 2010). Comparison of historical eel survey results have confirmed these observations (e.g. Beentjes et al 1997, Boubée et al 2000, Boubée & Hudson 2009, Crow & Jellyman 2010)

5. FUTURE RESEARCH NEEDS

- The “target species” reconstruction based on CELR data needs to be examined further by, for example, running sensitivities to determine the effect of different assumptions.
- For the Te Waihora shortfin CPUE, explore the possibility of developing an index of the ratio between the AS1 and AS2 catch as a potential explanatory variable.
- Investigate the utility of using more stringent criteria for choosing core permits.
- Examine trends over time for individual fishers; i.e. consider deriving fisher-based indices as an alternative way of standardising.
- Determine whether ancillary data exist that can be used to refine or verify the derived targets.
- Determine the proportion of fishers using destination code X to report the catches of legal-sized fish that are released.
- Identify the fishers who haven’t been using destination X correctly and fix this to the extent possible. Identify whether the issue is specific to certain areas. For some fishers it may be necessary to add the destination code X estimates from the ECLR forms to the catch estimates from the ECER forms to obtain a more accurate estimate of catch per day for the CPUE analyses.
- Investigate ways of compensating for the lack of recording of eels over 4 kg since 2007–08 (especially since this should be rectified once new forms are developed).
- For areas with few fishers or records, the Eel Working group should consider merging statistical areas and analysing at the QMA level. Alternatively the Working Group needs to consider ways of developing statements about stock status for areas with few fisheries or low effort.
- Investigate the possibility of augmenting the current data with information from customary fisheries.
- Calculate a weighted CPUE by QMA, with the weighting based on the amount of suitable habitat in each area.

6. STATUS OF THE STOCKS

There are no Level 1 Full Quantitative Stock Assessments on which to base specific recommendations on eel catch levels. Nevertheless, recruitment data, commercial CPUE indices, information on spawner escapement, and information on the proportion of longfin habitat fished allow for Level 2 Partial Quantitative Stock Assessments of longfin and shortfin eels.

Stock Structure Assumptions

Longfin and shortfin eels are considered to be New Zealand wide stocks, with common species-specific spawning grounds within the Fiji Basin. However, once recruited to a river system, eels do not move between catchments, so eels within each catchment may be regarded as separate sub-populations for management purposes. Maintaining sub-populations within each QMA at or above (sub-area proxies for) B_{MSY} , will ensure that the entire (national) stock of each species is maintained at that level. North Island QMAs have from two to four ESAs, and South Island QMAs all have two, except Westland (LFE 16 and SFE 16) which has one. ESAs also contain multiple catchments or subpopulations from which eels are harvested.


Status of South Island Eels

Level 2 Partial Quantitative Stock Assessments are conducted by statistical area and species, and are only possible where accepted indices of abundance are available; i.e. Westland, Otago, Southland and Te Waihora). Standardised CPUE provides information on the abundance of commercially harvested eels (300 g–4000 g) in areas that are fished commercially. Approximately 67% of currently available longfin habitat on the South Island is either in reserves or in areas rarely or never fished by commercial fishers.


- Westland (AX) longfin

Stock Status	
Year of Most Recent Assessment	2014
Assessment Runs Presented	Standardised CPUE
Reference Points	Target: B_{MSY} assumed, but not estimated Interim Soft Limit: Mean CPUE from 2001–02 to 2002–03 Hard Limit: 50% of Soft Limit Overfishing threshold: F_{MSY} assumed, but not estimated
Status in relation to Target	Unknown
Status in relation to Limits	Soft Limit: Unlikely (< 40%) to be below Hard Limit: Very Unlikely (< 10%) to be below
Status in relation to Overfishing	Unknown

Historical Stock Status Trajectory and Current Status



Comparison of standardised CPUE for longfin eels in Westland (AX) from 1990–91 to 1999–2000 (pre-QMS) and 2001–02 to 2012–13 (post-QMS) (from Beentjes & Dunn 2015). Also shown is the total estimated longfin catch in AX from ECERs. The two CPUE series have been scaled to the mean for each time series. Horizontal lines represent the soft and hard limits. 2000 = 1999–2000 fishing year. Error bars are 95% confidence intervals.



Annual relative exploitation rate for longfin eels in the Westland (AX) pre- and post-QMS. 2000 = 1999–2000 fishing year.

Fishery and Stock Trends	
Recent Trend in Biomass or Proxy	Pre-QMS CPUE declined from 1990–91 to 1992–93, and then increased steadily to 1999–2000. Post-QMS CPUE increased steadily from 2001–02 to 2012–13.
Recent Trend in Fishing intensity or Proxy	Relative exploitation rate declined steeply throughout the pre-QMS time series and generally declined from 2001–02 to 2008–09 before increasing to 2012–13 post-QMS.
Other Abundance Indices	-
Trends in Other Relevant Indicators or Variables	Catches of longfin elvers at primary monitoring sites have fluctuated without trend since the series of reliable data begins in 1995–96, suggesting no overall trend in recruitment.

Projections and Prognosis	
Stock Projections or Prognosis	Unlikely (< 40%) to decline in the medium term under current catch levels
Probability of Current Catch or TACC causing Biomass to remain below or to decline below Limits	Soft Limit: Unlikely (< 40%) if catch remains at current levels Hard Limit: Unlikely (< 40%) if catch remains at current levels South Island TACCs include both longfin and shortfin eels. As the TACC is substantially higher than the current longfin eel catch, it is not meaningful to evaluate potential impacts if catches of longfins increased to the level of the TACC.
Probability of Current Catch or TACC causing Overfishing to continue or to commence	Unknown if catch remains at current levels Likely (> 60%) if catch were to increase to the level of the TACC

Assessment Methodology and Evaluation		
Assessment Type	Level 2 – Partial Quantitative Stock Assessment	
Assessment Method	Standardised CPUE based on positive catches from commercial fyke net	
Assessment Dates	Latest assessment: 2014	Next assessment: 2019
Overall assessment quality rank	1 – High Quality	
Main data inputs (rank)	- Catch and effort data	1 – High Quality
Data not used (rank)	N/A	
Changes to Model Structure and Assumptions	-	
Major Sources of Uncertainty	<p>- Standardised CPUE only provides an index of abundance for eels in areas fished by commercial fishers. Other potential issues with the CPUE indices include:</p> <ul style="list-style-type: none"> • Low numbers of fishers • Uncertainty in target species after 2000 • Exclusion of zero catches • Changes in MLS and retention in early parts of the series (pre-QMS) 	


Qualifying Comments
<p>Because the commercial eel fishery has had a long history (beginning in the late 1960s), and indices of abundance are only available from the early 1990s, it is difficult to infer stock status from recent abundance trends, and these should therefore be interpreted with caution. Other sources of mortality, such as culling (primarily 1930s to 1950s) and habitat alteration (historical and current) have also reduced abundance prior to the CPUE series. The basis for the biological reference points is tenuous, and should be revised whenever new relevant information becomes available.</p> <p>The proportion of current longfin habitat in Westland (Statistical Area AX, ANG 11) fished commercially during the period 2009–10 and 2013–14 is estimated at 30% (Table 15). The proportion of virgin habitat impacted by hydro dams, commercial fishing and other anthropogenic activity was estimated to be 34%.</p>

Fishery Interactions
Bycatch of other species in the commercial eel fishery is low, and may include brown trout, galaxiids, yellow-eyed mullet, and koura in order of amount caught. Bycatch species are usually returned alive.


- Westland (AX) shortfin

Stock Status	
Year of Most Recent Assessment	2014
Assessment Runs Presented	Standardised CPUE
Reference Points	Target: B_{MSY} assumed, but not estimated Interim Soft Limit: Mean CPUE from 2001–02 to 2002–03 Hard Limit: 50% of Soft Limit Overfishing threshold: F_{MSY} assumed, but not estimated
Status in relation to Target	Unknown
Status in relation to Limits	Soft Limit: Unlikely (< 40%) to be below Hard Limit: Very Unlikely (< 10%) to be below
Status in relation to Overfishing	Unknown

Historical Stock Status Trajectory and Current Status



Comparison of standardised CPUE for shortfin eels in Westland (AX) from 1990–91 to 1999–2000 (pre-QMS) and 2001–02 to 2012–13 (post-QMS) (from Beentjes & Dunn 2015). Also shown is the total estimated shortfin catch in AX from ECERs. The two CPUE series have been scaled to the mean of each time series. Horizontal lines represent the soft and hard limits. 2000 = 1999–2000 fishing year. Error bars are 95% confidence intervals.



Annual relative exploitation rate for shortfin eels in the Westland (AX) pre- and post-QMS. 2000 = 1999–2000 fishing year.

Fishery and Stock Trends	
Recent Trend in Biomass or Proxy	Pre-QMS CPUE fluctuated without trend from 1990–91 to 1996–97 and then increased sharply to 1999–2000. Post-QMS CPUE increased steadily from 2001–02 to 2012–13.
Recent Trend in Fishing intensity or Proxy	Relative exploitation rate has shown large inter-annual fluctuations, with an increasing trend since 2003.
Other Abundance Indices	-
Trends in Other Relevant Indicators or Variables	Catches of shortfin elvers at primary monitoring sites have fluctuated without trend since the series of reliable data begins in 1995–96, suggesting no overall trend in recruitment.

Projections and Prognosis	
Stock Projections or Prognosis	Unlikely (< 40%) to decline in the medium term under current catch levels
Probability of Current Catch or TACC causing Biomass to remain below or to decline below Limits	<p>Soft Limit: Unlikely (< 40%) if catch remains at current levels</p> <p>Hard Limit: Very Unlikely (< 10%) if catch remains at current levels</p> <p>South Island TACCs include both longfin and shortfin eels. As the TACC is approximately 2–3 times higher than the current shortfin eel catch, it is not meaningful to evaluate potential impacts if catches of shortfins were to increase to the level of the TACC.</p>
Probability of Current Catch or TACC causing Overfishing to continue or to commence	<p>Unknown if catch remains at current levels</p> <p>Likely (> 60%) if catch were to increase to the level of the TACC</p>

Assessment Methodology and Evaluation		
Assessment Type	Level 2 - Partial Quantitative Stock Assessment	
Assessment Method	Standardised CPUE based on positive catches from commercial fyke net	
Assessment Dates	Latest assessment: 2014	Next assessment: 2019
Overall assessment quality rank	1 – High Quality	
Main data inputs (rank)	- Catch and effort data	1 – High Quality
Data not used (rank)	N/A	
Changes to Model Structure and Assumptions	-	
Major Sources of Uncertainty	<p>- Standardised CPUE only provides an index of abundance for eels in areas fished by commercial fishers. Other potential issues with the CPUE indices include:</p> <ul style="list-style-type: none"> • Low numbers of fishers • Uncertainty in target species after 2000 • Exclusion of zero catches • Changes in MLS and retention in early parts of the series (pre-QMS) 	


Qualifying Comments
Because the commercial eel fishery has had a long history (beginning in the late 1960s), and indices of abundance are only available from the early 1990s, it is difficult to infer stock status from recent abundance trends, and these should therefore be interpreted with caution. Other sources of mortality, such as culling (primarily 1930s to 1950s) and habitat alteration (historical and current) have also reduced abundance prior to the CPUE series. The basis for the biological reference points is tenuous, and should be revised whenever new relevant information becomes available.

Fishery Interactions
Bycatch of other species in the commercial eel fishery is low, and may include brown trout, galaxiids, yellow-eyed mullet, and koura in order of amount caught. Bycatch species are usually returned alive.


- Otago (AV) longfin

Stock Status	
Year of Most Recent Assessment	2014
Assessment Runs Presented	Standardised CPUE
Reference Points	Target: B_{MSY} assumed, but not estimated Interim Soft Limit: Mean CPUE from 2001–02 to 2002–03 Hard Limit: 50% of Soft Limit Overfishing threshold: F_{MSY} assumed, but not estimated
Status in relation to Target	Unknown
Status in relation to Limits	Soft Limit: Unlikely (< 40%) to be below Hard Limit: Very Unlikely (< 10%) to be below
Status in relation to Overfishing	Unknown

Historical Stock Status Trajectory and Current Status



Comparison of standardised CPUE for longfin eels in Otago (AV) from 1990–91 to 1999–2000 (pre-QMS) and 2001–02 to 2012–13 (post-QMS) (from Beentjes & Dunn 2015). Also shown is the total estimated longfin catch in AV from ECERs. The two CPUE series have been scaled to the mean of each time series. Horizontal lines represent the soft and hard limits. 2000 = 1999–2000 fishing year. Error bars are 95% confidence intervals.



Annual relative exploitation rate for longfin eels in the Otago (AV) pre- and post-QMS. 2000 = 1999–2000 fishing year.

Fishery and Stock Trends	
Recent Trend in Biomass or Proxy	Pre-QMS CPUE declined steadily from 1990–91 to 1995–96 and was stable to 1999–2000. Post-QMS CPUE is variable, but overall increased marginally from 2001–02 to 2012–13.
Recent Trend in Fishing intensity or Proxy	Relative exploitation rate declined markedly from 2002 to 2009 and then increased to the average for the post-QMS series.
Other Abundance Indices	-
Trends in Other Relevant Indicators or Variables	Catches of longfin elvers at primary monitoring sites have fluctuated without trend since the series of reliable data begins in 1995–96, suggesting no overall trend in recruitment.

Projections and Prognosis

Stock Projections or Prognosis	Unlikely (< 40%) to decline in the medium term if catch remains at current levels
--------------------------------	---

Probability of Current Catch or TACC causing Biomass to remain below or to decline below Limits	<p>Soft Limit: About as Likely as Not (40–60%) if catch remains at current levels</p> <p>Hard Limit: Unlikely (< 40%) if catch remains at current levels</p> <p>South Island TACCs include both longfin and shortfin eels. ANG 15 comprises statistical areas AV (Otago) and AW (Southland). As the TACC is substantially higher than the current longfin eel catch, it is not meaningful to evaluate potential impacts if catches were to increase to the level of the TACC.</p>
Probability of Current Catch or TACC causing Overfishing to continue or to commence	<p>Unknown if catch remains at current levels</p> <p>Unknown if catch were to increase to the level of the TACC</p>

Assessment Methodology		
Assessment Type	Level 2 – Partial Quantitative Stock Assessment	
Assessment Method	Standardised CPUE based on positive catches from commercial fyke net	
Assessment Dates	Latest assessment: 2014	Next assessment: 2019
Overall assessment quality rank	1 – High Quality	
Main data inputs (rank)	- Catch and effort data	1 – High Quality
Data not used (rank)	N/A	
Changes to Model Structure and Assumptions	-	
Major Sources of Uncertainty	<p>- Standardised CPUE only provides an index of abundance for eels in areas fished by commercial fishers. Other potential issues with the CPUE indices include:</p> <ul style="list-style-type: none"> • Low numbers of fishers • Uncertainty in target species after 2000 • Exclusion of zero catches • Changes in MLS and retention in early parts of the series (pre-QMS) 	


Qualifying Comments
<p>Because the commercial eel fishery has had a long history (beginning in the late 1960s), and indices of abundance are only available from the early 1990s, it is difficult to infer stock status from recent abundance trends, and these should therefore be interpreted with caution. Other sources of mortality, such as culling (primarily 1930s to 1950s) and habitat alteration (historical and current) have also reduced abundance prior to the CPUE series. The basis for the biological reference points is tenuous, and should be revised whenever new relevant information becomes available.</p> <p>The proportion of current longfin habitat in Otago (Statistical Area AV) fished commercially during the period 2009–10 and 2013–14 is estimated at 46% (Table 15). The proportion of virgin habitat impacted by hydro dams, commercial fishing and other anthropogenic activity was estimated to be 82.8%.</p>

Fishery Interactions
Bycatch of other species in the commercial eel fishery is low, and may include brown trout, galaxiids, yellow-eyed mullet, and koura in order of amount caught. Bycatch species are usually returned alive.


- Otago (AV) shortfin

Stock Status	
Year of Most Recent Assessment	2014
Assessment Runs Presented	Standardised CPUE
Reference Points	Target: B_{MSY} assumed, but not estimated Interim Soft Limit: Mean CPUE from 2001–02 to 2003–04 Hard Limit: 50% of Soft Limit Overfishing threshold: F_{MSY} assumed, but not estimated
Status in relation to Target	Unlikely (< 40%) to be at or above
Status in relation to Limits	Soft Limit: About as Likely as Not (40–60%) to be below Hard Limit: Unlikely (< 40%) to be below
Status in relation to Overfishing	Unknown

Historical Stock Status Trajectory and Current Status



Comparison of standardised CPUE for shortfin eels in Otago (AV) from 1990–91 to 1999–2000 (pre-QMS) and 2001–02 to 2012–13 (post-QMS) (from Beentjes & Dunn 2015). Also shown is the total estimated shortfin catch in AV from ECERs. The two CPUE series have been scaled to the mean of each time series. Horizontal lines represent the soft and hard limits. 2000 = 1999–2000 fishing year. Error bars are 95% confidence intervals.



Annual relative exploitation rate for shortfin eels in the Otago (AV) pre- and post-QMS. 2000 = 1999–2000 fishing year.

Fishery and Stock Trends	
Recent Trend in Biomass or Proxy	Pre-QMS CPUE declined steadily from 1990–91 to 1998–99 and then increased slightly to 1999–2000. Post-QMS CPUE increased steadily from 2001–02 to 2010–11, and then declined markedly to just below the long-term average.
Recent Trend in Fishing intensity or Proxy	Relative exploitation rate has fluctuated without trend since 2002.
Other Abundance Indices	-
Trends in Other Relevant Indicators or Variables	Catches of shortfin elvers at primary monitoring sites have fluctuated without trend since the series of reliable data begins in 1995–96, suggesting no overall trend in recruitment.

Projections and Prognosis	
Stock Projections or Prognosis	As both catch and exploitation rate show large inter-annual variation, it is not clear whether the population will continue to decline.
Probability of Current Catch or TACC causing Biomass to remain below or to decline below Limits	Soft Limit: About as Likely as Not (40–60%) if catch remains at current levels Hard Limit: Unlikely (< 40%) if catch remains at current levels South Island TACCs include both longfin and shortfin eels. ANG 15 comprises statistical areas AV (Otago) and AW (Southland). The TACC is 6–7 fold higher than the current shortfin eel catch in ANG 15. Catch at the level of the TACC is Likely (> 60%) to cause decline below both the soft and hard Limits
Probability of Current Catch or TACC causing Overfishing to continue or to commence	Unknown if catch remains at current levels Likely (> 40%) if catch were to increase to the level of the TACC

Assessment Methodology and Evaluation		
Assessment Type	Level 2 – Partial Quantitative Stock Assessment	
Assessment Method	Standardised CPUE based on positive catches from commercial fyke net	
Assessment Dates	Latest assessment: 2014	Next assessment: 2019
Overall assessment quality rank	1 – High Quality	
Main data inputs (rank)	- Catch and effort data	1 – High Quality
Data not used (rank)	N/A	
Changes to Model Structure and Assumptions	-	
Major Sources of Uncertainty	- Standardised CPUE only provides an index of abundance for eels in areas fished by commercial fishers. Other potential issues with the CPUE indices include: <ul style="list-style-type: none"> • Low numbers of fishers • Uncertainty in target species after 2000 • Exclusion of zero catches • Changes in MLS and retention in early parts of the series (pre-QMS) 	


Qualifying Comments
Because the commercial eel fishery has had a long history (beginning in the late 1960s), and indices of abundance are only available from the early 1990s, it is difficult to infer stock status from recent abundance trends, and these should therefore be interpreted with caution. Other sources of mortality, such as culling (primarily 1930s to 1950s) and habitat alteration (historical and current) have also reduced abundance prior to the CPUE series.

Fishery Interactions
Bycatch of other species in the commercial eel fishery is low, and may include: brown trout, black flounder, koura, yellow-eyed mullet, galaxiids, yellowbelly flounder, and bullies in order of amount caught. Bycatch species are usually returned alive.


- Southland (AW) longfin

Stock Status	
Year of Most Recent Assessment	2014
Assessment Runs Presented	Standardised CPUE
Reference Points	Target: B_{MSY} assumed, but not estimated Interim Soft Limit: Mean CPUE from 2006–07 to 2009–10 Hard Limit: 50% of Soft Limit Overfishing threshold: F_{MSY} assumed, but not estimated
Status in relation to Target	Unknown
Status in relation to Limits	Soft Limit: Unlikely (< 40%) to be below Hard Limit: Very Unlikely (< 10%) to be below
Status in relation to Overfishing	Unknown

Historical Stock Status Trajectory and Current Status



Comparison of standardised CPUE for longfin eels in Southland (AW) from 1990–91 to 1999–2000 (pre-QMS) and 2001–02 to 2012–13 (post-QMS) (from Beentjes & Dunn 2015). Also shown is the total estimated longfin catch in AW from ECERs. The two CPUE series have been scaled to the mean of each time series. Horizontal lines represent the soft and hard limits. 2000 = 1999–2000 fishing year. Error bars are 95% confidence intervals.



Annual relative exploitation rate for longfin eels in the Southland (AW) pre- and post-QMS. 2000 = 1999–2000 fishing year.

Fishery and Stock Trends	
Recent Trend in Biomass or Proxy	Pre-QMS CPUE declined steadily from 1990–91 to 1998–98 and increased to 1999–2000. Post-QMS CPUE is variable and showed a gradual decline from 2001–02 to 2009–10, then an increase since.
Recent Trend in Fishing intensity or Proxy	Relative exploitation rate declined from 2002 to 2010 and then increased steeply to well above the long-term average to 2013.
Other Abundance Indices	-
Trends in Other Relevant Indicators or Variables	Catches of longfin elvers at primary monitoring sites have fluctuated without trend since the series of reliable data begins in 1995–96, suggesting no overall trend in recruitment.

Projections and Prognosis

Stock Projections or Prognosis	Likely (> 60%) to decline under recent levels of catch and
--------------------------------	--

	exploitation rate
Probability of Current Catch or TACC causing Biomass to remain below or to decline below Limits	<p>Soft Limit: Unlikely (< 40%) if catch remains at current levels</p> <p>Hard Limit: Unlikely (< 40%) if catch remains at current levels</p> <p>South Island TACCs include both longfin and shortfin eels. ANG 15 comprises statistical areas AV (Otago) and AW (Southland). As the TACC is substantially higher than the current longfin eel catch, it is not meaningful to evaluate potential impacts if catches increased to the level of the TACC.</p>
Probability of Current Catch or TACC causing Overfishing to continue or to commence	<p>Unknown if catch remains at current levels</p> <p>Very Likely (> 90%) if catch were to increase to the level of the TACC</p>

Assessment Methodology and Evaluation		
Assessment Type	Level 2 - Partial Quantitative Stock Assessment	
Assessment Method	Standardised CPUE based on positive catches from commercial fyke net	
Assessment Dates	Latest assessment: 2014	Next assessment: 2019
Overall assessment quality rank	1 – High Quality	
Main data inputs (rank)	- Catch and effort data	1 – High Quality
Data not used (rank)	N/A	
Changes to Model Structure and Assumptions	-	
Major Sources of Uncertainty	<p>- Standardised CPUE only provides an index of abundance for eels in areas fished by commercial fishers. Other potential issues with the CPUE indices include:</p> <ul style="list-style-type: none"> • Low numbers of fishers • Uncertainty in target species after 2000 • Exclusion of zero catches • Changes in MLS and retention in early parts of the series (pre-QMS) 	


Qualifying Comments
<p>Because the commercial eel fishery has had a long history (beginning in the late 1960s), and indices of abundance are only available from the early 1990s, it is difficult to infer stock status from recent abundance trends, and these should therefore be interpreted with caution. Other sources of mortality, such as culling (primarily 1930s to 1950s) and habitat alteration (historical and current) have also reduced abundance prior to the CPUE series. The basis for the biological reference points is tenuous, and should be revised whenever new relevant information becomes available.</p> <p>The proportion of current longfin habitat in Southland (Statistical Area AW) fished commercially during the period 2009–10 and 2013–14 is estimated at 32% (Table 15). The proportion of virgin habitat impacted by hydro dams, commercial fishing and other anthropogenic activity was estimated to be 41%.</p>

Fishery Interactions
Bycatch of other species in the commercial eel fishery is low, and may include brown trout, giant bullies, koura, galaxiids, and common bullies in order of amount caught. Bycatch species are usually returned alive.


- Southland (AW) shortfin

Stock Status	
Year of Most Recent Assessment	2014
Assessment Runs Presented	Standardised CPUE
Reference Points	Target: B_{MSY} assumed, but not estimated Interim Soft Limit: Mean CPUE from 2001–02 to 2002–03 Hard Limit: 50% of Soft Limit Overfishing threshold: F_{MSY} assumed, but not estimated
Status in relation to Target	Unknown
Status in relation to Limits	Soft Limit: Unlikely (< 40%) to be below Hard Limit: Very Unlikely (< 10%) to be below
Status in relation to Overfishing	Unknown

Historical Stock Status Trajectory and Current Status



Comparison of standardised CPUE for shortfin eels in Southland (AW) from 1990–91 to 1999–2000 (pre-QMS) and 2001–02 to 2012–13 (post-QMS) (from Beentjes & Dunn 2015). Also shown is the total estimated shortfin catch in AW from ECERs. The two CPUE series have been scaled to the mean of each time series. Horizontal lines represent the soft and hard limits. 2000 = 1999–2000 fishing year. Error bars are 95% confidence intervals.



Annual relative exploitation rate for shortfin eels in the Southland (AW) pre- and post-QMS. 2000 = 1999–2000 fishing year.

Fishery and Stock Trends	
Recent Trend in Biomass or Proxy	Pre-QMS CPUE declined slowly from 1990–91 to 1996–97 and then gradually increased to 1999–2000. Post-QMS CPUE fluctuated but increased substantially from 2001–02 to 2012–13.
Recent Trend in Fishing intensity or Proxy	Relative exploitation rate shows high inter-annual variation, but a consistently declining trend since 2002.
Other Abundance Indices	-
Trends in Other Relevant Indicators or Variables	Catches of shortfin elvers at primary monitoring sites have fluctuated without trend since the series of reliable data begins in 1995–96, suggesting no overall trend in recruitment.

Projections and Prognosis	
Stock Projections or Prognosis	Likely (> 60%) to continue to increase in the medium term under current catch levels
Probability of Current Catch or TACC causing Biomass to remain below or to decline below Limits	Soft Limit: Unlikely (< 40%) if the catch remains at current levels Hard Limit: Very Unlikely (< 10%) if the catch remains at current levels South Island TACCs include both longfin and shortfin eels. ANG 15 comprises statistical areas AV (Otago) and AW (Southland). As the TACC is substantially higher than the current longfin eel catch, it is not meaningful to evaluate potential impacts if catches increased to the level of the TACC.
Probability of Current Catch or TACC causing Overfishing to continue or to commence	Unknown if catch remains at current levels Likely (> 60%) if catch were to increase to the level of the TACC

Assessment Methodology and Evaluation		
Assessment Type	Level 2 – Partial Quantitative Stock Assessment	
Assessment Method	Standardised CPUE based on positive catches from commercial fyke net	
Assessment Dates	Latest assessment: 2014	Next assessment: 2019
Overall assessment quality rank	1 – High Quality	
Main data inputs (rank)	- Catch and effort data	1 – High Quality
Data not used (rank)	N/A	
Changes to Model Structure and Assumptions	-	
Major Sources of Uncertainty	- Standardised CPUE only provides an index of abundance for eels in areas fished by commercial fishers. Other potential issues with the CPUE indices include: <ul style="list-style-type: none"> • Low numbers of fishers • Uncertainty in target species after 2000 • Exclusion of zero catches • Changes in MLS and retention in early parts of the series (pre-QMS) 	


Qualifying Comments
Because the commercial eel fishery has had a long history (beginning in the late 1960s), and indices of abundance are only available from the early 1990s, it is difficult to infer stock status from recent abundance trends, and these should therefore be interpreted with caution. Other sources of mortality, such as culling (primarily 1930s to 1950s) and habitat alteration (historical and current) have also reduced abundance prior to the CPUE series. The basis for the biological reference points is tenuous, and should be revised whenever new relevant information becomes available.

Fishery Interactions
Bycatch of other species in the commercial eel fishery is low, and may include brown trout, giant bullies, koura, galaxiids, and common bullies in order of amount caught. Bycatch species are usually returned alive.


- Te Waihora (AS1) shortfin

Stock Status	
Year of Most Recent Assessment	2014
Assessment Runs Presented	Standardised CPUE of feeder eels in AS1
Reference Points	Interim Target: B_{MSY} -compatible proxy based on mean CPUE for the period: 2006–07 to 2009–10. Soft Limit: 50% of target Hard Limit: 50% of soft limit Overfishing threshold: F_{MSY}
Status in relation to Target	Very Likely (> 60%) to be at or above B_{MSY}
Status in relation to Limits	Soft Limit: Very Unlikely (< 10%) to be below Hard Limit: Very Unlikely (< 10%) to be below
Status in relation to Overfishing	Overfishing is Very Unlikely (< 10%) to be occurring

Historical Stock Status Trajectory and Current Status



Comparison of standardised CPUE for shortfin eels in Te Waihora (AS1) from 2001–02 to 2012–13 (post-QMS) (from Beentjes & Dunn 2015). Also shown is the total estimated shortfin catch in AS1 from ECERs. The CPUE series have been scaled to the mean of each time series. Horizontal lines represent the target, and soft and hard limits. 2002 = 2001–2002 fishing year. Error bars are 95% confidence intervals.



Annual relative exploitation rate for shortfin eels in the Te Waihora (AS1) post-QMS. 2002 = 2001–02 fishing year.

Fishery and Stock Trends	
Recent Trend in Biomass or Proxy	CPUE of feeder shortfin eels in Te Waihora (AS1) increased 6-fold from 2001–02 to 2010–11, but showed no trend to 2012–13.
Recent Trend in Fishing intensity or Proxy	Relative exploitation rate has declined substantially (9-fold) since 2002, and is now well below the series average.

Other Abundance Indices	-
Trends in Other Relevant Indicators or Variables	Catches of shortfin elvers at primary monitoring sites have fluctuated without trend since the series of reliable data begins in 1995–96, suggesting no overall trend in recruitment. Increasing mean size since the mid-1990s suggests reduced exploitation rates.

Projections and Prognosis	
Stock Projections or Prognosis	Likely (> 60%) to remain well above the target in the medium term under current catch levels
Probability of Current Catch or TACC causing Biomass to remain below or to decline below Limits	Soft Limit: Very Unlikely (< 10%) if catch remains at current levels Hard Limit: Very Unlikely (< 10%) if catch remains at current levels Unlikely (< 40%) if catch were to increase to the level of the TACC, provided not all of the catch is taken from AS1
Probability of Current Catch or TACC causing Overfishing to continue or to commence	Unlikely (< 40%) if catch remains at current levels Unlikely (< 40%) if catch were to increase to the level of the TACC, provided not all of the catch is taken from AS1

Assessment Methodology and Evaluation		
Assessment Type	Level 2 – Partial Quantitative Stock Assessment	
Assessment Method	Standardised CPUE based on positive catches from commercial fyke net	
Assessment Dates	Latest assessment: 2014	Next assessment: 2019
Overall assessment quality rank	1 – High Quality	
Main data inputs (rank)	- Catch and effort data	1 – High Quality
Data not used (rank)	N/A	
Changes to Model Structure and Assumptions	-	
Major Sources of Uncertainty	- Standardised CPUE only provides an index of abundance for eels in areas fished by commercial fishers. Other potential issues with the CPUE indices include: <ul style="list-style-type: none"> • Low numbers of fishers • Exclusion of zero catches • Changes in MLS and retention in early parts of the series (pre-QMS) 	

Qualifying Comments
<p>Because the commercial eel fishery has had a long history (beginning in the late 1960s), and indices of abundance are only available from the early 1990s, it is difficult to infer stock status from recent abundance trends, and these should therefore be interpreted with caution. Other sources of mortality, such as habitat alteration (historical and current) have also reduced abundance prior to the CPUE series.</p> <p>The shortfin eel catch from Te Waihora comprises small migrant males from AS2 and feeder females from AS1. The index of abundance is based on the catch rates of feeder eels. The basis for the biological reference points is tenuous, and should be revised whenever new relevant information becomes available.</p> <p>Shortfin eels in Te Waihora have a markedly different (mostly strongly increasing) pattern in CPUE compared to other eel sub-populations. This could be due to a number of factors, both positive and negative, including eutrophication, and changes in productivity, lake opening regimes, and management measures.</p>

Fishery Interactions

Bycatch of other species in the commercial eel fishery may include: bullies, black flounder, yellowbelly flounder, sand flounder, and goldfish in order of the amount caught. The flatfish species are usually released alive or retained if caught under quota. Longfin eels are not abundant and are usually voluntarily released alive. All other bycatch is released alive.

Status of North Island Eels

Level 2 Partial Quantitative Stock Assessments are conducted by statistical area and species where accepted indices of abundance are available. Standardised CPUE provides information on the abundance of commercially harvested eels (300 g–4000 g) in areas that are fished commercially.


Approximately 73% of current longfin habitat on the North Island is either in reserves or in areas rarely or never fished by commercial fishers. Statements regarding the status of longfin eels in relation to reference points are made separately for the entire ESA and for the area commercially fished within it. There is no information available on the proportion of shortfin habitat in each ESA that is fished commercially.

QMA SFE 20 and LFE 20 (includes ESAs AA and AB)


- **Northland (AA) shortfin**

Stock Status

Year of Most Recent Assessment	2017
Assessment Runs Presented	Standardised CPUE on positive catch
Reference Points	Target: B_{MSY} proxy based on CPUE; not determined Default Soft Limit: 20% B_0 Default Hard Limit: 10% B_0 Overfishing threshold: F_{MSY} proxy based on relative exploitation rate; not determined
Status in relation to Target	Unknown
Status in relation to Limits	Soft Limit: Unknown Hard Limit: Unknown
Status in relation to Overfishing	Unknown

Historical Stock Status Trajectory and Current Status

Standardised CPUE for shortfin eels in Northland (AA) from 1990–91 to 2014–15 (from Beentjes & McKenzie 2017). Also shown is the total estimated core fisher shortfin catch in AA from ECERs. Error bars are 95% confidence intervals. Before 2001, 37% of the catch was recorded as EEU (unidentified) and these catches are omitted. 2000 = 1999–2000 fishing year.



Annual relative exploitation rate for shortfin eels in the Northland (AA). Because some catch of shortfin was reported as EEU (unidentified) and has not been allocated to species, the exploitation rate is likely to have been higher than shown before 2001.

Fishery and Stock Trends

Recent Trend in Biomass or Proxy	Increasing trend in CPUE since early 1990s, but relatively stable over the most recent 6 years
Recent Trend in Fishing intensity or Proxy	The relative exploitation rate has declined steeply since 2003 and in 2015 was well below the series mean
Other Abundance Indices	-
Trends in Other Relevant Indicators or Variables	Catches of shortfin elvers at primary monitoring sites have fluctuated without trend since the series of reliable data begins in 1995–96, suggesting no overall trend in recruitment.

Projections and Prognosis

Stock Projections or Prognosis	Unlikely (< 40%) to decline in the medium term under current catch levels
Probability of Current Catch or TACC causing Biomass to remain below or to decline below Limits	Unknown
Probability of Current Catch or TACC causing Overfishing to continue or to commence	Unknown

Assessment Methodology and Evaluation

Assessment Type	Level 2 – Partial Quantitative Stock Assessment	
Assessment Method	Standardised CPUE based on positive catches from commercial fyke net	
Assessment Dates	Latest assessment: 2017	Next assessment: 2019
Overall assessment quality rank	1 – High Quality	
Main data inputs (rank)	- Catch and effort data	1 – High Quality
Data not used (rank)	N/A	
Changes to Model Structure and Assumptions	-	
Major Sources of Uncertainty	- Standardised CPUE only provides an index of abundance for eels in areas fished by commercial fishers. Other potential issues with the CPUE indices include: <ul style="list-style-type: none"> • Low numbers of fishers • Uncertainty in target species after 2000 • Exclusion of zero catches 	

	<ul style="list-style-type: none"> Changes in MLS and retention in early parts of the series and increased escape tube size from 25 mm to 31 mm in 2012–13 Failure of some fishers to record on ECE returns all legal sized eels caught, not just those retained
--	--

Qualifying Comments

Because the commercial eel fishery has had a long history (beginning in the late 1960s), and indices of abundance are only available from the early 1990s, it is difficult to infer stock status from recent abundance trends, and these should therefore be interpreted with caution. Other sources of mortality, such as habitat alteration (historical and current) have also reduced abundance prior to the CPUE series.


Fishery Interactions

Bycatch of other species in the commercial Northland eel fishery includes mainly catfish, with lesser quantities of koura, goldfish and perch. Most bycatch species are usually returned alive.


- Northland (AA) longfin**

Stock Status

Year of Most Recent Assessment	2017
Assessment Runs Presented	Standardised CPUE on positive catch
Reference Points	<p>For ESA, Interim Target is 40% B_0</p> <p>For commercially fished area, Target is B_{MSY} proxy based on CPUE; not determined</p> <p>Default Soft Limit: 20% B_0</p> <p>Default Hard Limit: 10% B_0</p> <p>For ESA, Overfishing threshold is F_{MSY}</p> <p>For commercially fished area, Overfishing threshold is F_{MSY} proxy based on relative exploitation rate; not determined</p>
Status in relation to Target	<p>For total ESA: Likely (> 60%) to be at or above</p> <p>For fished area: Unknown</p>
Status in relation to Limits	<p>For ESA, Soft Limit: Very Unlikely (< 10%) to be below</p> <p>For ESA, Hard Limit: Very Unlikely (< 10%) to be below</p>
Status in relation to Overfishing	<p>For ESA: Unlikely (< 40%) to be overfishing</p> <p>For fished area: Unknown</p>

Historical Stock Status Trajectory and Current Status

Standardised CPUE for longfin eels in Northland (AA) from 1990–91 to 2014–15 (from Beentjes & McKenzie 2017). Also shown is the total estimated core fisher longfin catch in AA from ECERs. Error bars are 95% confidence intervals. Vertical dashed line indicates when the 4 kg maximum size was introduced in 2007–08 after which longfin eels 4 kg and over are not recorded on ECERs. Before 2001, 37% of the catch was recorded as EEU (unidentified) and these catches are omitted.



Annual relative exploitation rate for longfin eels in the Northland (AA). Because some catch of longfin was reported as EEU (unidentified) and has not been allocated to species, the exploitation rate is likely to have been higher than shown before 2001.

Fishery and Stock Trends

Recent Trend in Biomass or Proxy	Very slight downward trend in CPUE over the time series
Recent Trend in Fishing intensity or Proxy	The relative exploitation rate has declined steeply since 2002 and in 2015 was well below the series mean.
Other Abundance Indices	-
Trends in Other Relevant Indicators or Variables	Catches of longfin elvers at primary monitoring sites have fluctuated without trend since the series of reliable data begins in 1995–96, suggesting no overall trend in recruitment.

Projections and Prognosis

Stock Projections or Prognosis	Unlikely (< 40%) to decline in the medium term under current catch levels
Probability of Current Catch or TACC causing Biomass to remain below or to decline below Limits	For ESA, Soft Limit: Very Unlikely (< 10%) if catch remains at current levels For ESA, Hard Limit: Very Unlikely (< 10%) if catch remains at current levels
Probability of Current Catch or TACC causing Overfishing to continue or to commence	For ESA, Unlikely (< 40%) if catch remains at current levels

Assessment Methodology and Evaluation

Assessment Type	Level 2 – Partial Quantitative Stock Assessment	
Assessment Method	Standardised CPUE based on positive catches from commercial fyke net	
Assessment Dates	Latest assessment: 2017	Next assessment: 2019
Overall assessment quality rank	1 – High Quality	
Main data inputs (rank)	- Catch and effort data	1 – High Quality
Data not used (rank)	N/A	
Changes to Model Structure and Assumptions	-	
Major Sources of Uncertainty	- Standardised CPUE only provides an index of abundance for eels in areas fished by commercial fishers. Other potential issues with the CPUE indices include: <ul style="list-style-type: none"> • Low numbers of fishers (for some ESAs) • Uncertainty in the method used to derive target species 	

	<ul style="list-style-type: none"> • Changes in MLS and retention in early parts of the series and increased escape tube size from 25 mm to 31 mm in 2012–13 • Failure of some fishers to record on ECE returns all legal sized eels caught, not just those retained • Unrecorded release of > 4kg eels since 2007–08
--	---

Qualifying Comments

Because the commercial eel fishery has had a long history (beginning in the late 1960s), and indices of abundance are only available from the early 1990s, it is difficult to infer stock status from recent abundance trends, and these should therefore be interpreted with caution. Other sources of mortality, such as habitat alteration (historical and current) have also reduced abundance prior to the CPUE series.

The proportion of current longfin habitat in Northland (Statistical Area AA) fished commercially during the period 2009–10 and 2013–14 is estimated at 36% (Table 15) The proportion of virgin habitat impacted by hydro dams, commercial fishing and other anthropogenic activity was estimated to be 40%.


Fishery Interactions

Bycatch of other species in the commercial Northland eel fishery includes mainly catfish, with lesser quantities of koura, goldfish and perch. Most bycatch species are usually returned alive.


• Auckland (AB) shortfin

Stock Status	
Year of Most Recent Assessment	2017
Assessment Runs Presented	Standardised CPUE on positive catch
Reference Points	Target: B_{MSY} proxy based on CPUE; not determined Default Soft Limit: 20% B_0 Default Hard Limit: 10% B_0 Overfishing threshold: F_{MSY} proxy based on relative exploitation rate; not determined
Status in relation to Target	Unknown
Status in relation to Limits	Soft Limit: Unknown Hard Limit: Unknown
Status in relation to Overfishing	Unknown

Historical Stock Status Trajectory and Current Status



Standardised CPUE for shortfin eels in Auckland (AB) from 1990–91 to 2014–15 (from Beentjes & McKenzie 2017). Also shown is the total estimated core fisher shortfin catch in AB from ECERs. Error bars are 95% confidence intervals. Before 2000, 26% of the catch was recorded as EEU (unidentified) and these catches are omitted.



Annual relative exploitation rate for shortfin eels in the Auckland (AB). Because some catch of shortfin was reported as EEU (unidentified) and has not been allocated to species, the exploitation rate is likely to have been higher than shown before 2000.

Fishery and Stock Trends

Recent Trend in Biomass or Proxy	No trend in CPUE until 2003, after which it increases consistently
Recent Trend in Fishing intensity or Proxy	The relative exploitation rate declined from 2012 and in 2015 was below the series mean.
Other Abundance Indices	-
Trends in Other Relevant Indicators or Variables	Catches of shortfin elvers at primary monitoring sites have fluctuated without trend since the series of reliable data begins in 1995–96, suggesting no overall trend in recruitment.

Projections and Prognosis

Stock Projections or Prognosis	Unlikely (< 40%) to decline in the medium term under current catch levels
Probability of Current Catch or TACC causing Biomass to remain below or to decline below Limits	Unknown
Probability of Current Catch or TACC causing Overfishing to continue or to commence	Unknown

Assessment Methodology and Evaluation

Assessment Type	Level 2 – Partial Quantitative Stock Assessment	
Assessment Method	Standardised CPUE based on positive catches from commercial fyke net	
Assessment Dates	Latest assessment: 2017	Next assessment: 2019
Overall assessment quality rank	1 – High Quality	
Main data inputs (rank)	- Catch and effort data	1 – High Quality
Data not used (rank)	N/A	
Changes to Model Structure and Assumptions	-	
Major Sources of Uncertainty	- Standardised CPUE only provides an index of abundance for eels in areas fished by commercial fishers. Other potential issues with the CPUE indices include: <ul style="list-style-type: none"> • Low numbers of fishers • Uncertainty in target species after 2000 • Exclusion of zero catches 	

	<ul style="list-style-type: none"> • Changes in MLS and retention in early parts of the series and increased escape tube size from 25 mm to 31 mm in 2012–13 • Failure of some fishers to record on ECE returns all legal sized eels caught, not just those retained
--	--

Qualifying Comments

Because the commercial eel fishery has had a long history (beginning in the late 1960s), and indices of abundance are only available from the early 1990s, it is difficult to infer stock status from recent abundance trends, and these should therefore be interpreted with caution. Other sources of mortality, such as habitat alteration (historical and current) have also reduced abundance prior to the CPUE series.


Fishery Interactions

Bycatch of other species in the commercial Auckland eel fishery includes mainly catfish, with lesser quantities of Koi carp, goldfish, koura, grey mullet and yellowbelly flounder. Most bycatch species are usually returned alive.

- **Auckland (AB) longfin**


Stock Status

Year of Most Recent Assessment	2017
Assessment Runs Presented	Standardised CPUE on positive catch
Reference Points	<p>For ESA, Interim Target is 40% B_0</p> <p>For commercially fished area, Target is B_{MSY} proxy based on CPUE; not determined</p> <p>Default Soft Limit: 20% B_0</p> <p>Default Hard Limit: 10% B_0</p> <p>For ESA, Overfishing threshold is F_{MSY}</p> <p>For commercially fished area, Overfishing threshold is F_{MSY} proxy based on relative exploitation rate; not determined</p>
Status in relation to Target	<p>For total ESA: Likely (> 60%) to be at or above</p> <p>For fished area: Unknown</p>
Status in relation to Limits	<p>For ESA, Soft Limit: Very Unlikely (< 10%) to be below</p> <p>For ESA, Hard Limit: Very Unlikely (< 10%) to be below</p>
Status in relation to Overfishing	<p>For ESA: Unlikely (< 40%) to be overfishing</p> <p>For fished area: Unknown</p>

Historical Stock Status Trajectory and Current Status

Comparison of standardised CPUE for longfin eels in Northland (AB) from 1990–91 to 2014–15 (from Beentjes & McKenzie 2017). Also shown is the total estimated core fisher longfin catch in AB from ECERs. Vertical dashed line indicates when the 4 kg maximum size was introduced in 2007–08 after which longfin eels 4 kg and over are

not recorded on ECERs. Error bars are 95% confidence intervals. Before 2000, 26% of the catch was recorded as EEU (unidentified) and these catches are omitted.



Annual relative exploitation rate for longfin eels in the Auckland (AB). Because some catch of longfin was reported as EEU (unidentified) and has not been allocated to species, the exploitation rate is likely to have been higher than shown before 2000.

Fishery and Stock Trends

Recent Trend in Biomass or Proxy	A slight decline in CPUE to 2005, but stable thereafter
Recent Trend in Fishing intensity or Proxy	The relative exploitation rate has declined since 2013 and in 2015 was below the series mean.
Other Abundance Indices	-
Trends in Other Relevant Indicators or Variables	Catches of longfin elvers at primary monitoring sites have fluctuated without trend since the series of reliable data begins in 1995–96, suggesting no overall trend in recruitment.

Projections and Prognosis

Stock Projections or Prognosis	Unlikely (< 40%) to decline in the medium term under current catch levels
Probability of Current Catch or TACC causing Biomass to remain below or to decline below Limits	For ESA, Soft Limit: Very Unlikely (< 10%) if catch remains at current levels For ESA, Hard Limit: Very Unlikely (< 10%) if catch remains at current levels
Probability of Current Catch or TACC causing Overfishing to continue or to commence	For ESA, Unlikely (< 40%) if catch remains at current levels

Assessment Methodology and Evaluation

Assessment Type	Level 2 – Partial Quantitative Stock Assessment	
Assessment Method	Standardised CPUE based on positive catches from commercial fyke net	
Assessment Dates	Latest assessment: 2017	Next assessment: 2019
Overall assessment quality rank	1 – High Quality	
Main data inputs (rank)	- Catch and effort data	1 – High Quality
Data not used (rank)	N/A	
Changes to Model Structure and Assumptions	-	
Major Sources of Uncertainty	- Standardised CPUE only provides an index of abundance for eels in areas fished by commercial fishers. Other potential issues with the CPUE indices include: <ul style="list-style-type: none"> • Low numbers of fishers • Uncertainty in target species after 2000 • Exclusion of zero catches 	

FRESHWATER EELS (SFE, LFE, ANG)

	<ul style="list-style-type: none"> • Changes in MLS and retention in early parts of the series and increased escape tube size from 25 mm to 31 mm in 2012–13 • Failure of some fishers to record on ECE returns all legal sized eels caught, not just those retained • Unrecorded release of > 4kg eels since 2007–08
--	---

Qualifying Comments

Because the commercial eel fishery has had a long history (beginning in the late 1960s), and indices of abundance are only available from the early 1990s, it is difficult to infer stock status from recent abundance trends, and these should therefore be interpreted with caution. Other sources of mortality, such as habitat alteration (historical and current) have also reduced abundance prior to the CPUE series.

The proportion of current longfin habitat in Auckland (Statistical Area AB) fished commercially during the period 2009–10 and 2013–14 is estimated at 35% (Table 15). The proportion of virgin habitat impacted by hydro dams, commercial fishing and other anthropogenic activity was estimated to be 38%.

Fishery Interactions


Bycatch of other species in the commercial Auckland eel fishery includes mainly catfish, with lesser quantities of Koi carp, goldfish, koura, grey mullet and yellowbelly flounder. Most bycatch species are usually returned alive.

QMA SFE 21 and LFE 21 (includes ESAs AC, AD, AE and AF)


• Hauraki (AC) shortfin

Stock Status	
Year of Most Recent Assessment	2017
Assessment Runs Presented	Standardised CPUE on positive catch
Reference Points	Target: B_{MSY} proxy based on CPUE; not determined Default Soft Limit: 20% B_0 Default Hard Limit: 10% B_0 Overfishing threshold: F_{MSY} proxy based on relative exploitation rate; not determined
Status in relation to Target	Unknown
Status in relation to Limits	Soft Limit: Unknown Hard Limit: Unknown
Status in relation to Overfishing	Unknown

Historical Stock Status Trajectory and Current Status



Standardised CPUE for shortfin eels in Hauraki (AC) from 1990–91 to 2014–15 (from Beentjes & McKenzie 2017). Also shown is the total estimated core fisher shortfin catch in AC from ECERs. Error bars are 95% confidence intervals. Before 2002, 16% of the catch was recorded as EEU (unidentified) and these catches are omitted.



Annual relative exploitation rate for shortfin eels in the Hauraki (AC). Because some catch of shortfin was reported as EEU (unidentified) and has not been allocated to species, the exploitation rate is likely to have been higher than shown before 2002.

Fishery and Stock Trends

Recent Trend in Biomass or Proxy	No trend in CPUE until 2010, after which it has increased
Recent Trend in Fishing intensity or Proxy	The relative exploitation rate has declined since 2006, and in 2015 was below the series mean.
Other Abundance Indices	-
Trends in Other Relevant Indicators or Variables	Catches of shortfin elvers at primary monitoring sites have fluctuated without trend since the series of reliable data begins in 1995–96, suggesting no overall trend in recruitment.

Projections and Prognosis

Stock Projections or Prognosis	Unlikely (< 40%) to decline in the medium term under current catch levels
Probability of Current Catch or TACC causing Biomass to remain below or to decline below Limits	Unknown
Probability of Current Catch or TACC causing Overfishing to continue or to commence	Unknown

Assessment Methodology and Evaluation

Assessment Type	Level 2 – Partial Quantitative Stock Assessment	
Assessment Method	Standardised CPUE based on positive catches from commercial fyke net	
Assessment Dates	Latest assessment: 2017	Next assessment: 2019
Overall assessment quality rank	1 – High Quality	
Main data inputs (rank)	- Catch and effort data	1 – High Quality
Data not used (rank)	N/A	
Changes to Model Structure and Assumptions	-	
Major Sources of Uncertainty	- Standardised CPUE only provides an index of abundance for eels in areas fished by commercial fishers. Other potential issues with the CPUE indices include: <ul style="list-style-type: none"> • Low numbers of fishers • Uncertainty in target species after 2000 • Exclusion of zero catches 	

	<ul style="list-style-type: none"> • Changes in MLS and retention in early parts of the series and increased escape tube size from 25 mm to 31 mm in 2012–13 • Failure of some fishers to record on ECE returns all legal sized eels caught, not just those retained
--	--

Qualifying Comments

Because the commercial eel fishery has had a long history (beginning in the late 1960s), and indices of abundance are only available from the early 1990s, it is difficult to infer stock status from recent abundance trends, and these should therefore be interpreted with caution. Other sources of mortality, such as habitat alteration (historical and current) have also reduced abundance prior to the CPUE series.


Fishery Interactions

Bycatch of other species in the commercial Hauraki eel fishery includes mainly catfish, with lesser quantities of brown trout, goldfish, koi carp, and kokopu. Most bycatch species are usually returned alive.

- **Hauraki (AC) longfin**


Stock Status

Year of Most Recent Assessment	2017
Assessment Runs Presented	Standardised CPUE on positive catch
Reference Points	<p>For ESA, Interim Target is 40% B_0</p> <p>For commercially fished area, Target is B_{MSY} proxy based on CPUE; not determined</p> <p>Default Soft Limit: 20% B_0</p> <p>Default Hard Limit: 10% B_0</p> <p>For ESA, Overfishing threshold is F_{MSY}</p> <p>For commercially fished area, Overfishing threshold is F_{MSY} proxy based on relative exploitation rate; not determined</p>
Status in relation to Target	<p>For total ESA: Likely (> 60%) to be at or above</p> <p>For fished area: Unknown</p>
Status in relation to Limits	<p>For ESA, Soft Limit: Very Unlikely (< 10%) to be below</p> <p>For ESA, Hard Limit: Very Unlikely (< 10%) to be below</p>
Status in relation to Overfishing	<p>For ESA: Unlikely (< 40%) to be overfishing</p> <p>For fished area: Unknown</p>

Historical Stock Status Trajectory and Current Status

Standardised CPUE for longfin eels in Hauraki (AC) from 1990–91 to 2014–15 (from Beentjes & McKenzie 2017). Also shown is the total estimated core fisher longfin catch in AC from ECERs. Vertical dashed line indicates when the 4 kg maximum size was introduced in 2007–08 after which longfin eels 4 kg and over are not recorded on

ECERs. Error bars are 95% confidence intervals. Before 2002, 16% of the catch was recorded as EEU (unidentified) and these catches are omitted.



Annual relative exploitation rate for longfin eels in the Hauraki (AC). Because some catch of longfin was reported as EEU (unidentified) and has not been allocated to species, the exploitation rate is likely to have been higher than shown before 2002.

Fishery and Stock Trends

Recent Trend in Biomass or Proxy	Steep decline in CPUE to 2000–01, and then without trend/stable to 2014–15
Recent Trend in Fishing intensity or Proxy	The relative exploitation rate has declined steeply since 2012 and in 2015 was well below the average for the series.
Other Abundance Indices	-
Trends in Other Relevant Indicators or Variables	Catches of longfin elvers at primary monitoring sites have fluctuated without trend since the series of reliable data begins in 1995–96, suggesting no overall trend in recruitment.

Projections and Prognosis

Stock Projections or Prognosis	Unlikely (< 40%) to decline in the medium term under current catch levels
Probability of Current Catch or TACC causing Biomass to remain below or to decline below Limits	For ESA, Soft Limit: Very Unlikely (< 10%) if catch remains at current levels For ESA, Hard Limit: Very Unlikely (< 10%) if catch remains at current levels
Probability of Current Catch or TACC causing Overfishing to continue or to commence	For ESA, Unlikely (< 40%) if catch remains at current levels

Assessment Methodology and Evaluation

Assessment Type	Level 2 – Partial Quantitative Stock Assessment	
Assessment Method	Standardised CPUE based on positive catches from commercial fyke net	
Assessment Dates	Latest assessment: 2017	Next assessment: 2019
Overall assessment quality rank	1 – High Quality	
Main data inputs (rank)	- Catch and effort data	1 – High Quality
Data not used (rank)	N/A	
Changes to Model Structure and Assumptions	-	
Major Sources of Uncertainty	- Standardised CPUE only provides an index of abundance for eels in areas fished by commercial fishers. Other potential issues with the CPUE indices include: <ul style="list-style-type: none"> • Low numbers of fishers • Uncertainty in target species after 2000 • Exclusion of zero catches 	

	<ul style="list-style-type: none"> • Changes in MLS and retention in early parts of the series and increased escape tube size from 25 mm to 31 mm in 2012–13 • Failure of some fishers to record on ECE returns all legal sized eels caught, not just those retained • Unrecorded release of > 4kg eels since 2007–08
--	---

Qualifying Comments

Because the commercial eel fishery has had a long history (beginning in the late 1960s), and indices of abundance are only available from the early 1990s, it is difficult to infer stock status from recent abundance trends, and these should therefore be interpreted with caution. Other sources of mortality, such as habitat alteration (historical and current) have also reduced abundance prior to the CPUE series.

The proportion of current longfin habitat in Hauraki (Statistical Area AC) fished commercially during the period 2009–10 and 2013–14 is estimated at 50% (Table 15). The proportion of virgin habitat impacted by hydro dams, commercial fishing and other anthropogenic activity was estimated to be 55%.

Fishery Interactions


Bycatch of other species in the commercial Hauraki eel fishery includes mainly catfish, with lesser quantities of Koi carp, goldfish, koura, grey mullet and yellowbelly flounder. Most bycatch species are usually returned alive.

• Waikato (AD) shortfin


Stock Status

Year of Most Recent Assessment	2017
Assessment Runs Presented	Standardised CPUE on positive catch
Reference Points	Target: B_{MSY} proxy based on CPUE; not determined Default Soft Limit: 20% B_0 Default Hard Limit: 10% B_0 Overfishing threshold: F_{MSY} proxy based on relative exploitation rate; not determined
Status in relation to Target	Unknown
Status in relation to Limits	Soft Limit: Unknown Hard Limit: Unknown
Status in relation to Overfishing	Unknown

Historical Stock Status Trajectory and Current Status



Standardised CPUE for shortfin eels in Waikato (AD) from 1990–91 to 2014–15 (from Beentjes & McKenzie 2017). Also shown is the total estimated core fisher shortfin catch in AD from ECERs. Error bars are 95% confidence intervals. Before 2002, 71% of the catch was recorded as EEU (unidentified) and these catches are omitted.



Annual relative exploitation rate for shortfin eels in the Waikato (AD). Because considerable catch of shortfin was reported as EEU (unidentified) and has not been allocated to species, the exploitation rate is likely to have been much higher than shown before 2002.

Fishery and Stock Trends

Recent Trend in Biomass or Proxy	No long-term trend in CPUE until 2003, after which it increased
Recent Trend in Fishing intensity or Proxy	The relative exploitation rate has declined since 2009 and in 2015 was at the series mean.
Other Abundance Indices	-
Trends in Other Relevant Indicators or Variables	Catches of shortfin elvers at primary monitoring sites have fluctuated without trend since the series of reliable data begins in 1995–96, suggesting no overall trend in recruitment.

Projections and Prognosis

Stock Projections or Prognosis	Unlikely (< 40%) to decline in the medium term under current catch levels
Probability of Current Catch or TACC causing Biomass to remain below or to decline below Limits	Unknown
Probability of Current Catch or TACC causing Overfishing to continue or to commence	Unknown

Assessment Methodology and Evaluation

Assessment Type	Level 2 – Partial Quantitative Stock Assessment	
Assessment Method	Standardised CPUE based on positive catches from commercial fyke net	
Assessment Dates	Latest assessment: 2017	Next assessment: 2019
Overall assessment quality rank	1 – High Quality	
Main data inputs (rank)	- Catch and effort data	1 – High Quality
Data not used (rank)	N/A	
Changes to Model Structure and Assumptions	-	
Major Sources of Uncertainty	- Standardised CPUE only provides an index of abundance for eels in areas fished by commercial fishers. Other potential issues with the CPUE indices include: <ul style="list-style-type: none"> • Low numbers of fishers • Uncertainty in target species after 2000 • Exclusion of zero catches 	

	<ul style="list-style-type: none"> Changes in MLS and retention in early parts of the series and increased escape tube size from 25 mm to 31 mm in 2012–13 Failure of some fishers to record on ECE returns all legal sized eels caught, not just those retained
--	--

Qualifying Comments

Because the commercial eel fishery has had a long history (beginning in the late 1960s), and indices of abundance are only available from the early 1990s, it is difficult to infer stock status from recent abundance trends, and these should therefore be interpreted with caution. Other sources of mortality, such as habitat alteration (historical and current) have also reduced abundance prior to the CPUE series.

Fishery Interactions


Bycatch of other species in the commercial Waikato eel fishery includes large quantities of catfish and koi carp, as well as goldfish, rudd, koura, brown trout, perch, and kokopu. Most bycatch species are usually returned alive.

• Waikato (AD) longfin


Stock Status

Year of Most Recent Assessment	2017
Assessment Runs Presented	Standardised CPUE on positive catch
Reference Points	<p>For ESA, Interim Target is 40% B_0</p> <p>For commercially fished area, Target is B_{MSY} proxy based on CPUE; not determined</p> <p>Default Soft Limit: 20% B_0</p> <p>Default Hard Limit: 10% B_0</p> <p>For ESA, Overfishing threshold is F_{MSY}</p> <p>For commercially fished area, Overfishing threshold is F_{MSY} proxy based on relative exploitation rate; not determined</p>
Status in relation to Target	<p>For total ESA: Likely (> 60%) to be at or above</p> <p>For fished area: Unknown</p>
Status in relation to Limits	<p>For ESA, Soft Limit: Very Unlikely (< 10%) to be below</p> <p>For ESA, Hard Limit: Very Unlikely (< 10%) to be below</p>
Status in relation to Overfishing	<p>For ESA: Unlikely (< 40%) to be overfishing</p> <p>For fished area: Unknown</p>

Historical Stock Status Trajectory and Current Status



Standardised CPUE for longfin eels in Waikato (AD) from 1990–91 to 2014–15 (from Beentjes & McKenzie 2017). Also shown is the total estimated core fisher longfin catch in AD from ECERs. Vertical dashed line indicates when the 4 kg maximum size was introduced in 2007–08 after which longfin eels 4 kg and over are not recorded on ECERs. Error bars are 95% confidence intervals. Before 2002, 71% of the catch was recorded as EEU (unidentified) and these catches are omitted.



Annual relative exploitation rate for longfin eels in the Waikato (AD). Because considerable catch of longfin was reported as EEU (unidentified) and has not been allocated to species, the exploitation rate is likely to have been much higher than shown before 2002.

Fishery and Stock Trends

Recent Trend in Biomass or Proxy	A moderate decline in CPUE to 1998, and then a gradual increase to around the level of the former peak by 2014–15
Recent Trend in Fishing intensity or Proxy	The relative exploitation rate has declined steeply since 2002 and in 2015 was well below the series mean.
Other Abundance Indices	-
Trends in Other Relevant Indicators or Variables	Catches of longfin elvers at primary monitoring sites have fluctuated without trend since the series of reliable data begins in 1995–96, suggesting no overall trend in recruitment.

Projections and Prognosis

Stock Projections or Prognosis	Unlikely (< 40%) to decline in the medium term under current catch levels
Probability of Current Catch or TACC causing Biomass to remain below or to decline below Limits	For ESA, Soft Limit: Very Unlikely (< 10%) if catch remains at current levels For ESA, Hard Limit: Very Unlikely (< 10%) if catch remains at current levels
Probability of Current Catch or TACC causing Overfishing to continue or to commence	For ESA, Unlikely (< 40%) if catch remains at current levels

Assessment Methodology and Evaluation

Assessment Type	Level 2 – Partial Quantitative Stock Assessment	
Assessment Method	Standardised CPUE based on positive catches from commercial fyke net	
Assessment Dates	Latest assessment: 2017	Next assessment: 2019
Overall assessment quality rank	1 – High Quality	
Main data inputs (rank)	- Catch and effort data	1 – High Quality
Data not used (rank)	N/A	
Changes to Model Structure and Assumptions	-	
Major Sources of Uncertainty	- Standardised CPUE only provides an index of abundance for eels in areas fished by commercial fishers. Other potential issues with the CPUE indices include: <ul style="list-style-type: none"> • Low numbers of fishers • Uncertainty in target species after 2000 • Exclusion of zero catches 	

	<ul style="list-style-type: none"> • Changes in MLS and retention in early parts of the series and increased escape tube size from 25 mm to 31 mm in 2012–13 • Failure of some fishers to record on ECE returns all legal sized eels caught, not just those retained • Unrecorded release of > 4kg eels since 2007–08
--	---

Qualifying Comments

Because the commercial eel fishery has had a long history (beginning in the late 1960s), and indices of abundance are only available from the early 1990s, it is difficult to infer stock status from recent abundance trends, and these should therefore be interpreted with caution. Other sources of mortality, such as habitat alteration (historical and current) have also reduced abundance prior to the CPUE series.

The proportion of current longfin habitat in Waikato (Statistical Area AD) fished commercially during the period 2009–10 and 2013–14 is estimated at 43% (Table 15). The proportion of virgin habitat impacted by hydro dams, commercial fishing and other anthropogenic activity was estimated to be 56%.

Fishery Interactions


Bycatch of other species in the commercial Waikato eel fishery includes large quantities of catfish and koi carp, as well as goldfish, rudd, koura, brown trout, perch, and kokopu. Most bycatch species are usually returned alive.

• Bay of Plenty (AE) shortfin


Stock Status

Year of Most Recent Assessment	2017
Assessment Runs Presented	Standardised CPUE on positive catch
Reference Points	Target: B_{MSY} proxy based on CPUE; not determined Default Soft Limit: 20% B_0 Default Hard Limit: 10% B_0 Overfishing threshold: F_{MSY} proxy based on relative exploitation rate; not determined
Status in relation to Target	Unknown
Status in relation to Limits	Soft Limit: Unknown Hard Limit: Unknown
Status in relation to Overfishing	Unknown

Historical Stock Status Trajectory and Current Status



Standardised CPUE for shortfin eels in Bay of Plenty (AE) from 1990–91 to 2014–15 (from Beentjes & McKenzie 2017). Also shown is the total estimated core fisher shortfin catch in AE from ECERs. Error bars are 95% confidence intervals. Before 2000, 13% of the catch was recorded as EEU (unidentified) and these catches are omitted.



Annual relative exploitation rate for shortfin eels in the Bay of Plenty (AE). Because some catch of shortfin was reported as EEU (unidentified) and has not been allocated to species, the exploitation rate is likely to have been higher than shown before 2000.

Fishery and Stock Trends

Recent Trend in Biomass or Proxy	No trend in CPUE until 2002, after which it increases steeply to a peak in 2012
Recent Trend in Fishing intensity or Proxy	Relative exploitation rate has declined since 2002, and in 2015 was well below the series mean.
Other Abundance Indices	-
Trends in Other Relevant Indicators or Variables	Catches of shortfin elvers at primary monitoring sites have fluctuated without trend since the series of reliable data begins in 1995–96, suggesting no overall trend in recruitment.

Projections and Prognosis

Stock Projections or Prognosis	Unlikely (< 40%) to decline in the medium term under current catch levels
Probability of Current Catch or TACC causing Biomass to remain below or to decline below Limits	Unknown
Probability of Current Catch or TACC causing Overfishing to continue or to commence	Unknown

Assessment Methodology and Evaluation

Assessment Type	Level 2 – Partial Quantitative Stock Assessment	
Assessment Method	Standardised CPUE based on positive catches from commercial fyke net	
Assessment Dates	Latest assessment: 2017	Next assessment: 2019
Overall assessment quality rank	1 – High Quality	
Main data inputs (rank)	- Catch and effort data	1 – High Quality
Data not used (rank)	N/A	
Changes to Model Structure and Assumptions	-	
Major Sources of Uncertainty	- Standardised CPUE only provides an index of abundance for eels in areas fished by commercial fishers. Other potential issues with the CPUE indices include: <ul style="list-style-type: none"> • Low numbers of fishers • Uncertainty in target species after 2000 • Exclusion of zero catches 	

	<ul style="list-style-type: none"> Changes in MLS and retention in early parts of the series and increased escape tube size from 25 mm to 31 mm in 2012–13 Failure of some fishers to record on ECE returns all legal sized eels caught, not just those retained
--	--

Qualifying Comments

Because the commercial eel fishery has had a long history (beginning in the late 1960s), and indices of abundance are only available from the early 1990s, it is difficult to infer stock status from recent abundance trends, and these should therefore be interpreted with caution. Other sources of mortality, such as habitat alteration (historical and current) have also reduced abundance prior to the CPUE series.

Fishery Interactions


Bycatch of other species in the commercial Bay of Plenty eel fishery includes very small quantities of goldfish and bullies. Most bycatch species are usually returned alive.

• Bay of Plenty (AE) longfin


Stock Status

Year of Most Recent Assessment	2017
Assessment Runs Presented	Standardised CPUE on positive catch
Reference Points	<p>For ESA, Interim Target is 40% B_0</p> <p>For commercially fished area, Target is B_{MSY} proxy based on CPUE; not determined</p> <p>Default Soft Limit: 20% B_0</p> <p>Default Hard Limit: 10% B_0</p> <p>For ESA, Overfishing threshold is F_{MSY}</p> <p>For commercially fished area, Overfishing threshold is F_{MSY} proxy based on relative exploitation rate; not determined</p>
Status in relation to Target	<p>For total ESA: Likely (> 60%) to be at or above</p> <p>For fished area: Unknown</p>
Status in relation to Limits	<p>For ESA, Soft Limit: Very Unlikely (< 10%) to be below</p> <p>For ESA, Hard Limit: Very Unlikely (< 10%) to be below</p>
Status in relation to Overfishing	<p>For ESA: Unlikely (< 40%) to be overfishing</p> <p>For fished area: Unknown</p>

Historical Stock Status Trajectory and Current Status



Standardised CPUE for longfin eels in Bay of Plenty (AE) from 1990–91 to 2014–15 (from Beentjes & McKenzie 2017). Also shown is the total estimated core fisher longfin catch in AE from ECERs. Vertical dashed line indicates when the 4 kg maximum size was introduced in 2007–08 after which longfin eels 4 kg and over are not recorded on ECERs. Error bars are 95% confidence intervals. Before 2000, 13% of the catch was recorded as EEU (unidentified) and these catches are omitted.



Annual relative exploitation rate for longfin eels in the Bay of Plenty (AE). Because some catch of longfin was reported as EEU (unidentified) and has not been allocated to species, the exploitation rate is likely to have been higher than shown before 2000.

Fishery and Stock Trends

Recent Trend in Biomass or Proxy	A steep decline in CPUE to 2000, and then a gradual increase to a peak in 2012–13
Recent Trend in Fishing intensity or Proxy	The relative exploitation rate has declined since 2005, and since 2007 has been below the series mean.
Other Abundance Indices	-
Trends in Other Relevant Indicators or Variables	Catches of longfin elvers at primary monitoring sites have fluctuated without trend since the series of reliable data begins in 1995–96, suggesting no overall trend in recruitment.

Projections and Prognosis

Stock Projections or Prognosis	Unlikely (< 40%) to decline in the medium term under current catch levels
Probability of Current Catch or TACC causing Biomass to remain below or to decline below Limits	For ESA, Soft Limit: Very Unlikely (< 10%) if catch remains at current levels For ESA, Hard Limit: Very Unlikely (< 10%) if catch remains at current levels
Probability of Current Catch or TACC causing Overfishing to continue or to commence	For ESA, Unlikely (< 40%) if catch remains at current levels

Assessment Methodology and Evaluation

Assessment Type	Level 2 – Partial Quantitative Stock Assessment	
Assessment Method	Standardised CPUE based on positive catches from commercial fyke net	
Assessment Dates	Latest assessment: 2017	Next assessment: 2019
Overall assessment quality rank	1 – High Quality	
Main data inputs (rank)	- Catch and effort data	1 – High Quality
Data not used (rank)	N/A	
Changes to Model Structure and Assumptions	-	
Major Sources of Uncertainty	- Standardised CPUE only provides an index of abundance for eels in areas fished by commercial fishers. Other potential issues with the CPUE indices include: <ul style="list-style-type: none"> • Low numbers of fishers • Uncertainty in target species after 2000 • Exclusion of zero catches 	

	<ul style="list-style-type: none"> • Changes in MLS and retention in early parts of the series and increased escape tube size from 25 mm to 31 mm in 2012–13 • Failure of some fishers to record on ECE returns all legal sized eels caught, not just those retained • Unrecorded release of > 4kg eels since 2007–08
--	---

Qualifying Comments

Because the commercial eel fishery has had a long history (beginning in the late 1960s), and indices of abundance are only available from the early 1990s, it is difficult to infer stock status from recent abundance trends, and these should therefore be interpreted with caution. Other sources of mortality, such as habitat alteration (historical and current) have also reduced abundance prior to the CPUE series.

The proportion of current longfin habitat in Bay of Plenty (Statistical Area AE) fished commercially during the period 2009–10 and 2013–14 is estimated at 17% (Table 15). The proportion of virgin habitat impacted by hydro dams, commercial fishing and other anthropogenic activity was estimated to be 24%.

Fishery Interactions


Bycatch of other species in the commercial Bay of Plenty eel fishery includes very small quantities of goldfish and bullies. Most bycatch species are usually returned alive.

QMA SFE 22 and LFE 22 (includes ESAs AG, AK, AL and AM)


• Hawkes Bay (AG) shortfin

Stock Status	
Year of Most Recent Assessment	2017
Assessment Runs Presented	Standardised CPUE on positive catch
Reference Points	Target: B_{MSY} proxy based on CPUE; not determined Default Soft Limit: 20% B_0 Default Hard Limit: 10% B_0 Overfishing threshold: F_{MSY} proxy based on relative exploitation rate; not determined
Status in relation to Target	Unknown
Status in relation to Limits	Soft Limit: Unknown Hard Limit: Unknown
Status in relation to Overfishing	Unknown

Historical Stock Status Trajectory and Current Status



Standardised CPUE for shortfin eels in Hawkes Bay (AG) from 1990–91 to 2014–15 (from Beentjes & McKenzie 2017). Also shown is the total estimated core fisher shortfin catch in AG from ECERs. Error bars are 95% confidence intervals. Before 2001, 5% of the catch was recorded as EEU (unidentified) and these catches are omitted.



Annual relative exploitation rate for shortfin eels in the Hawkes Bay (AG). Because some catch of shortfin was reported as EEU (unidentified) and has not been allocated to species, the exploitation rate is likely to have been higher than shown before 2001.

Fishery and Stock Trends

Recent Trend in Biomass or Proxy	CPUE declined until 2002, followed by a steep increase.
Recent Trend in Fishing intensity or Proxy	The relative exploitation rate has declined since 2007, and from 2009 has been below the series mean.
Other Abundance Indices	-
Trends in Other Relevant Indicators or Variables	Catches of shortfin elvers at primary monitoring sites have fluctuated without trend since the series of reliable data begins in 1995–96, suggesting no overall trend in recruitment.

Projections and Prognosis

Stock Projections or Prognosis	Unlikely (< 40%) to decline in the medium term under current catch levels
Probability of Current Catch or TACC causing Biomass to remain below or to decline below Limits	Unknown
Probability of Current Catch or TACC causing Overfishing to continue or to commence	Unknown

Assessment Methodology and Evaluation

Assessment Type	Level 2 – Partial Quantitative Stock Assessment	
Assessment Method	Standardised CPUE based on positive catches from commercial fyke net	
Assessment Dates	Latest assessment: 2017	Next assessment: 2019
Overall assessment quality rank	1 – High Quality	
Main data inputs (rank)	- Catch and effort data	1 – High Quality
Data not used (rank)	N/A	
Changes to Model Structure and Assumptions	-	
Major Sources of Uncertainty	<p>- Standardised CPUE only provides an index of abundance for eels in areas fished by commercial fishers. Other potential issues with the CPUE indices include:</p> <ul style="list-style-type: none"> • Low numbers of fishers • Uncertainty in target species after 2000 • Exclusion of zero catches • Changes in MLS and retention in early parts of the series and increased escape tube size from 25 mm to 31 mm in 2012–13 	

- Failure of some fishers to record on ECE returns all legal sized eels caught, not just those retained

Qualifying Comments

Because the commercial eel fishery has had a long history (beginning in the late 1960s), and indices of abundance are only available from the early 1990s, it is difficult to infer stock status from recent abundance trends, and these should therefore be interpreted with caution. Other sources of mortality, such as habitat alteration (historical and current) have also reduced abundance prior to the CPUE series.

Fishery Interactions


Bycatch of other species in the commercial Hawkes Bay eel fishery includes mostly goldfish and small quantities of brown trout. Most bycatch species are usually returned alive.

• Hawkes Bay (AG) longfin


Stock Status

Year of Most Recent Assessment	2017
Assessment Runs Presented	Standardised CPUE on positive catch
Reference Points	For ESA, Interim Target is 40% B_0 For commercially fished area, Target is B_{MSY} proxy based on CPUE; not determined Default Soft Limit: 20% B_0 Default Hard Limit: 10% B_0 For ESA, Overfishing threshold is F_{MSY} For commercially fished area, Overfishing threshold is F_{MSY} proxy based on relative exploitation rate; not determined
Status in relation to Target	For total ESA: Likely (> 60%) to be at or above For fished area: Unknown
Status in relation to Limits	For ESA, Soft Limit: Very Unlikely (< 10%) to be below For ESA, Hard Limit: Very Unlikely (< 10%) to be below
Status in relation to Overfishing	For ESA: Unlikely (< 40%) to be overfishing For fished area: Unknown

Historical Stock Status Trajectory and Current Status



Standardised CPUE for longfin eels in Hawkes Bay (AG) from 1990–91 to 2014–15 (from Beentjes & McKenzie 2017). Also shown is the total estimated core fisher longfin catch in AG from ECERs. Vertical dashed line indicates when the 4 kg maximum size was introduced in 2007–08 after which longfin eels 4 kg and over are not recorded on ECERs. Error bars are 95% confidence intervals. Before 2001, 5% of the catch was recorded as EEU (unidentified) and these catches are omitted.



Annual relative exploitation rate for longfin eels in the Hawke's Bay (AG). Because some catch of longfin was reported as EEU (unidentified) and has not been allocated to species, the exploitation rate is likely to have been higher than shown before 2001.

Fishery and Stock Trends

Recent Trend in Biomass or Proxy	CPUE declined until 1997, was stable until 2008 and then increased
Recent Trend in Fishing intensity or Proxy	The relative exploitation rate has declined steeply since 2007, and in 2015 was well below the series mean.
Other Abundance Indices	-
Trends in Other Relevant Indicators or Variables	Catches of longfin elvers at primary monitoring sites have fluctuated without trend since the series of reliable data begins in 1995–96, suggesting no overall trend in recruitment.

Projections and Prognosis

Stock Projections or Prognosis	Unlikely (< 40%) to decline in the medium term under current catch levels
Probability of Current Catch or TACC causing Biomass to remain below or to decline below Limits	For ESA, Soft Limit: Very Unlikely (< 10%) if catch remains at current levels For ESA, Hard Limit: Very Unlikely (< 10%) if catch remains at current levels
Probability of Current Catch or TACC causing Overfishing to continue or to commence	For ESA, Unlikely (< 40%) if catch remains at current levels

Assessment Methodology and Evaluation

Assessment Type	Level 2 – Partial Quantitative Stock Assessment	
Assessment Method	Standardised CPUE based on positive catches from commercial fyke net	
Assessment Dates	Latest assessment: 2017	Next assessment: 2019
Overall assessment quality rank	1 – High Quality	
Main data inputs (rank)	- Catch and effort data	1 – High Quality
Data not used (rank)	N/A	
Changes to Model Structure and Assumptions	-	
Major Sources of Uncertainty	- Standardised CPUE only provides an index of abundance for eels in areas fished by commercial fishers. Other potential issues with the CPUE indices include: <ul style="list-style-type: none"> • Low numbers of fishers • Uncertainty in target species after 2000 • Exclusion of zero catches 	

	<ul style="list-style-type: none"> • Changes in MLS and retention in early parts of the series and increased escape tube size from 25 mm to 31 mm in 2012–13 • Failure of some fishers to record on ECE returns all legal sized eels caught, not just those retained • Unrecorded release of > 4kg eels since 2007–08
--	---

Qualifying Comments

Because the commercial eel fishery has had a long history (beginning in the late 1960s), and indices of abundance are only available from the early 1990s, it is difficult to infer stock status from recent abundance trends, and these should therefore be interpreted with caution. Other sources of mortality, such as habitat alteration (historical and current) have also reduced abundance prior to the CPUE series.

The proportion of current longfin habitat in Hawkes Bay (Statistical Area AG) fished commercially during the period 2009–10 and 2013–14 is estimated at 17% (Table 15). The proportion of virgin habitat impacted by hydro dams, commercial fishing and other anthropogenic activity was estimated to be 25%.

Fishery Interactions


Bycatch of other species in the commercial Hawkes Bay eel fishery includes mostly goldfish and small quantities of brown trout. Most bycatch species are usually returned alive.

• Manawatu (AK) shortfin


Stock Status

Year of Most Recent Assessment	2017
Assessment Runs Presented	Standardised CPUE on positive catch
Reference Points	Target: B_{MSY} proxy based on CPUE; not determined Default Soft Limit: 20% B_0 Default Hard Limit: 10% B_0 Overfishing threshold: F_{MSY} proxy based on relative exploitation rate; not determined
Status in relation to Target	Unknown
Status in relation to Limits	Soft Limit: Unknown Hard Limit: Unknown
Status in relation to Overfishing	Unknown

Historical Stock Status Trajectory and Current Status



Standardised CPUE for shortfin eels in Manawatu (AK) from 1990–91 to 2014–15 (from Beentjes & McKenzie 2017). Also shown is the total estimated core fisher shortfin catch in AK from ECERs. Error bars are 95% confidence intervals. Before 2001, 56% of the catch was recorded as EEU (unidentified) and these catches are omitted.



Annual relative exploitation rate for shortfin eels in the Manawatu (AK). Because some catch of shortfin was reported as EEU (unidentified) and has not been allocated to species, the exploitation rate is likely to have been higher than shown before 2001.

Fishery and Stock Trends

Recent Trend in Biomass or Proxy	CPUE dropped markedly from 1992 to 1994, was stable until an increase in 2004, and has fluctuated without trend since then
Recent Trend in Fishing intensity or Proxy	The relative exploitation rate has declined since 2013, and in 2015 was below the series mean.
Other Abundance Indices	-
Trends in Other Relevant Indicators or Variables	Catches of shortfin elvers at primary monitoring sites have fluctuated without trend since the series of reliable data begins in 1995–96, suggesting no overall trend in recruitment.

Projections and Prognosis

Stock Projections or Prognosis	Unlikely (< 40%) to decline in the medium term under current catch levels
Probability of Current Catch or TACC causing Biomass to remain below or to decline below Limits	Unknown
Probability of Current Catch or TACC causing Overfishing to continue or to commence	Unknown

Assessment Methodology and Evaluation

Assessment Type	Level 2 – Partial Quantitative Stock Assessment	
Assessment Method	Standardised CPUE based on positive catches from commercial fyke net	
Assessment Dates	Latest assessment: 2017	Next assessment: 2019
Overall assessment quality rank	1 – High Quality	
Main data inputs (rank)	- Catch and effort data	1 – High Quality
Data not used (rank)	N/A	
Changes to Model Structure and Assumptions	-	
Major Sources of Uncertainty	- Standardised CPUE only provides an index of abundance for eels in areas fished by commercial fishers. Other potential issues with the CPUE indices include: <ul style="list-style-type: none"> • Low numbers of fishers • Uncertainty in target species after 2000 • Exclusion of zero catches 	

	<ul style="list-style-type: none"> Changes in MLS and retention in early parts of the series and increased escape tube size from 25 mm to 31 mm in 2012–13 Failure of some fishers to record on ECE returns all legal sized eels caught, not just those retained
--	--

Qualifying Comments

Because the commercial eel fishery has had a long history (beginning in the late 1960s), and indices of abundance are only available from the early 1990s, it is difficult to infer stock status from recent abundance trends, and these should therefore be interpreted with caution. Other sources of mortality, such as habitat alteration (historical and current) have also reduced abundance prior to the CPUE series.

Fishery Interactions


Bycatch in the commercial Manawatu eel fishery include small quantities of koi carp, black flounder, yellowbelly flounder, and perch. Most bycatch species are usually returned alive.

• Manawatu (AK) longfin


Stock Status

Year of Most Recent Assessment	2017
Assessment Runs Presented	Standardised CPUE on positive catch
Reference Points	<p>For ESA, Interim Target is 40% B_0</p> <p>For commercially fished area, Target is B_{MSY} proxy based on CPUE; not determined</p> <p>Default Soft Limit: 20% B_0</p> <p>Default Hard Limit: 10% B_0</p> <p>For ESA, Overfishing threshold is F_{MSY}</p> <p>For commercially fished area, Overfishing threshold is F_{MSY} proxy based on relative exploitation rate; not determined</p>
Status in relation to Target	<p>For total ESA: Likely (> 60%) to be at or above</p> <p>For fished area: Unknown</p>
Status in relation to Limits	<p>For ESA, Soft Limit: Very Unlikely (< 10%) to be below</p> <p>For ESA, Hard Limit: Very Unlikely (< 10%) to be below</p>
Status in relation to Overfishing	<p>For ESA: Unlikely (< 40%) to be overfishing</p> <p>For fished area: Unknown</p>

Historical Stock Status Trajectory and Current Status



Standardised CPUE for longfin eels in Manawatu (AK) from 1990–91 to 2014–15 (from Beentjes & McKenzie 2017). Also shown is the total estimated core fisher longfin catch in AK from ECERs. Vertical dashed line indicates when the 4 kg maximum size was introduced in 2007–08 after which longfin eels 4 kg and over are not recorded on ECERs. Error bars are 95% confidence intervals. Before 2001, 56% of the catch was recorded as EEU (unidentified) and these catches are omitted.



Annual relative exploitation rate for longfin eels in the Manawatu (AK). Because some catch of longfin was reported as EEU (unidentified) and has not been allocated to species, the exploitation rate is likely to have been higher than shown before 2001.

Fishery and Stock Trends

Recent Trend in Biomass or Proxy	CPUE declined steeply until 2003, increased in 2004 and has fluctuated without trend since then.
Recent Trend in Fishing intensity or Proxy	The relative exploitation rate has declined since 2013, and in 2015 was just below the series mean.
Other Abundance Indices	-
Trends in Other Relevant Indicators or Variables	Catches of longfin elvers at primary monitoring sites have fluctuated without trend since the series of reliable data begins in 1995–96, suggesting no overall trend in recruitment.

Projections and Prognosis

Stock Projections or Prognosis	Unlikely (< 40%) to decline in the medium term under current catch levels
Probability of Current Catch or TACC causing Biomass to remain below or to decline below Limits	For ESA, Soft Limit: Very Unlikely (< 10%) if catch remains at current levels For ESA, Hard Limit: Very Unlikely (< 10%) if catch remains at current levels
Probability of Current Catch or TACC causing Overfishing to continue or to commence	For ESA, Unlikely (< 40%) if catch remains at current levels

Assessment Methodology and Evaluation

Assessment Type	Level 2 – Partial Quantitative Stock Assessment	
Assessment Method	Standardised CPUE based on positive catches from commercial fyke net	
Assessment Dates	Latest assessment: 2017	Next assessment: 2019
Overall assessment quality rank	1 – High Quality	
Main data inputs (rank)	- Catch and effort data	1 – High Quality
Data not used (rank)	N/A	
Changes to Model Structure and Assumptions	-	
Major Sources of Uncertainty	- Standardised CPUE only provides an index of abundance for eels in areas fished by commercial fishers. Other potential issues with the CPUE indices include: <ul style="list-style-type: none"> • Low numbers of fishers • Uncertainty in target species after 2000 • Exclusion of zero catches 	

	<ul style="list-style-type: none"> • Changes in MLS and retention in early parts of the series and increased escape tube size from 25 mm to 31 mm in 2012–13 • Failure of some fishers to record on ECE returns all legal sized eels caught, not just those retained • Unrecorded release of > 4kg eels since 2007–08
--	---

Qualifying Comments

Because the commercial eel fishery has had a long history (beginning in the late 1960s), and indices of abundance are only available from the early 1990s, it is difficult to infer stock status from recent abundance trends, and these should therefore be interpreted with caution. Other sources of mortality, such as habitat alteration (historical and current) have also reduced abundance prior to the CPUE series.

The proportion of current longfin habitat in Manawatu (Statistical Area AK) fished commercially during the period 2009–10 and 2013–14 is estimated at 36% (Table 15). The proportion of virgin habitat impacted by hydro dams, commercial fishing and other anthropogenic activity was estimated to be 41%.

Fishery Interactions


Bycatch in the commercial Manawatu eel fishery include small quantities of koi carp, black flounder, yellowbelly flounder, and perch. Most bycatch species are usually returned alive.

• Wairarapa (AL) shortfin


Stock Status

Year of Most Recent Assessment	2017
Assessment Runs Presented	Standardised CPUE on positive catch
Reference Points	Target: B_{MSY} proxy based on CPUE; not determined Default Soft Limit: 20% B_0 Default Hard Limit: 10% B_0 Overfishing threshold: F_{MSY} proxy based on relative exploitation rate; not determined
Status in relation to Target	Unknown
Status in relation to Limits	Soft Limit: Unknown Hard Limit: Unknown
Status in relation to Overfishing	Unknown

Historical Stock Status Trajectory and Current Status



Standardised CPUE for shortfin eels in Wairarapa (AL) from 1990–91 to 2014–15 (from Beentjes & McKenzie 2017). Also shown is the total estimated core fisher shortfin catch in AL from ECERs. Error bars are 95% confidence intervals. Before 1999, 33% of the catch was recorded as EEU (unidentified) and these catches are omitted.



Annual relative exploitation rate for shortfin eels in the Wairarapa (AL). Because some catch of shortfin was reported as EEU (unidentified) and has not been allocated to species, the exploitation rate is likely to have been higher than shown before 1999.

Fishery and Stock Trends

Recent Trend in Biomass or Proxy	CPUE declined from 1995 to 2003, increased in 2005 and has fluctuated without trend since then.
Recent Trend in Fishing intensity or Proxy	The relative exploitation rate declined steeply after 2003, and has been below the series mean since 2005.
Other Abundance Indices	-
Trends in Other Relevant Indicators or Variables	Catches of shortfin elvers at primary monitoring sites have fluctuated without trend since the series of reliable data begins in 1995–96, suggesting no overall trend in recruitment.

Projections and Prognosis

Stock Projections or Prognosis	Unlikely (< 40%) to decline in the medium term under current catch levels
Probability of Current Catch or TACC causing Biomass to remain below or to decline below Limits	Unknown
Probability of Current Catch or TACC causing Overfishing to continue or to commence	Unknown

Assessment Methodology and Evaluation

Assessment Type	Level 2 – Partial Quantitative Stock Assessment	
Assessment Method	Standardised CPUE based on positive catches from commercial fyke net	
Assessment Dates	Latest assessment: 2017	Next assessment: 2019
Overall assessment quality rank	1 – High Quality	
Main data inputs (rank)	- Catch and effort data	1 – High Quality
Data not used (rank)	N/A	
Changes to Model Structure and Assumptions	-	
Major Sources of Uncertainty	<p>- Standardised CPUE only provides an index of abundance for eels in areas fished by commercial fishers. Other potential issues with the CPUE indices include:</p> <ul style="list-style-type: none"> • Low numbers of fishers • Uncertainty in target species after 2000 • Exclusion of zero catches • Changes in MLS and retention in early parts of the series and increased escape tube size from 25 mm to 31 mm in 2012–13 	

- Failure of some fishers to record on ECE returns all legal sized eels caught, not just those retained

Qualifying Comments

Because the commercial eel fishery has had a long history (beginning in the late 1960s), and indices of abundance are only available from the early 1990s, it is difficult to infer stock status from recent abundance trends, and these should therefore be interpreted with caution. Other sources of mortality, such as habitat alteration (historical and current) have also reduced abundance prior to the CPUE series.

Fishery Interactions


Bycatch in the commercial Wairarapa eel fishery include mostly rudd and perch, with smaller quantities of flatfish and goldfish. Most bycatch species are usually returned alive.

• Wairarapa (AL) longfin


Stock Status

Year of Most Recent Assessment	2017
Assessment Runs Presented	Standardised CPUE on positive catch
Reference Points	For ESA, Interim Target is 40% B_0 For commercially fished area, Target is B_{MSY} proxy based on CPUE; not determined Default Soft Limit: 20% B_0 Default Hard Limit: 10% B_0 For ESA, Overfishing threshold is F_{MSY} For commercially fished area, Overfishing threshold is F_{MSY} proxy based on relative exploitation rate; not determined
Status in relation to Target	For total ESA: Likely (> 60%) to be at or above For fished area: Unknown
Status in relation to Limits	For ESA, Soft Limit: Very Unlikely (< 10%) to be below For ESA, Hard Limit: Very Unlikely (< 10%) to be below
Status in relation to Overfishing	For ESA: Unlikely (< 40%) to be overfishing For fished area: Unknown

Historical Stock Status Trajectory and Current Status



Standardised CPUE for longfin eels in Wairarapa (AL) from 1990–91 to 2014–15 (from Beentjes & McKenzie 2017). Also shown is the total estimated core fisher longfin catch in AL from ECERs. Vertical dashed line indicates when the 4 kg maximum size was introduced in 2007–08 after which longfin eels 4 kg and over are not recorded on ECERs. Error bars are 95% confidence intervals. Before 1999, 33% of the catch was recorded as EEU (unidentified) and these catches are omitted.



Annual relative exploitation rate for longfin eels in the Wairarapa (AL). Because some catch of longfin was reported as EEU (unidentified) and has not been allocated to species, the exploitation rate is likely to have been higher than shown before 1999.

Fishery and Stock Trends

Recent Trend in Biomass or Proxy	CPUE declined until 2003, increased in 2004 and has been fluctuated without trend since then.
Recent Trend in Fishing intensity or Proxy	The relative exploitation rate declined steeply after 2003, and has been below the series mean since 2005.
Other Abundance Indices	-
Trends in Other Relevant Indicators or Variables	Catches of longfin elvers at primary monitoring sites have fluctuated without trend since the series of reliable data begins in 1995–96, suggesting no overall trend in recruitment.

Projections and Prognosis

Stock Projections or Prognosis	Unlikely (< 40%) to decline in the medium term under current catch levels
Probability of Current Catch or TACC causing Biomass to remain below or to decline below Limits	For ESA, Soft Limit: Very Unlikely (< 10%) if catch remains at current levels For ESA, Hard Limit: Very Unlikely (< 10%) if catch remains at current levels
Probability of Current Catch or TACC causing Overfishing to continue or to commence	For ESA, Unlikely (< 40%) if catch remains at current levels

Assessment Methodology and Evaluation

Assessment Type	Level 2 – Partial Quantitative Stock Assessment	
Assessment Method	Standardised CPUE based on positive catches from commercial fyke net	
Assessment Dates	Latest assessment: 2017	Next assessment: 2019
Overall assessment quality rank	1 – High Quality	
Main data inputs (rank)	- Catch and effort data	1 – High Quality
Data not used (rank)	N/A	
Changes to Model Structure and Assumptions	-	
Major Sources of Uncertainty	<p>- Standardised CPUE only provides an index of abundance for eels in areas fished by commercial fishers. Other potential issues with the CPUE indices include:</p> <ul style="list-style-type: none"> • Low numbers of fishers • Uncertainty in target species after 2000 • Exclusion of zero catches • Changes in MLS and retention in early parts of the series and increased escape tube size from 25 mm to 31 mm in 2012–13 	

FRESHWATER EELS (SFE, LFE, ANG)

	<ul style="list-style-type: none"> • Failure of some fishers to record on ECE returns all legal sized eels caught, not just those retained • Unrecorded release of > 4kg eels since 2007–08
--	--

Qualifying Comments

Because the commercial eel fishery has had a long history (beginning in the late 1960s), and indices of abundance are only available from the early 1990s, it is difficult to infer stock status from recent abundance trends, and these should therefore be interpreted with caution. Other sources of mortality, such as habitat alteration (historical and current) have also reduced abundance prior to the CPUE series.

The proportion of current longfin habitat in Wairarapa (Statistical Area AL) fished commercially during the period 2009–10 and 2013–14 is estimated at 4% (Table 15) (Beentjes et al 2016). The proportion of virgin habitat impacted by hydro dams, commercial fishing and other anthropogenic activity was estimated to be 5%.

Fishery Interactions

There has been no recorded bycatch in the commercial Wairarapa eel fishery since 2000–01. Most bycatch species are usually returned alive.


QMA SFE 23 and LFE 23 (includes ESAs AH, AJ)

• Rangitikei-Wanganui (AH) shortfin


Stock Status

Year of Most Recent Assessment	2017
Assessment Runs Presented	Standardised CPUE on positive catch
Reference Points	Target: B_{MSY} proxy based on CPUE; not determined Default Soft Limit: 20% B_0 Default Hard Limit: 10% B_0 Overfishing threshold: F_{MSY} proxy based on relative exploitation rate; not determined
Status in relation to Target	Unknown
Status in relation to Limits	Soft Limit: Unknown Hard Limit: Unknown
Status in relation to Overfishing	Unknown

Historical Stock Status Trajectory and Current Status



Standardised CPUE for shortfin eels in Rangitikei-Wanganui (AH) from 1990–91 to 2014–15 (from Beentjes & McKenzie 2017). Also shown is the total estimated core fisher shortfin catch in AH from ECERs. Error bars are 95% confidence intervals. Before 2001, 7% of the catch was recorded as EEU (unidentified) and these catches are omitted.



Annual relative exploitation rate for shortfin eels in the Rangitikei-Wanganui (AH). Because some catch of shortfin was reported as EEU (unidentified) and has not been allocated to species, the exploitation rate is likely to have been higher than shown before 2001.

Fishery and Stock Trends

Recent Trend in Biomass or Proxy	CPUE declined until 2004, increased steeply to 2012, and then declined to 2015
Recent Trend in Fishing intensity or Proxy	The relative exploitation rate declined steeply after 2003, and has been below the series mean since 2004.
Other Abundance Indices	-
Trends in Other Relevant Indicators or Variables	Catches of shortfin elvers at primary monitoring sites have fluctuated without trend since the series of reliable data begins in 1995–96, suggesting no overall trend in recruitment.

Projections and Prognosis

Stock Projections or Prognosis	Unlikely (< 40%) to decline in the medium term under current catch levels
Probability of Current Catch or TACC causing Biomass to remain below or to decline below Limits	Unknown
Probability of Current Catch or TACC causing Overfishing to continue or to commence	Unknown

Assessment Methodology and Evaluation

Assessment Type	Level 2 – Partial Quantitative Stock Assessment	
Assessment Method	Standardised CPUE based on positive catches from commercial fyke net	
Assessment Dates	Latest assessment: 2017	Next assessment: 2019
Overall assessment quality rank	1 – High Quality	
Main data inputs (rank)	- Catch and effort data	1 – High Quality
Data not used (rank)	N/A	
Changes to Model Structure and Assumptions	-	
Major Sources of Uncertainty	<p>- Standardised CPUE only provides an index of abundance for eels in areas fished by commercial fishers. Other potential issues with the CPUE indices include:</p> <ul style="list-style-type: none"> • Low numbers of fishers • Uncertainty in target species after 2000 • Exclusion of zero catches • Changes in MLS and retention in early parts of the series and increased escape tube size from 25 mm to 31 mm in 2012–13 	

FRESHWATER EELS (SFE, LFE, ANG)

	<ul style="list-style-type: none"> Failure of some fishers to record on ECE returns all legal sized eels caught, not just those retained
--	---

Qualifying Comments

Because the commercial eel fishery has had a long history (beginning in the late 1960s), and indices of abundance are only available from the early 1990s, it is difficult to infer stock status from recent abundance trends, and these should therefore be interpreted with caution. Other sources of mortality, such as habitat alteration (historical and current) have also reduced abundance prior to the CPUE series.

Fishery Interactions


The only recorded bycatch in the commercial Rangitikei-Wanganui eel fishery since 2000–01 has been brown trout. Most bycatch species are usually returned alive.

• Taranaki (AJ) shortfin


Stock Status

Year of Most Recent Assessment	2017
Assessment Runs Presented	Standardised CPUE on positive catch
Reference Points	Target: B_{MSY} proxy based on CPUE; not determined Default Soft Limit: 20% B_0 Default Hard Limit: 10% B_0 Overfishing threshold: F_{MSY} proxy based on relative exploitation rate; not determined
Status in relation to Target	Unknown
Status in relation to Limits	Soft Limit: Unknown Hard Limit: Unknown
Status in relation to Overfishing	Unknown

Historical Stock Status Trajectory and Current Status



Standardised CPUE for shortfin eels in Taranaki (AJ) from 1990–91 to 2014–15 (from Beentjes & McKenzie 2017). Also shown is the total estimated core fisher shortfin catch in AJ from ECERs. Error bars are 95% confidence intervals. Before 2001, 16% of the catch was recorded as EEU (unidentified) and these catches are omitted.



Annual relative exploitation rate for shortfin eels in the Taranaki (AJ). Because some catch of shortfin was reported as EEU (unidentified) and has not been allocated to species, the exploitation rate is likely to have been higher than shown before 2001.

Fishery and Stock Trends

Recent Trend in Biomass or Proxy	CPUE declined to 2003, followed by a gradual increase
Recent Trend in Fishing intensity or Proxy	Relative exploitation rate declined steeply after 2002, and has been below the series mean since 2005.
Other Abundance Indices	-
Trends in Other Relevant Indicators or Variables	Catches of shortfin elvers at primary monitoring sites have fluctuated without trend since the series of reliable data begins in 1995–96, suggesting no overall trend in recruitment.

Projections and Prognosis

Stock Projections or Prognosis	Unlikely (< 40%) to decline in the medium term under current catch levels
Probability of Current Catch or TACC causing Biomass to remain below or to decline below Limits	Unknown
Probability of Current Catch or TACC causing Overfishing to continue or to commence	Unknown

Assessment Methodology and Evaluation

Assessment Type	Level 2 – Partial Quantitative Stock Assessment	
Assessment Method	Standardised CPUE based on positive catches from commercial fyke net	
Assessment Dates	Latest assessment: 2017	Next assessment: 2019
Overall assessment quality rank	1 – High Quality	
Main data inputs (rank)	- Catch and effort data	1 – High Quality
Data not used (rank)	N/A	
Changes to Model Structure and Assumptions	-	
Major Sources of Uncertainty	<p>- Standardised CPUE only provides an index of abundance for eels in areas fished by commercial fishers. Other potential issues with the CPUE indices include:</p> <ul style="list-style-type: none"> • Low numbers of fishers • Uncertainty in target species after 2000 • Exclusion of zero catches • Changes in MLS and retention in early parts of the series and increased escape tube size from 25 mm to 31 mm in 2012–13 	

	<ul style="list-style-type: none"> Failure of some fishers to record on ECE returns all legal sized eels caught, not just those retained
--	---

Qualifying Comments

Because the commercial eel fishery has had a long history (beginning in the late 1960s), and indices of abundance are only available from the early 1990s, it is difficult to infer stock status from recent abundance trends, and these should therefore be interpreted with caution. Other sources of mortality, such as habitat alteration (historical and current) have also reduced abundance prior to the CPUE series.

Fishery Interactions


There has been no recorded bycatch in the commercial Taranaki eel fishery since 2000–01. Most bycatch species are usually returned alive.

• Taranaki (AJ) longfin


Stock Status

Year of Most Recent Assessment	2017
Assessment Runs Presented	Standardised CPUE on positive catch
Reference Points	<p>For ESA, Interim Target is 40% B_0</p> <p>For commercially fished area, Target is B_{MSY} proxy based on CPUE; not determined</p> <p>Default Soft Limit: 20% B_0</p> <p>Default Hard Limit: 10% B_0</p> <p>For ESA, Overfishing threshold is F_{MSY}</p> <p>For commercially fished area, Overfishing threshold is F_{MSY} proxy based on relative exploitation rate; not determined</p>
Status in relation to Target	<p>For total ESA: Likely (> 60%) to be at or above</p> <p>For fished area: Unknown</p>
Status in relation to Limits	<p>For ESA, Soft Limit: Very Unlikely (< 10%) to be below</p> <p>For ESA, Hard Limit: Very Unlikely (< 10%) to be below</p>
Status in relation to Overfishing	<p>For ESA: Unlikely (< 40%) to be overfishing</p> <p>For fished area: Unknown</p>

Historical Stock Status Trajectory and Current Status



Standardised CPUE for longfin eels in Taranaki (AJ) from 1990–91 to 2014–15 (from Beentjes & McKenzie 2017). Also shown is the total estimated core fisher longfin catch in AJ from ECERs. Vertical dashed line indicates when the 4 kg maximum size was introduced in 2007–08 after which longfin eels 4 kg and over are not recorded on ECERs. Error bars are 95% confidence intervals. Before 2001, 16% of the catch was recorded as EEU (unidentified) and these catches are omitted.



Annual relative exploitation rate for longfin eels in the Taranaki (AJ). Because some catch of longfin was reported as EEU (unidentified) and has not been allocated to species, the exploitation rate is likely to have been higher than shown before 2001.

Fishery and Stock Trends

Recent Trend in Biomass or Proxy	Moderate decline in CPUE until 2003, increasing to 2012, and then declining to 2015
Recent Trend in Fishing intensity or Proxy	The relative exploitation rate declined steeply after 2003, and in 2015 was well below the series mean.
Other Abundance Indices	-
Trends in Other Relevant Indicators or Variables	Catches of longfin elvers at primary monitoring sites have fluctuated without trend since the series of reliable data begins in 1995–96, suggesting no overall trend in recruitment.

Projections and Prognosis

Stock Projections or Prognosis	Unlikely (< 40%) to decline in the medium term under current catch levels
Probability of Current Catch or TACC causing Biomass to remain below or to decline below Limits	For ESA, Soft Limit: Very Unlikely (< 10%) if catch remains at current levels For ESA, Hard Limit: Very Unlikely (< 10%) if catch remains at current levels
Probability of Current Catch or TACC causing Overfishing to continue or to commence	For ESA, Unlikely (< 40%) if catch remains at current levels

Assessment Methodology and Evaluation

Assessment Type	Level 2 – Partial Quantitative Stock Assessment	
Assessment Method	Standardised CPUE based on positive catches from commercial fyke net	
Assessment Dates	Latest assessment: 2017	Next assessment: 2019
Overall assessment quality rank	1 – High Quality	
Main data inputs (rank)	- Catch and effort data	1 – High Quality
Data not used (rank)	N/A	
Changes to Model Structure and Assumptions	-	
Major Sources of Uncertainty	- Standardised CPUE only provides an index of abundance for eels in areas fished by commercial fishers. Other potential issues with the CPUE indices include: <ul style="list-style-type: none"> • Low numbers of fishers • Uncertainty in target species after 2000 • Exclusion of zero catches 	

	<ul style="list-style-type: none"> • Changes in MLS and retention in early parts of the series and increased escape tube size from 25 mm to 31 mm in 2012–13 • Failure of some fishers to record on ECE returns all legal sized eels caught, not just those retained • Unrecorded release of > 4kg eels since 2007–08
--	---

Qualifying Comments

Because the commercial eel fishery has had a long history (beginning in the late 1960s), and indices of abundance are only available from the early 1990s, it is difficult to infer stock status from recent abundance trends, and these should therefore be interpreted with caution. Other sources of mortality, such as habitat alteration (historical and current) have also reduced abundance prior to the CPUE series.

The proportion of current longfin habitat in Taranaki (Statistical Area AJ) fished commercially during the period 2009–10 and 2013–14 is estimated at 17% (Table 15). The proportion of virgin habitat impacted by hydro dams, commercial fishing and other anthropogenic activity was estimated to be 24%.

Fishery Interactions

There has been no recorded bycatch in the commercial Taranaki eel fishery since 2000–01. Most bycatch species are usually returned alive.


6. FOR FURTHER INFORMATION

- Beentjes, M P (1998) Enhancement of Lake Hawea eel stocks by transfer of juveniles. *NIWA Technical Report 41*. 15 p.
- Beentjes, M P (1999) Size, age, and species composition of South Island commercial eel catches from market sampling (1997–98). *NIWA Technical Report 51*. 51 p.
- Beentjes, M P (2005) Monitoring commercial eel fisheries in 2003–04. *New Zealand Fisheries Assessment Report 2005/39*. 57 p.
- Beentjes, M P (2008) Monitoring commercial eel fisheries in 2003–04 and 2004–05. *New Zealand Fisheries Assessment Report 2008/19*. 43 p.
- Beentjes, M P (2011) Monitoring commercial eel fisheries in 2007–08 and 2008–09. *New Zealand Fisheries Assessment Report 2011/50*.
- Beentjes, M P (2013) Monitoring commercial eel fisheries: 2009–10 to 2011–12. *New Zealand Fisheries Assessment Report 2013/47*. 76 p.
- Beentjes, M.P. (2016). Monitoring commercial eel fisheries: 2003–04 to 2014–15. *New Zealand Fisheries Assessment Report 2016/50*. 83 p.
- Beentjes, M P; Boubée, J A T; Jellyman, D J; Graynoth, E (2005) Non-fishing mortality of freshwater eels (*Anguilla* spp.). *New Zealand Fisheries Assessment Report 2005/34*. 38 p.
- Beentjes, M P; Bull, B (2002) CPUE analyses of the commercial freshwater eel fishery. *New Zealand Fisheries Assessment Report 2002/18*. 55 p.
- Beentjes, M P; Chisnall, B L (1997) Trends in size and species composition and distribution of commercial eel catches. *New Zealand Fisheries Data Report 89*. 71 p.
- Beentjes, M P; Chisnall, B L (1998) Size, age, and species composition of commercial eel catches from market sampling, 1996–97. *NIWA Technical Report 29*. 124 p.
- Beentjes, M P; Chisnall, B L; Boubée, J A; Jellyman, D J (1997) Enhancement of the New Zealand eel fishery by elver transfers. *New Zealand Fisheries Technical Report 45*. 44 p.
- Beentjes, M P; Dunn, A (2003) CPUE analysis of the commercial freshwater eel fishery in selected areas, 1990–91 to 2000–01. *New Zealand Fisheries Assessment Report 2003/54*. 47 p.
- Beentjes, M P; Dunn, A (2008) Catch per unit effort (CPUE) analyses of the South Island commercial freshwater eel fishery, 1990–91 to 2005–06. *New Zealand Fisheries Assessment Report 2008/51*. 109 p.
- Beentjes, M P; Dunn, A (2010) CPUE analyses of the North Island commercial freshwater eel fishery, 1990–91 to 2006–07. *New Zealand Fisheries Assessment Report 2010/5*. 101 p.
- Beentjes, M P; Dunn, A (2013a) Catch per unit effort (CPUE) analyses and characterisation of the South Island commercial freshwater eel fishery, 1990–91 to 2009–10. *New Zealand Fisheries Assessment Report 2013/11*. 211 p.
- Beentjes, M P; Dunn, A (2013b) Catch per unit effort (CPUE) analyses and characterisation of the North Island commercial freshwater eel fishery, 1990–91 to 2011–12. *New Zealand Fisheries Assessment Report 2013/62*. 292 p.
- Beentjes, M P; Dunn, A (2014) Catch per unit effort (CPUE) analyses and characterisation of Te Waihora commercial freshwater eel fishery, 1990–91 to 2011–12. *New Zealand Fisheries Assessment Report 2014/17*. 46 p.
- Beentjes, M P; Dunn, A (2015) Catch per unit effort (CPUE) analyses and characterisation of the South Island commercial freshwater eel fishery, 1990–91 to 2012–13. *New Zealand Fisheries Assessment Report 2015/30*. 177 p.
- Beentjes, M P; McKenzie, A (2017) Catch per unit effort (CPUE) analyses and characterisation of the North Island commercial freshwater eel fishery, 1990–91 to 2014–15. *New Zealand Fisheries Assessment Report 2017/61*.
- Beentjes, M P; Jellyman, D J (2002) Evaluation of eel enhancement in Coopers Lagoon, South Canterbury. *New Zealand Fisheries Assessment Report 2002/7*. 18 p.
- Beentjes, M P; Jellyman, D J (2003) Enhanced growth of longfin eels, *Anguilla dieffenbachii*, transplanted into Lake Hawea, a high country lake in South Island, New Zealand. *New Zealand Journal of Marine and Freshwater Research* 37: 1–11.
- Beentjes, M P; Jellyman, D J (2011) Evaluation of the 1998 transfer of juvenile longfin eels into Lake Hawea from recaptures, and ageing validation based on otolith annual ring deposition. *New Zealand Fisheries Assessment Report 2011/19*. 37 p.
- Beentjes, M P; Sykes, J; Crow, S K (2016) GIS mapping of the longfin eel commercial fishery throughout New Zealand, and estimates of longfin habitat and proportion fished. *New Zealand Fisheries Assessment Report 2016/32*. 53 p.

- Boubée, J; Hudson, N (2009) Assessment of the state of the Rangitaiki River within Ngati Manawa rohe. NIWA client report HAM2009-100. 350 p.
- Chisnall, B L (2000) Enhancement of Maori customary eel fisheries in Lakes Taharoa and Harihari by transfer of juveniles. *NIWA Technical Report 91*. 16 p.
- Chisnall, B L; Beentjes, M P; Boubée, J A T; West, D W (1998) Enhancement of the New Zealand Eel Fisheries by transfers of elvers, 1996–97. *NIWA Technical Report: 37*. 55 p.
- Chisnall, B L; Jellyman, D J; Bonnett, M L; Sykes, J R E (2002) Spatial and temporal variability in size of glass eels (*Anguilla* spp.) in New Zealand. *New Zealand Journal of Marine and Freshwater Research* 36: 89–104.
- Chisnall, B L; Kalish, J M (1993) Age validation and movement of freshwater eels (*Anguilla dieffenbachii* and *A. australis*) in a New Zealand pastoral stream. *New Zealand Journal of Marine and Freshwater Research*. 27: 333–338.
- Crow, S; Jellyman, D (2010) Assessment of the eel population in the Pātea River catchment. Final Research Project for Ministry of Fisheries Project EEL2009-04. 38 p. (Unpublished report held by Fisheries New Zealand Wellington.)
- Davey, A J H; Jellyman, D J (2005) Sex determination in freshwater eels and management options for manipulation of sex. *Reviews in Fish Biology and Fisheries* 15, 37–52.
- Dekker, W (2001) Status of the European eel stock and fisheries. Proceedings of the International Symposium, Advances in Eel Biology, University of Tokyo, 28–30 September 2001: 50–52.
- Dekker, W; Casselman, J (2014) The 2003 Québec Declaration of Concern About Eel Declines—11 Years Later: Are Eels Climbing Back up the Slippery Slope? *Fisheries*, 39:12, 613–614, DOI: 10.1080/03632415.2014.979342
- Dunn, A; Beentjes, M P; Graynoth, E (2009) Preliminary investigations into the feasibility of assessment models for New Zealand longfin eels (*Anguilla dieffenbachii*). *New Zealand Fisheries Assessment Report 2009/30*. 42 p.
- Durif, C M F; Knutsen, J A; Johannessen, T; Vollestad, L A (2008) Analysis of European eel (*Anguilla anguilla*) time series from Norway. Report No 8/2008. Institute of Marine Research, Oslo. 22 p.
- Fu, D; Beentjes, M P; Dunn, A (2012) Further investigations into the feasibility of assessment models for New Zealand longfin eels (*Anguilla dieffenbachii*). Final Research Report for Project EEL200702. 77 p. (Unpublished report held by Fisheries New Zealand Wellington.)
- Graynoth, E; Booker, D J (2009) Biomass of longfin eels in medium to large rivers. *New Zealand Fisheries Assessment Report. No. 44*. 24 p.
- Graynoth, E; Jellyman, D J; Bonnett, M (2008) Spawning escapement of female longfin eels. *New Zealand Fisheries Assessment Report 2008/07*. 57 p.
- Horn, P L (1996) A review of age and growth data for New Zealand freshwater eels (*Anguilla* sp). New Zealand Fisheries Assessment Research Document 1996/6. 26 p. (Unpublished document held by NIWA library, Wellington.)
- Hoyle, S D; Jellyman, D J (2002) Longfin eels need reserves: modelling the effects of commercial harvest on stocks of New Zealand eels. *Marine and Freshwater Research*, 53: 887–895.
- Jellyman, D J (1993) A review of the fishery for freshwater eels in New Zealand. *NIWA. New Zealand Freshwater Fisheries Report 10*, 51 p.
- Jellyman, D J (1995) Longevity of longfinned eels *Anguilla dieffenbachia* in a New Zealand high country lake. *Ecology of Freshwater Fish* 4: 106–112.
- Jellyman, D J; Beentjes, M P (1998) Enhancement of the eel stocks of Coopers Lagoon, Canterbury Bight, by transfer of juvenile eels. *NIWA Technical Report 22*, 18 p.
- Jellyman, D J; Booker, D J; Watene, E (2009) Recruitment of glass eels (*Anguilla* spp.) in the Waikato River, New Zealand. Evidence of declining migrations? *Journal of Fish Biology* 74: 2014–2033.
- Jellyman, D J; Bowen, M (2009) Modelling larval migration routes and spawning areas of Anguillid eels of New Zealand and Australia. pp 255–274. In: Haro, A J; Smith, K L; Rulifson, R A; Moffitt, C M; Klauda, R J; Dadswell, M J; Cunjak, R A; Cooper, J E; Beal, K L; Avery, T S (eds). Challenges for Diadromous Fishes in a Dynamic Global Environment *American Fisheries Society Symposium* 69. Bethesda, Maryland.
- Jellyman, D J; Chisnall, B L; Dijkstra, L H; Boubée, J A T (1996) First record of the Australian longfinned eel, *Anguilla reinhardtii*, in New Zealand. *New Zealand Journal of Marine and Freshwater Research*, 47: 1037–1340.
- Jellyman, D J; Chisnall, B L; Sykes, J R E; Bonnett, M L (2002) Variability in spatial and temporal abundance of glass eels (*Anguilla* spp.) in New Zealand waterways. *New Zealand Journal of Marine and Freshwater Research* 36: 511–517.
- Jellyman, D J; Chisnall, B L; Todd, P R (1995) The status of the eel stocks of Lake Ellesmere. *NIWA science and technology series* 26, 62 p.
- Jellyman, D J; Graynoth, E; Francis, R I C C; Chisnall, B L; Beentjes, M P (2000) A review of the evidence for a decline in the abundance of longfinned eels (*Anguilla dieffenbachii*) in New Zealand. Final Research Report, Ministry of Fisheries Research Project EEL9802. 76 p. (Unpublished report held by Fisheries New Zealand Wellington.)
- Jellyman, D J; Sykes, J R E (2004) Review of glass eel recruitment, 1996–2002. Final Research Report, Ministry of Fisheries Research Project 13 p. (Unpublished report held by Fisheries New Zealand Wellington.)
- Martin, M L; Boubée, J A T; Bowman, E J (2013) Recruitment of freshwater elvers, 1995–2012. *New Zealand Fisheries Assessment Report 2013/50*. 111 p.
- Martin, M L; Boubée, J A T; Bowman, E J (2014) Recruitment of freshwater elvers. New Zealand Fisheries Progress Report. Ministry for Primary Industries. Wellington. 32 p. (Unpublished report held by Fisheries New Zealand Wellington.)
- Martin, M L; Boubée, J A T; Bowman, E J (2016) Recruitment of freshwater elvers 1995–2015. *New Zealand Fisheries Assessment Report 2016/46* 98 p.
- McCleave, J D; Jellyman, D J (2004) Male dominance in the New Zealand longfin eel population of a New Zealand river: Probable causes and implications for management. *North American Journal of Fisheries Management* 24, 490–505.
- Ministry of Fisheries. (2004) Setting of Sustainability and Other Management Controls for Stocks to be Introduced into the QMS on 1 October 2004: North Island shortfin and longfin eels (SFE, LFE). Final Advice Paper 25 June 2004. 232 p.
- Ministry of Fisheries. (2007) Review of Sustainability Measures and Other management Controls for 1 October 2007. Volume 1. Final Advice Paper and Summary of Recommendations. 5 September 2007. pp. 210–251.
- Moriarty, C; Dekker, W (1997) Management of the European eel. *Fisheries Bulletin (Dublin)* 15: 125 p.
- Quebec Declaration of Concern (2003) Worldwide decline of eel resources necessitates immediate action. pp 447–449 In: Casselman, J M; Cairns, D K (editors.) Eels at the edge: science status and conservation concerns. *American Fisheries Society Symposium* 58, Bethesda, Maryland, USA. 460 p.
- Richkus, W A; Whalen, K (2000) Evidence for a decline in the abundance of the American eel, *Anguilla rostrata* (LeSueur), in North America since the early 1980s. *Dana* 12: 83–97.
- Speed, S R; Browne, G N; Boyd, R O (2001) Assessment and Monitoring of Commercial Eel Fisheries. Final Research Report for Ministry of Fisheries Research Project EEL9801. 178 p. (Unpublished report held by Fisheries New Zealand Wellington.)
- Todd, P R (1980) Size and age of migrating New Zealand freshwater eels (*Anguilla* spp.). *New Zealand Journal of Marine and Freshwater Research* 14, 283–293.
- Townsend, A J; de Lange, P J; Duffy, C A J; Miskelly, C M; Molloy, J; Norton, D A (2008) New Zealand Threat Classification System manual. Department of Conservation, Wellington. 35 p.
- Vollestad, L A; Jonsson, B (1988) A 13-year study of the population dynamics and growth of the European eel (*Anguilla Anguilla*) in a Norwegian River: evidence for density-dependent mortality, and development of a model for predicting yield. *Journal of Animal Ecology* 88: 983–997.

FROSTFISH (FRO)*(Lepidopus caudatus)*

Para, Taharangi, Hikau

**1. FISHERY SUMMARY****1.1 Commercial fisheries**

Frostfish are predominantly taken as bycatch from target trawl fisheries on jack mackerel and hoki and to a lesser extent, arrow squid, barracouta and gemfish. These fisheries are predominantly targeted by larger vessels owned or chartered by New Zealand fishing companies. Target fishing for frostfish is reported from the west coast of both the South Island and North Island and at Puysegur Bank, with the best catches taken from the west coast of the South Island.

The main areas reporting frostfish catches are to the west of New Zealand primarily in QMA 7 on the west coast of the South Island and to a lesser extent QMA 8 and 9 in the north and south Taranaki Bight. The highest annual catches are associated with hoki fishing during winter (since 1986–87) and jack mackerel fishing during late spring and early summer. The proportion of catch coming from these two main fisheries has varied over time. Sources of error in the catch figures include unreported catch and discarded catch. Compliance investigations have shown that damaged and small hoki were recorded as frostfish by some vessels.

Since the mid-2000s, most frostfish landings have come from the trawl fishery targeting jack mackerel (JMA) in the North and South Taranaki Bights and off the west coast of the South Island (Statistical Areas 035 to 041; FRO 7, 8, 9). In 2009–10, over 80% of the national frostfish landings came from this fishery. Since 1999–2000, the fishery has been dominated by seven vessels which use midwater trawling exclusively. Catches of frostfish have become more concentrated on two distinct periods, October to January and June to July, and in the north and south Taranaki Bight (Statistical Areas 037, 040, 041) rather than the west coast of the South Island (Statistical Areas 034, 035, 036).

No catch data from deepwater vessels for frostfish are available prior to the introduction of the EEZ in 1978 (Table 1). Frostfish were introduced into the QMS from 1 October 1998. The total reported landings and TACCs for each QMA are given in Table 1 and 2, while Figure 1 shows the historical landings and TACC values for the main FRO stocks. An allowance of 2 t was made for non-commercial catch in each of FRO (1, 2, 7 and 9) and therefore TACs for these stocks are 2 t higher than the TACCs. TACCs were increased from 1 October 2006 in FRO 2 to 110 t, in FRO 3 to 176 t

FROSTFISH (FRO)

and in FRO 4 to 28 t. In these stocks landings were above the TACC for a number of years and the TACCs were increased to the average of the previous seven years plus an additional 10% (Table 3).

1.2 Recreational fisheries

Frostfish are occasionally taken by recreational fishers. Small numbers have been reported from recreational diary surveys, mainly in QMA 1, and rarely in QMA 2 and 9.

1.3 Customary non-commercial fisheries

No quantitative information is available on the current level of customary non-commercial take. Maori have collected beach cast frostfish in the past (Graham 1956).

1.4 Illegal catch

No information is available.

1.5 Other sources of mortality

No information is available on other sources of mortality.

Table 1: Reported landings (t) of frostfish by fishing year and area, by foreign licensed and joint venture vessels, 1978–79 to 1983–83. The EEZ areas (see figure 2 of Baird & McKoy 1988) correspond approximately to the QMAs as indicated. Fishing years are from 1 April to 31 March. The 1983–83 is a 6 month transitional period from 1 April to 30 September. No data are available for the 1980–81 fishing year.

EEZ area	B	C(M)	C(-)	D	E	F	G	H	Total
QMA	1 & 2	3	3	4	6	5	7	8 & 9	
1978–79	5	1	6	0	1	0	1 283	226	1 522
1979–80	13	0	1	23	1	1	26	151	216
1980–81	-	-	-	-	-	-	-	-	-
1981–82	0	5	2	19	1	4	55	464	550
1982–83	0	1	0	9	3	1	56	1 545	1 615
1983–83	0	1	1	1	1	1	22	123	150

Table 2: Reported landings (t) for the main QMAs from 1931 to 1982 [Continued on next page].

Year	FRO 1	FRO 2	FRO 3	FRO 4	FRO 5	Year	FRO 1	FRO 2	FRO 3	FRO 4	FRO5
1931–32	0	0	0	0	0	1957	0	0	0	0	0
1932–33	0	0	0	0	0	1958	0	0	0	0	0
1933–34	0	0	0	0	0	1959	0	0	0	0	0
1934–35	0	0	0	0	0	1960	0	0	0	0	0
1935–36	0	0	0	0	0	1961	0	0	0	0	0
1936–37	0	0	0	0	0	1962	0	0	0	0	0
1937–38	0	0	0	0	0	1963	0	0	0	0	0
1938–39	0	0	0	0	0	1964	0	0	0	0	0
1939–40	0	0	0	0	0	1965	0	0	0	0	0
1940–41	0	0	0	0	0	1966	0	5	0	0	0
1941–42	0	1	0	0	0	1967	0	0	0	0	0
1942–43	0	0	0	0	0	1968	0	0	0	0	0
1943–44	0	0	0	0	0	1969	0	0	0	0	0
1944	0	0	0	0	0	1970	0	0	0	0	0
1945	0	0	0	0	0	1971	0	0	0	0	0
1946	0	0	0	0	0	1972	0	0	0	0	0
1947	3	0	0	0	0	1973	0	0	0	0	0
1948	0	0	0	0	0	1974	0	0	0	0	0
1949	0	0	0	0	0	1975	0	0	0	0	0
1950	0	0	0	0	0	1976	0	0	0	0	0
1951	0	0	0	0	0	1977	0	0	0	0	0
1952	0	0	0	0	0	1978	1	4	2	0	0
1953	0	0	0	0	0	1979	1	14	4	19	1
1954	0	0	0	0	0	1980	0	0	2	20	7
1955	0	0	0	0	0	1981	0	0	6	25	3
1956	0	0	0	0	0	1982	4	0	0	8	13

Table 2 continued.

Year	FRO 6	FRO 7	FRO 8	FRO 9	Year	FRO 6	FRO 7	FRO 8	FRO 9
1931–32	0	0	0	0	1957	0	0	0	0
1932–33	0	0	0	0	1958	0	0	0	0
1933–34	0	0	0	0	1959	0	0	0	0
1934–35	0	0	0	0	1960	0	0	0	0
1935–36	0	0	0	0	1961	0	0	0	0
1936–37	0	0	0	0	1962	0	0	0	0
1937–38	0	0	0	0	1963	0	0	0	0
1938–39	0	0	0	0	1964	0	0	0	0
1939–40	0	0	0	0	1965	0	0	0	0
1940–41	0	0	0	0	1966	0	0	0	0
1941–42	0	0	0	0	1967	0	0	0	0
1942–43	0	0	0	0	1968	0	0	0	0
1943–44	0	0	0	0	1969	0	0	1	0
1944	0	0	0	0	1970	0	0	1	0
1945	0	0	0	0	1971	0	0	0	0
1946	0	0	0	0	1972	0	0	0	0
1947	0	0	0	1	1973	0	0	0	0
1948	0	0	0	0	1974	0	0	0	0
1949	0	0	0	0	1975	0	0	0	0
1950	0	0	0	0	1976	0	0	0	0
1951	0	0	0	0	1977	0	0	0	0
1952	0	0	0	0	1978	0	782	30	16
1953	0	0	0	0	1979	1	614	93	88
1954	0	0	0	0	1980	1	41	54	10
1955	0	0	0	0	1981	0	327	226	209
1956	0	0	0	0	1982	0	132	385	546

Notes:

The 1931–1943 years are April–March but from 1944 onwards are calendar years. Data up to 1985 are from fishing returns: Data from 1986 to 1990 are from Quota Management Reports. Data for the period 1931 to 1982 are based on reported landings by harbour and are likely to be underestimated as a result of under-reporting and discarding practices. Data includes both foreign and domestic landings.

Table 3: Reported landings (t) of frostfish by QMA and fishing year, 1983–84 to 2016–17. The data in this table has been updated from that published in the 1998 Plenary Report by using the data up to 1996–97 in table 26 on p. 244 of the “Review of Sustainability Measures and Other Management Controls for the 1998–99 Fishing Year - Final Advice Paper” dated 6 August 1998. Data since 1997–98 based on catch and effort returns (where area was not reported catch was pro-rated across all QMAs). There are no landings reported from QMA 10. [Continued on next page].

Fishstock FMA	FRO 1		FRO 2		FRO 3		FRO 4		FRO 5	
	Landings	TACC	Landings	TACC	Landings	TACC	Landings	TACC	Landings	TACC
1983–84	2	-	0	-	0	-	10	-	28	-
1984–85	0	-	0	-	2	-	1	-	100	-
1985–86	0	-	0	-	9	-	2	-	258	-
1986–87	4	-	4	-	5	-	6	-	71	-
1987–88	2	-	0	-	3	-	1	-	20	-
1988–89	115	-	0	-	1	-	0	-	15	-
1989–90	397	-	0	-	58	-	0	-	146	-
1990–91	45	-	24	-	224	-	0	-	496	-
1991–92	46	-	3	-	143	-	0	-	337	-
1992–93	80	-	9	-	51	-	0	-	0	-
1993–94	100	-	19	-	168	-	0	-	0	-
1994–95	55	-	14	-	120	-	0	-	87	-
1995–96	80	-	40	-	72	-	29	-	0	-
1996–97	198	-	6	-	12	-	4	-	8	-
1997–98	309	-	273	-	35	-	< 1	-	9	-
1998–99	146	149	134	20	39	128	< 1	5	19	135
1999–00	84	149	161	20	97	128	< 1	5	57	135
2000–01	76	149	194	20	107	128	48	5	33	135
2001–02	64	149	67	20	176	128	81	5	59	135
2002–03	127	149	66	20	268	128	15	5	63	135
2003–04	98	149	52	20	19	128	7	5	14	135
2004–05	130	149	38	20	427	128	15	5	20	135
2005–06	132	149	40	20	45	128	31	5	17	135
2006–07	76	149	31	110	21	176	13	28	16	135
2007–08	44	149	30	110	31	176	7	28	5	135
2008–09	36	149	24	110	6	176	10	28	2	135
2009–10	36	149	24	110	15	176	3	28	4	135
2010–11	52	149	41	110	< 1	176	4	28	14	135
2011–12	34	149	15	110	8	176	14	28	3	135
2012–13	21	149	18	110	32	176	2	28	4	135
2013–14	40	149	34	110	63	176	15	28	11	135
2014–15	54	149	41	110	13	176	69	28	14	135
2015–16	70	149	46	110	10	176	13	28	8	135
2016–17	75	149	52	110	9	176	9	28	27	135

FROSTFISH (FRO)

Table 3 Continued

Fishstock FMA	FRO 6		FRO 7		FRO 8		FRO 9		Total	
	Landings	TACC	Landings	TACC	Landings	TACC	Landings	TACC	Landings	TACC
1983-84	7	-	432	-	539	-	457	-	1 475	-
1984-85	0	-	214	-	455	-	129	-	901	-
1985-86	0	-	344	-	574	-	226	-	1 415	-
1986-87	4	-	1 089	-	898	-	190	-	2 272	-
1987-88	0	-	3 466	-	875	-	22	-	4 391	-
1988-89	3	-	1 950	-	413	-	455	-	2 952	-
1989-90	29	-	1 370	-	132	-	0	-	2 132	-
1990-91	67	-	3 029	-	539	-	0	-	4 424	-
1991-92	7	-	2 295	-	750	-	1	-	3 582	-
1992-93	0	-	1 360	-	1 165	-	0	-	2 665	-
1993-94	0	-	1 998	-	696	-	12	-	2 993	-
1994-95	0	-	3 069	-	388	-	7	-	3 740	-
1995-96	0	-	1 536	-	22	-	9	-	1 788	-
1996-97	0	-	2 881	-	126	-	93	-	3 328	-
1997-98	0	-	2 590	-	143	-	205	-	3 564	-
1998-99	0	11	2 461	2 623	156	649	33	138	2 989	3 858
1999-00	< 1	11	917	2 623	28	649	48	138	1 392	3 858
2000-01	< 1	11	1 620	2 623	303	649	43	138	2 424	3 858
2001-02	< 1	11	2 303	2 623	138	649	25	138	2 913	3 858
2002-03	< 1	11	1 025	2 623	621	649	67	138	2 252	3 858
2003-04	< 1	11	959	2 623	293	649	367	138	1 809	3 858
2004-05	< 1	11	934	2 623	770	649	327	138	2 661	3 858
2005-06	< 1	11	888	2 623	787	649	181	138	2 119	3 858
2006-07	< 1	11	951	2 623	722	649	142	138	1 972	4 019
2007-08	< 1	11	906	2 623	678	649	136	138	1 837	4 019
2008-09	< 1	11	576	2 623	605	649	110	138	1 369	4 019
2009-10	< 1	11	382	2 623	686	649	238	138	1 389	4 019
2010-11	< 1	11	248	2 623	578	649	167	138	1 106	4 019
2011-12	< 1	11	500	2 623	893	649	198	138	1 665	4 019
2012-13	< 1	11	570	2 623	890	649	278	138	1 814	4 019
2013-14	< 1	11	880	2 623	814	649	261	138	2 120	4 019
2014-15	< 1	11	1 027	2 623	732	649	373	138	2 322	4 019
2015-16	< 1	11	1 063	2 623	692	649	310	138	2 212	4 019
2016-17	< 1	11	1 164	2 623	553	649	96	138	1 986	4 019




Figure 1: Reported commercial landings and TACC for the eight main FRO stocks. FRO 1 (Auckland East)[Continued on next page].




Figure 1: Reported commercial landings and TACC for the eight main FRO stocks. From top: FRO 2 (Central East), FRO 3 (South East Coast), FRO 4 (South East Chatham Rise) [Continued on next page].

FROSTFISH (FRO)




Figure 1: Reported commercial landings and TACC for the eight main FRO stocks. From top: FRO 5 (Southland), FRO 7 (Challenger), FRO 8 (Central West). [Continued on next page].




Figure 1 [Continued]: Reported commercial landings and TACC for the eight main FRO stocks. FRO 9 (Auckland West). Note that these figures do not show data prior to entry into the QMS.

2. BIOLOGY

Frostfish are widely distributed throughout the continental shelf and upper slopes of all oceans, except the North Pacific, and have a benthopelagic lifestyle. In New Zealand, frostfish are found from about 34°S to 49°S, but are most common between 36°S and 44°S. They occur mainly in depths of 50–600 m with the largest catches made at around 200 m bottom depth. Preferred bottom temperatures range between 10 and 16°C. There is one species of *Lepidopus* recorded from New Zealand waters. However, scabbardfishes (*Benthodesmus* species) and the false frostfish (*Paradiplospinosus gracilis*) may be confused with small *Lepidopus caudatus*.

Frostfish reach a maximum length of 165 cm (fork length) around New Zealand, although the same species may reach 205 cm and 8 kg weight in the eastern North Atlantic (Nakamura & Parin 1993). In the northwestern Mediterranean males reach sexual maturity at 97 cm and a maximum length of 176 cm, whilst females reach sexual maturity at 111 cm and a maximum length of 196 cm (Demestre et al 1993).

The adults probably congregate in the late spring months, and spawn during the summer and autumn over the mid to outer shelf. Fertilisation has been calculated to take place between noon and sunset at depths greater than 50 m where the surface waters have a temperature of 17.5 to 22.0°C (Robertson 1980).

A recent study developed ageing methods and estimated growth rates for frostfish from the west coast of New Zealand (Horn 2013). This study confirmed that frostfish are fast growing and relatively short lived. Most fish reach 100 cm FL (fork length) by the end of their third year and the maximum estimated age for both sexes was 10.6 years. The von Bertalanffy parameters estimated for both sexes combined were: $L_{\infty}=137$ cm, $k=0.505$ yr⁻¹, $t_0=0.07$ yr. The estimated growth curves were similar, for the first four years, to those estimated for northern hemisphere frostfish, although the asymptotic length is lower. Horn (2013) estimated the instantaneous rate of natural mortality to be 0.6 yr⁻¹ based on 1% of the population reaching 7–8 years of age.

A length-weight relationship for New Zealand frostfish is available from the *Kaharoa* trawl surveys (Horn 2013).

Frostfish migrate into mid-water at night and feed on crustaceans, small fish and squid (Nakamura & Parin 1993). Euphausiids and *Pasiphaea* spp. (both crustaceans) are the most common prey of frostfish in the northwest Mediterranean (Demestre et al 1993). In Tasmanian waters, the diet of frostfish consists mainly of myctophids and euphausiids (Blaber & Bulman 1987).

Frostfish are distributed widely in temperate seas but are most commonly reported in the north-

FROSTFISH (FRO)

eastern Atlantic (including the Mediterranean), in the southern Atlantic off Namibia and South Africa, and in the south-west Pacific around Australia and New Zealand (Nakamura & Parin 1993, Froese & Pauly 2012). Morphometric studies have shown differences in dorsal-fin pigmentation and meristic characteristics between north-eastern Atlantic and southern Atlantic populations (Mikhailin 1977). Genome sequencing of frostfish showed strong genetic differentiation between the northern and southern hemisphere populations and suggests that there are two distinct biological species (Ward et al 2008).

Robertson (1980) examined the seasonality and location of frostfish spawning based on the occurrence of planktonic eggs. He concluded that spawning probably occurs around all of New Zealand except for the south-east coast and adults probably congregate in the late spring months, and spawn during the summer and autumn over the mid to outer shelf. Fertilisation was calculated to take place between noon and sunset at depths greater than 50 m where the surface waters have a temperature of 17.5 to 22.0°C. Analysis of data on female gonad stages from the scientific observer programme (see Section 6.1) suggests that for the west coast of both the North and South Islands frostfish have a protracted spawning period starting in mid-winter with a peak from summer to early autumn.

Biological parameters relevant to the stock assessment are shown in Table 4.

Table 4: Estimates of biological parameters for frostfish.

Fishstock	Estimate						Source
1. Natural mortality (<i>M</i>)							
All stocks	<i>M</i> = 0.6 y ⁻¹ considered best estimate for all areas for both sexes						Horn (2013)
2. Weight = <i>a</i> (length) ^b (Weight in g, length in cm fork length)							
	<i>a</i>		<i>b</i>				
WCSI trawl surveys	0.000407		3.155		Horn (2013)		
3. von Bertalanffy growth parameters							
	Male			Female			
	<i>L</i> _∞	<i>k</i>	<i>t</i> ₀	<i>L</i> _∞	<i>k</i>	<i>t</i> ₀	
WCSI	129.2	0.56	0.08	143.5	0.457	-0.04	Horn (2013)

3. STOCKS AND AREAS

Spawning areas identified from eggs taken in plankton tows include the outer shelf from the Bay of Islands to south of East Cape, and an area off Fiordland (Robertson 1980). No eggs were recorded from the south-east coast of the South Island and no spawning has been recorded on the Chatham Rise. Spawning is also known to take place on the west coast of the South Island in March.

Juvenile frostfish (less than 30 cm) have been reported from trawl surveys in the Bay of Plenty, the Hauraki Gulf, off Northland, the west coast of the North Island and the west coast of the South Island.

The occurrence of spawning in three areas at similar times of year and the distribution of frostfish from catches suggest that there may be at least three separate stocks. A fourth stock is also possible based on known distribution of juveniles and adults and analogies with other species which often have a separate Chatham Rise stock. Bagley et al (1998) proposed the following Fishstock areas for management of frostfish: FRO 1: (FMA 1 and 2); FRO 3: (FMA 3 and 4); FRO 5: (FMA 5 and 6) and FRO 7: (FMA 7, 8, and 9). There have been no reported landings from QMA 10. TACs were set for each QMA (1–9) in 1998 and each FMA is managed separately.

4. STOCK ASSESSMENT

There are no stock assessments available for any stocks of frostfish and therefore estimates of biomass and yields are not available.

4.1 Estimates of fishery parameters and abundance

No estimates of fishery parameters are available for frostfish.

Biomass indices on frostfish are available from trawl surveys carried out by different vessels (Table 5). Few surveys cover the central west coast of New Zealand where the commercial catch records highest landings. The catchability of frostfish is not known but, because they are known to occur frequently well off the bottom, catchability is expected to be low and variable between surveys.

Table 5: Doorspread biomass indices (t) and CVs (%) of frostfish from random stratified trawl surveys 1981–2013.

Vessel	Trip Code	Depth Range (m)	Biomass index (t)	CV (%)	Date
QMA 1					
Bay of Plenty					
<i>Kaharoa</i>	KAH9004	10–150	246	87	February/March 1990
<i>Kaharoa</i>	KAH9202	10–150	92	48	February 1992
<i>Kaharoa</i>	KAH9601	10–250	328	49	February 1996
<i>Kaharoa</i>	KAH9902		193	34	February 1999
QMA 2					
<i>Kaharoa</i>	KAH9304	20–400	573	38	March/April 1993
<i>Kaharoa</i>	KAH9402	20–400	1 079	40	February/March 1994
<i>Kaharoa</i>	KAH9502	20–400	493	22	February/March 1995
<i>Kaharoa</i>	KAH9602	20–400	693	17	February/March 1996
QMA 7 & 8					
<i>Tomi Maru</i>		30–300	2 173	22	December 1980 - January 1981
<i>Shinkai Maru</i>	SHI8102	20–300	6 638	12	October/November 1981
<i>Cordella</i>	COR9001	25–300	2 189	20	February/March 1990
QMA 7 (WCSI)					
<i>Kaharoa</i>	KAH9006	20–400	121	27	March/April 1990
<i>Kaharoa</i>	KAH9204	20–400	24	29	March/April 1992
<i>Kaharoa</i>	KAH9404	20–400	53	37	March/April 1994
<i>Kaharoa</i>	KAH9504	20–400	89	31	March/April 1995
<i>Kaharoa</i>	KAH9701	20–400	259	32	March/April 1997
<i>Kaharoa</i>	KAH0004	20–400	316	16	March/April 2000
<i>Kaharoa</i>	KAH0304	20–400	494	22	March/April 2003
<i>Kaharoa</i>	KAH0504	20–400	423	45	March/April 2005
<i>Kaharoa</i>	KAH0704	20–400	529	38	March/April 2007
<i>Kaharoa</i>	KAH0904	20–400	835	34	March/April 2009
<i>Kaharoa</i>	KAH1104	20–400	251	28	March/April 2011
<i>Kaharoa</i>	KAH1305	20–400	424	24	March/April 2013
WCSI south of 41° 30'					
<i>James Cook</i>	JCO8311	25–450	183	34	September/October 1983
<i>James Cook</i>	JCO8415	25–450	181	25	August/September 1985

4.2 Biomass estimates

No biomass estimates are available for frostfish.

4.3 Yield estimates and projections

MCY cannot be determined as only a small percentage (less than 2%) of the reported catch in recent years is from target fishing. Annual catches are likely to vary according to effort targeting other species in areas of frostfish abundance. It is therefore not possible to choose a catch history which represents a period of stable and unrestricted effort in order to estimate yields. Other problems include under-reporting of frostfish catches and restrictions targeting frostfish in QMAs 3, 4, 5, and 6.

There are no reliable data on current biomass; CAY was therefore not estimated.

FROSTFISH (FRO)

4.4 Other factors

None available.

5. STATUS OF THE STOCKS

Estimates of current and reference biomass are not available. The stock structure is uncertain; the fishery is variable and almost entirely a bycatch of other target fisheries. No age data or estimates of abundance are available.

It is therefore not possible to estimate yields. It is not known if recent catches are sustainable or whether they are at levels that will allow the stock to move towards a size that will support the maximum sustainable yield.

TACCs and reported landings for the 2016–17 fishing year are summarised in Table 6.

Table 6: Summary of TACCs (t), and reported landings (t) of frostfish for the most recent fishing year.


Fishstock		FMA	2016–17 Actual TACC	2016–17 Reported landings
FRO 1	Auckland (East)	1	149	75
FRO 2	Central (East)	2	110	52
FRO 3	South-east (Coast)	3	176	9
FRO 4	South-east (Chatham)	4	28	9
FRO 5	Southland	5	135	27
FRO 6	Sub-Antarctic	6	11	< 1
FRO 7	Challenger	7	2 623	1 164
FRO 8	Central (West)	8	649	553
FRO 9	Auckland (West)	9	138	96
FRO 10	Kermadec	10	0	0
Total			4 019	1 986

6. FOR FURTHER INFORMATION

- Bagley, N W; Schofield, K A; Colman, J A (1998) A summary of biological and commercial landings, and a stock assessment of the frostfish *Lepidopus caudatus* Euphrasen, 1788 (Pisces: Trichiuridae), in New Zealand waters. New Zealand Fisheries Assessment Research Document 1998/23. 28 p. (Unpublished document held in NIWA library.)
- Baird, G G; McKoy, J L (Comps. and Eds.) (1988) Papers from the workshop to review fish stock assessments for the 1987–88 New Zealand fishing year. 300 p. (Preliminary discussion paper, held in NIWA library, Wellington.)
- Bentley, N; Kendrick, T H; MacGibbon, D J (2014) Fishery characterisation and catch-per-unit-effort analyses for frostfish (*Lepidopus caudatus*), 1989–90 to 2009–10. *New Zealand Fisheries Assessment Report* 2014/25. 131 p.
- Blaber, S J M; Bulman, C M (1987) Diets of fishes of the upper continental slope of eastern Tasmania: Content, caloric values, dietary overlap and trophic relationships. *Marine Biology* 95(3): 345–356.
- Demestre, M; Moli, B; Recasens, P; Sanchez, P (1993) Life history and fishery of *Lepidopus caudatus* (Pisces: Trichiuridae) in the Catalan Sea (Northwestern Mediterranean). *Marine Biology* 115: 23–32.
- FAO (1993) Yearbook of Fishery Statistics, 1993. Catches and landings (Vol. 76) *FAO Statistics Series* No. 123.
- Froese, R; Pauly, D (editors). (2012) FishBase. World Wide Web electronic publication. www.fishbase.org
- Graham, D H (1956) *A Treasury of New Zealand fishes*. A.H. & A.W. Reed Ltd. 424 p.
- Horn, P L (2013) Age determination of frostfish (*Lepidopus caudatus*) off west coast South Island. *New Zealand Fisheries Assessment Report* 2013/21. 24 p.
- Mikhailin, S V (1977) The intraspecific variability of the frostfish, *Lepidopus caudatus*. *Vopr Ikhtiol.* 17(2):226–236.
- Nakamura, I; Parin, N V (1993) FAO species catalogue vol. 15. Snake mackerels and cutlassfishes of the world. *FAO Fisheries Synopsis*. No. 125. 136 p.
- Robertson, D A (1980) Spawning of the frostfish, *Lepidopus caudatus* (Pisces: Trichiuridae), in New Zealand waters. *New Zealand Journal of Marine and Freshwater Research* 14: 129–136.
- Schofield, K A; Bagley, N W; Colman, J A (1998) A summary of biology and commercial landings, and a stock assessment of the frostfish, *Lepidopus caudatus* Euphrasen, 1788 (Pisces: Trichiuridae) in New Zealand waters. New Zealand Fisheries Assessment Research Document 1998/23. 28 p. (Unpublished report held by NIWA library, Wellington.)
- Ward, R D; Costa F O; Holmes, B H; Steinke, D (2008) DNA barcoding of shared fish species from the North Atlantic and Australasia: minimal divergence for most taxa, but *Zeus faber* and *Lepidopus caudatus* each probably constitute two species. *Aquatic Biology* 3: 71–78.

GARFISH (GAR)*(Hyporhamphus ihi)*

Takeke

**1. FISHERY SUMMARY**

Garfish was introduced into the QMS from 1 October 2002 with allowances, TACCs and TACs as shown in Table 1. These have not changed.

Table 1: Recreational and Customary non-commercial allowances, TACCs and TACs (t) of garfish by Fishstock.

Fishstock	Recreational Allowance	Customary Non-Commercial Allowance	TACC	TAC
GAR 1	20	10	25	55
GAR 2	8	4	5	17
GAR 3	2	1	5	8
GAR 4	1	1	2	4
GAR 7	10	5	8	23
GAR 8	8	4	5	17
GAR 10	0	0	0	0

1.1 Commercial fisheries

Garfish landings were first recorded in 1933, and a minor fishery must have existed before this (Table 2). Moderate quantities of garfish can be readily caught by experienced fishers, it is a desirable food fish, and informal sales at beaches or from wharves are likely to have been made from the late 1800s onwards. Reported landings to 1990 almost certainly understate the actual “commercial” catch.

Table 2: Reported total New Zealand landings (t) of garfish from 1931 to 1990.

Year	Landings	Year	Landings	Year	Landings	Year	Landings	Year	Landings	Year	Landings
1931	—	1941	1	1951	4	1961	3	1971	11	1981	7
1932	—	1942	1	1952	7	1962	4	1972	4	1982	11
1933	1	1943	1	1953	6	1963	4	1973	10	1983	12
1934	—	1944	2	1954	8	1964	2	1974	6	1984	13
1935	—	1945	9	1955	9	1965	2	1975	2	1975	8
1936	—	1946	3	1956	7	1966	3	1976	5	1986	14
1937	—	1947	2	1957	2	1967	4	1977	5	1987	36
1938	—	1948	1	1958	2	1968	3	1978	15	1988	20
1939	4	1949	6	1959	4	1969	5	1979	12	1989	15
1940	6	1950	2	1960	6	1970	13	1980	12	1990	24

Source: Annual Reports on Fisheries (Marine Department/Ministry of Agriculture & Fisheries) to 1974, and subsequent MAF data.

GARFISH (GAR)

By 1990 reported landings were in the range 20–40 t, and the total catches may have reached 50 t. Reported catches and landings through the 1990s have been of a similar order of magnitude although catches have declined since the 2000–01 fishing season (Table 3).

Largest catches and landings (8–31 t) were made in FMA 1, mostly in Statistical Area 003 (southern east Northland) and 009 (central Bay of Plenty). Small (2–6 t) quantities were taken in FMA 7, almost entirely in area 017 (Marlborough Sounds). Only minor and intermittent catches and landings were made elsewhere. The most consistent catches were taken by beach seine, with some catches by lampara net. Most of the catch is reported as targeted.

In the early 1990s about 50 vessels reported a catch or landing in a year; by the late 1990s this had declined to 20–30. Most vessels reported garfish in only a few years. Annual reported landings have fluctuated between 9 and 26 tonnes since 2010–11.

Table 3: Reported catches or landings (t) of garfish by Fishstock from 1990–91 to 2016–17*. Prior to 2001–02 the catches or landings (t) of garfish were reported by FMA.

Fishstock FMA (s)	GAR 1		GAR 2		GAR 3		GAR 4	
	Landings	TACC	Landings	TACC	Landings	TACC	Landings	TACC
1990–91†	31	-	< 1	-	2	-	-	-
1991–92†	22	-	< 1	-	1	-	-	-
1992–93†	14	-	< 1	-	1	-	-	-
1993–94†	23	-	0	-	2	-	-	-
1994–95†	17	-	< 1	-	< 1	-	-	-
1995–96†	15	-	< 1	-	1	-	-	-
1996–97†	15	-	< 1	-	1	-	-	-
1997–98†	21	-	< 1	-	< 1	-	-	-
1998–99†	19	-	< 1	-	< 1	-	-	-
1999–00†	17	-	< 1	-	< 1	-	-	-
2000–01†	11	-	0	-	< 1	-	-	-
2001–02†	8	25	0	5	< 1	5	0	2
2002–03†	6	25	0	5	< 1	5	0	2
2003–04†	11	25	0	5	0	5	0	2
2004–05†	13	25	< 1	5	0	5	0	2
2005–06†	7	25	< 1	5	1	5	0	2
2006–07†	10	25	0	5	0	5	0	2
2007–08†	8	25	0	5	0	5	< 1	2
2008–09†	10	25	0	5	0	5	0	2
2009–10†	9	25	0	5	0	5	0	2
2010–11†	11	25	0	5	< 1	5	0	2
2011–12†	8	25	0	5	0	5	0	2
2012–13	12	25	< 1	5	< 1	5	0	2
2013–14	15	25	0	5	0	5	0	2
2014–15	16	25	0	5	0	5	0	2
2015–16	25	25	0	5	0	5	0	2
2016–17	26	25	0	5	0	5	0	2

Fishstock FMA (s)	GAR 7		GAR 8		GAR 10		Total	
	Landings	TACC	Landings	TACC	Landings	TACC	Landings [#]	TACC
1990–91†	4	-	1	-	0	-	38	-
1991–92†	6	-	0	-	0	-	29	-
1992–93†	2	-	2	-	0	-	18	-
1993–94†	2	-	0	-	0	-	26	-
1994–95†	2	-	0	-	0	-	19	-
1995–96†	3	-	< 1	-	0	-	19	-
1996–97†	5	-	< 1	-	0	-	20	-
1997–98†	4	-	1	-	0	-	27	-
1998–99†	6	-	1	-	0	-	26	-
1999–00†	4	-	< 1	-	0	-	21	-
2000–01†	2	-	0	-	0	-	13	-
2001–02†	3	8	0	5	0	0	11	50
2002–03†	< 1	8	0	5	0	0	6	50
2003–04†	1	8	< 1	5	0	0	12	50
2004–05†	0	8	< 1	5	0	0	13	50
2005–06†	0	8	0	5	0	0	9	50
2006–07†	< 1	8	< 1	5	0	0	10	50
2007–08†	< 1	8	0	5	0	0	8	50
2008–09†	1	8	0	5	0	0	11	50
2009–10†	3	8	0	5	0	0	12	50

Table 3 [Continued]

Fishstock FMA (s)	GAR 7		GAR 8		GAR 10		Total	
	Landings	TACC	Landings	TACC	Landings	TACC	Landings [#]	TACC
2010–11†	1	8	0	5	0	0	13	50
2011–12†	< 1	8	< 1	5	0	0	9	50
2012–13	0	8	0	5	0	0	12	50
2013–14	0	8	0	5	0	0	15	50
2014–15	<1	8	0	5	0	0	16	50
2015–16	<1	8	0	5	0	0	25	50
2016–17	0	8	0	5	0	0	26	50

* Listed as landings, but are the higher of catch or landing values. There were relatively small differences between the two series.

† CELR data.

Note totals may not match figures in the tables due to rounding errors.

1.2 Recreational fisheries

There is a small and specific recreational fishery using beach seines, but no information on the size of catch.

1.3 Customary non-commercial fisheries

Quantitative information on the current level of customary non-commercial catch is not available.

1.4 Illegal catch

Estimates of illegal catch are not available, but this is probably insignificant or nil.

1.5 Other sources of mortality

There may be some accidental catches of garfish in small-mesh nets (purse seines, lampara nets, and beach seines) used in the fisheries for pilchard and yellow-eyed mullet.

2. BIOLOGY

Only one species of garfish or piper is common in New Zealand waters, *Hyporhamphus ihi*. It is endemic, but very similar species occur in Australia. A larger garfish, *Euleptorhamphus viridis*, is occasionally recorded in northern New Zealand. The common garfish is not closely related to the ocean piper or saury, *Scomberexox saurus*. Garfish occur around most of New Zealand, and are present at the Chatham Islands. They are most abundant in sheltered gulfs, bays, and large estuaries, particularly near seagrass beds in shallow water, and over shallow reefs. The pale green, almost transparent colouring, and localised schooling behaviour of garfish makes them difficult to see and their abundance difficult to estimate.

Spawning occurs during spring and summer probably in suitable shallow bays; the eggs sink to the seafloor and adhere to vegetation. Larvae are seldom taken in coastal plankton surveys.

Patterns of age and growth are not known in New Zealand, but likely to be similar to Australia, where the larger of two closely related species (southern garfish, *H. melanochir*) matures at 25 cm (2–3 years) and reaches 52 cm (10 years). The New Zealand garfish matures at 22 cm, and with a maximum size of 40 cm may have a lower maximum age. Average size is 20–30 cm.

Garfish feed on zooplankton. They form single-species schools, but occur in close proximity with other small pelagic fishes in shallow coastal waters, particularly yellow-eyed mullet.

There have been no biological studies that are directly relevant to the recognition of separate stocks, or to yield estimates. Consequently no estimates of biological parameters are available.

3. STOCKS AND AREAS

There is no information on whether separate biological stocks occur in New Zealand. Given their preferred habitat of shallow sheltered waters, and the mode of reproduction in which the eggs are attached to the seafloor rather than free-floating, it is probable that localised populations occur, and possible that these may differ in some biological parameters (e.g., growth and recruitment). Consequently these populations may be susceptible to local depletion.

Garfish are sometimes taken as a non-target catch in the pilchard fishery, but this catch is likely to be very small. Although the target fisheries for these two species are quite separate, it is convenient for their Fishstocks to have the same boundaries.

4. STOCK ASSESSMENT

There have been no previous stock assessments of garfish.

4.1 Estimates of fishery parameters and abundance

No fishery parameters are available.

4.2 Biomass estimates

No estimates of biomass (B_0 , B_{MSY} , or $B_{current}$) are available.

4.3 Yield estimates and projections

MCY cannot be determined.

Current biomass cannot be estimated, so CAY cannot be determined.

4.4 Other yield estimates and stock assessment results

No information is available.

4.5 Other factors

The extent of natural variability in the size of garfish populations is not known, but from their very shallow inshore distribution, and demersal rather than pelagic eggs, it is suspected that they are less variable than other small pelagic species. However, these features also suggest localised populations, susceptible to local depletion.

There is anecdotal information that garfish are very abundant in some localities. It is not known whether this represents similar abundance over a larger region, or a tendency for a few schools to become concentrated in these localities. Apparent abundance, and initial catches, may be misleading in terms of sustainable yields.

The maximum age of 10 years proposed for a similar Australian garfish implies that productivity might not be as high as would be expected from a small pelagic species.

There is no reliable information on catches from the recreational fishery for garfish, or even their size relative to that of the commercial fishery.

5. STATUS OF THE STOCKS

No estimates of current biomass are available. A fishery has existed for several decades, but it is not known how heavily this has exploited the stock. It is not possible to determine if recent catch levels will allow the stock(s) to move towards a size that would support the MSY .

TACCs and reported landings by Fishstock are summarised in Table 4.

Table 4: Summary of yield estimates (t), TACCs (t), and reported landings (t) for garfish for the most recent fishing year.

Fishstock	QMA	FMAs	MCY estimates	2016–17 Actual	2016–17 Reported
				TACC	Landings
GAR 1	Auckland (East)	1	—	25	26
GAR 2	Central (East)	2	—	5	0
GAR 3	South East (Coast), Southland, Sub-Antarctic	3, 5, 6	—	5	0
GAR 4	South East (Chatham)	4	—	2	0
GAR 7	Challenger	7	—	8	0
GAR 8	Auckland (West), Central (West)	8, 9	—	5	0
GAR 10	Kermadec	10	—	0	0
Total			—	50	26

6. FOR FURTHER INFORMATION


Abel, K; Kailola, P (1993) Garfish. In Kailola, P J et al (comps). 1993. Australian fisheries resources. pp. 225–227. Bureau of Resource Sciences and the Fisheries Research and Development Corporation. Canberra. 422 p.

Collette, B B (1974) The garfishes (Hemiramphidae) of Australia and New Zealand. *Records of the Australian Museum* 29(2): 11–105.

Paul, L (2000) *New Zealand fishes. Identification, natural history & fisheries*. Reed Books, Auckland. 253 p.

GEMFISH (SKI)*(Rexea solandri)*

Maka-tikati

**1. FISHERY SUMMARY****1.1 Commercial fisheries**

Gemfish are caught in coastal waters around mainland New Zealand down to about 550 m. Historical estimated and recent reported gemfish landings and TACCs are shown in Tables 1 and 2, while Figure 1 shows the historical and recent landings and TACC values for the main gemfish stocks. Annual catches increased significantly in the early 1980s and peaked at about 8250 t in 1985–86 (Table 1). In the late 1980s, annual catches generally ranged from about 4200 to 4800 t per annum, but since then have steadily declined, with landings of less than 1000 t reported in most recent years (Table 3). TACCs were reduced in SKI 3 and SKI 7 for the 1996–97 fishing year and have been progressively reduced in SKI 1 and SKI 2 since 1997–98. TACs and TACCs are 218 t and 210 t for SKI 1, and 248 t and 240 t for SKI 2, respectively. Both SKI 1 and SKI 2 were allocated customary and recreational allowances of 3 t and 5 t respectively.

Table 1: Reported gemfish catch (t) from 1978–79 to 1987–88. Source - MAF and FSU data.

Fishing year Year	New Zealand		Foreign Licensed			Total
	Domestic	Chartered	Japan	Korea	USSR	
1978–79*	352	53	1 509	1 079	0	2 993
1979–80*	423	1 174	1 036	78	60	2 771
1980–81*	1 050	N/A	N/A	N/A	N/A	> 1 050
1981–82*	1 223	1 845	391	16	0	3 475
1982–83*	822	1 368	274	567	0	3 031
1983–83†	1 617	1 799	57	37	0	3 510
1983–84‡	1 982	3 532	819	305	0	6 638
1984–85‡	1 360	2 993	470	223	0	5 046
1985–86‡	1 696	4 056	2 059	442	0	8 253
1986–87‡	1 603	2 277	269	76	0	4 225 §
1987–88‡	1 016	2 331	90	35	0	3 472 §

* 1 April–31 March.

‡ 1 October–30 September.

† 1 April–30 September.

§ These totals do not match those in Table 2 due to under-reporting to the FSU.

N/A Unknown.

GEMFISH (SKI)

Table 2: Reported landings (t) for the main QMAs from 1931 to 1982.

Year	SKI 1	SKI 2	SKI 3	SKI 7	Year	SKI 1	SKI 2	SKI 3	SKI 7
1931–32	0	0	0	0	1957	2	12	21	10
1932–33	0	0	0	0	1958	5	34	19	28
1933–34	0	42	0	66	1959	2	40	58	38
1934–35	0	70	0	105	1960	3	61	65	39
1935–36	0	39	0	59	1961	6	42	14	19
1936–37	0	37	13	57	1962	5	58	49	27
1937–38	0	86	19	130	1963	19	72	19	38
1938–39	0	50	47	66	1964	17	48	20	29
1939–40	0	48	47	72	1965	19	96	11	28
1940–41	0	58	72	87	1966	12	102	15	26
1941–42	1	63	50	96	1967	32	173	14	46
1942–43	0	47	22	71	1968	18	183	15	33
1943–44	0	15	15	23	1969	60	308	11	22
1944	0	14	15	23	1970	50	281	22	28
1945	6	19	13	30	1971	52	315	24	59
1946	5	20	30	33	1972	85	261	15	37
1947	0	23	74	32	1973	56	237	46	102
1948	1	28	51	44	1974	21	150	14	89
1949	4	19	48	28	1975	2	96	172	37
1950	15	32	59	30	1976	11	108	8	36
1951	5	29	35	27	1977	22	118	4	74
1952	1	21	45	22	1978	36	235	411	1069
1953	1	13	42	10	1979	82	235	2104	628
1954	2	31	12	38	1980	278	287	1899	924
1955	0	25	22	23	1981	236	350	1369	1669
1956	0	31	27	35	1982	546	219	971	676

Notes:

1. The 1931–1943 years are April–March but from 1944 onwards are calendar years.
2. Data up to 1985 are from fishing returns: Data from 1986 to 1990 are from Quota Management Reports.
3. Data for the period 1931 to 1982 are based on reported landings by harbour and are likely to be underestimated as a result of under-reporting and discarding practices. Data includes both foreign and domestic landings.

Table 3: Reported landings (t) of gemfish by Fishstock from 1983–84 to present and actual TACs from 1986–87.

Fishstock FMA (s)	SKI 1 1 & 9		SKI 2 2		SKI 3 3, 4, 5, & 6		SKI 7 7 & 8		SKI 10 10	Total	
	Landings	TAC	Landings	TAC	Landings	TAC	Landings	TAC	TAC	Landings	TAC
1983–84*	588	-	632	-	3 481	-	1 741	-	† -	6 442 §	-
1984–85*	388	-	381	-	2 533	-	1 491	-	† -	4 793 §	-
1985–86*	716	-	381	-	5 446	-	1 468	-	† -	8 011 §	-
1986–87	773	550	896	860	2 045	2 840	1 069	1 490	†10	4 783	5 750
1987–88	696	632	1 095	954	1 664	2 852	1 073	1 543	†10	4 528	5 991
1988–89	1 023	1 139	1 011	1 179	1 126	2 922	1 083	1 577	†10	4 243	6 827
1989–90	1 230	1 152	1 043	1 188	1 164	3 259	932	1 609	†10	4 369	7 218
1990–91	1 058	1 152	949	1 188	616	3 339	325	1 653	†10	2 948	7 342
1991–92	1 017	1 152	1 208	1 197	287	3 339	584	1 653	†10	3 096	7 350
1992–93	1 292	1 152	1 020	1 230	371	3 345	469	1 663	†10	3 152	7 401
1993–94	1 156	1 152	1 058	1 300	75	3 345	321	1 663	†10	2 616	7 470
1994–95	1 032	1 152	905	1 300	160	3 355	103	1 663	†10	2 169	7 480
1995–96	801	1 152	789	1 300	49	3 355	81	1 663	†10	1 720	7 480
1996–97	965	1 152	978	1 300	58	1 500	238	900	†10	2 240	4 862
1997–98	627	752	671	849	27	300	44	300	†10	1 369	2 211
1998–99	413	460	336	520	17	300	59	300	†10	825	1 590
1999–00	409	460	506	520	62	300	107	300	†10	1 083	1 590
2000–01	335	460	330	520	47	300	87	300	†10	799	1 590
2001–02	201	210	268	240	72	300	123	300	†10	664	1 060
2002–03	206	210	313	240	115	300	268	300	†10	902	1 060
2003–04	221	210	301	240	78	300	542	300	†10	1 142	1 060
2004–05	234	210	259	240	72	300	635	300	†10	1 199	1 060
2005–06	230	210	182	240	27	300	248	300	†10	687	1 060
2006–07	215	210	317	240	26	300	209	300	†10	767	1 060
2007–08	216	210	249	240	18	300	179	300	†10	662	1 060
2008–09	191	210	191	240	11	300	213	300	†10	606	1 060
2009–10	247	210	176	240	20	300	144	300	†10	587	1 060
2010–11	226	210	300	240	33	300	301	300	†10	860	1 060
2011–12	212	210	155	240	11	300	260	300	†10	638	1 060
2012–13	182	210	140	240	23	300	234	300	†10	580	1 060
2013–14	198	210	268	240	39	300	268	300	†10	764	1 060
2014–15	83	210	168	240	21	300	231	300	†10	503	1 060
2015–16	188	210	224	240	80	300	186	300	†10	677	1 060
2016–17	244	210	236	240	248	300	431	300	10	1 159	1 060


FSU data.

§ The totals do not match those in Table 1 as some fish were not reported by area (FSU data prior to 1986–87).

† No recorded landings

Table 4: Catch history for gemfish stocks, divided into pre-spawning and spawning seasons (t). N/A - not available.

Year	SKI 1 (spawn)			SKI 2 (pre-spawn)	Total SKI 1 & 2	Year	SKI 1 (spawn)			SKI 2 (pre-spawn)	Total SKI 1 & 2
	SKI 1E	SKI 1W	Total				SKI 1E	SKI 1W	Total		
1952	5	0	5	50	55	1984	588	0	588	632	1 220
1953	5	0	5	25	30	1985	388	0	388	381	769
1954	5	0	5	60	65	1986	716	0	716	381	1 097
1955	5	0	5	35	40	1987	773	0	773	896	1 669
1956	5	0	5	35	40	1988	696	0	696	1 095	1 791
1957	5	0	5	55	60	1989	1 023	0	1 023	1 011	2 034
1958	5	0	5	30	35	1990	1 230	0	1 230	1 043	2 273
1959	5	0	5	45	50	1991	1 048	10	1 058	949	2 007
1960	5	0	5	85	90	1992	940	77	1 017	1 208	2 225
1961	5	0	5	70	75	1993	1 137	155	1 292	1 020	2 312
1962	5	0	5	60	65	1994	606	550	1 156	1 058	2 214
1963	15	0	15	70	85	1995	438	594	1 032	906	1 938
1964	15	0	15	65	80	1996	485	316	801	789	1 590
1965	20	0	20	130	150	1997	385	580	965	978	1 943
1966	15	0	15	140	155	1998	N/A	N/A	627	671	1 298
1967	35	0	35	240	275	1999	N/A	N/A	413	335	748
1968	40	0	40	250	290	2000	N/A	N/A	409	506	915
1969	100	0	100	375	475	2001	N/A	N/A	335	330	665
1970	95	0	95	400	495	2002	N/A	N/A	201	268	487
1971	100	0	100	420	520	2003	N/A	N/A	206	313	519
1972	130	0	130	400	530	2004	N/A	N/A	221	301	522
1973	45	0	45	300	345	2005	N/A	N/A	234	259	493
1974	35	0	35	230	265	2006	N/A	N/A	230	182	412
1975	10	0	10	170	180	2007	N/A	N/A	215	317	532
1976	30	0	30	190	220	2008	N/A	N/A	216	249	465
1978	90	0	90	240	330	2009	N/A	N/A	191	191	382
1979	120	0	120	200	320	2010	N/A	N/A	247	176	424
1980	140	0	140	450	590	2011	N/A	N/A	226	300	525
1981	120	0	120	500	620	2012	N/A	N/A	212	155	367
1982	100	0	100	320	420	2013	N/A	N/A	182	140	322
1983	360	0	360	730	1 090						

Figure 1: Reported commercial landings and TACC for the four main SKI stocks. SKI 1 (Auckland East).
[Continued on next page].

GEMFISH (SKI)




Figure 1 [Continued]: Reported commercial landings and TACC for the four main SKI stocks. From top to bottom right: SKI 2 (Central East), SKI 3 (South East Coast) and SKI 7 (Challenger).

Most of the recorded catch is taken by trawlers. Target fisheries developed off the eastern and northern coasts of the North Island. From 1993 to 2000 there was a major shift in effort from east of North Cape to the west (Table 4), and over 50% of the SKI 1 catch was taken from QMA 9 in some years. However, the distribution of fishing changed substantially after 2001 when the quota was last reduced. The west coast fishery has since virtually disappeared, as has the fishery off East Northland, each accounting for less than 10% of the SKI 1 catch since 2001–02. The Bay of Plenty fishery has

correspondingly increased, accounting for over 80% of the SKI 1 landings in the same period. While landings in SKI 1 are almost entirely concentrated in the months of May and June, landings in SKI 2 are spread fairly evenly from October to May. SKI 2 landings occur as a bycatch in a range of trawl fisheries, including tarakihi, barracouta, scampi and hoki, although over 80% of the SKI 2 landings are targeted at gemfish. Catches off the west and southern coasts of the South Island are primarily bycatch of hoki and squid target fisheries. Reported landings in SKI 7 were more than double the level of the TACC in 2004–05, but decreased in the following years until again exceeding the TACC in 2016–17. Landings in SKI 3 were at very low levels from about 1993–94 to 2015–16 until increasing in 2016–17. Figure 1 shows the historical landings and TACC values for the main SKI stocks.

1.2 Recreational fisheries

Little or no recreational catch was reported in marine recreational fishing telephone-diary surveys between 1992 and 2001 but the harvest estimates provided by these surveys are no longer considered reliable. A Recreational Technical Working Group concluded that these harvest estimates should be used only with the following qualifications: a) they may be very inaccurate; b) the 1996 and earlier surveys contain a methodological error; and c) the 2000 and 2001 estimates are implausibly high for many important fisheries. In response to these problems and the cost and scale challenges associated with onsite methods, a National Panel Survey was conducted for the first time throughout the 2011–12 fishing year. The panel survey used face-to-face interviews of a random sample of 30 390 New Zealand households to recruit a panel of fishers and non-fishers for a full year. The panel members were contacted regularly about their fishing activities and harvest information collected in standardised phone interviews. Harvest estimates for gemfish (in numbers of fish) are given in Table 5 (from Wynne-Jones et al 2014).

Table 5: Recreational harvest estimates for gemfish stocks (Wynne-Jones et al 2014). Mean fish weights were not available from boat ramp surveys so catches cannot be converted to weight.

Stock	Year	Method	Number of fish	Total weight (t)	CV
SKI 1	2011/12	Panel survey	2 752	-	0.39
SKI 7	2011/12	Panel survey	137	-	1.03
SKI total	2011/12	Panel survey	2 889	-	0.39

A repeat of the National Panel Survey is being conducted over the 2017–18 October fishing year. Results are expected in early 2019.

1.3 Customary non-commercial fisheries

Quantitative information on the current level of customary non-commercial take is not available and is assumed to be negligible.

1.4 Illegal catch

The amount of gemfish misreported is not available and is assumed to be negligible.

1.5 Other sources of mortality

There may have been some gemfish discarded prior to the introduction of the EEZ, but this is likely to have been minimal since the early 1980s as gemfish is a medium value species.

2. BIOLOGY

Gemfish occur on the continental shelf and slope, from about 50–550 m depth. They are known to undertake spawning migrations and the pre-spawning runs have formed the basis of winter target fisheries, but exact times and locations of spawning are not well known. Spawning probably takes place about July near North Cape and late August/September on the west coast of the South Island.

Ageing of southern gemfish indicate that fish attain about 30 cm at the end of the first year, 45 cm at the end of the second year, 53 cm at the end of the third year and 63 cm at the end of the fourth year.

GEMFISH (SKI)

Both sexes display similar growth rates until age 5, but subsequently, females grow larger. The maximum ages recorded for gemfish (from 1989 to 1994) are 17 years for both sexes. In the northern fishery (SKI 1, SKI 2), males and females appear to recruit into the fishery from age 3 but are probably not fully recruited until about age 5 (SKI 2) and age 7 or 8 (spawning fishery in SKI 1). In the southern fishery, gemfish start to recruit at age 2 into spawning and non-spawning fisheries but age at full recruitment is difficult to determine because of large variation in year class strength.

Recruitment variability in SKI 3 and SKI 7 has been correlated to wind and sea surface temperature patterns during the spawning season (Renwick et al 1998). No significant correlations were found between SKI 1 and SKI 2 recruitment indices and a range of climate variables (Hurst et al 1999).

Biological parameters relevant to the stock assessment are shown in Table 6.

Table 6: Estimates of biological parameters for gemfish.

Fishstock				Source			
1. Natural mortality (<i>M</i>)							
All stocks		<i>M</i> = 0.25 y ⁻¹ considered best estimate for all areas for both sexes		Horn & Hurst (1999)			
2. Weight = <i>a</i> (length) ^b (Weight in g, length in cm fork length)							
	Male		Female				
	<i>a</i>	<i>b</i>	<i>a</i>	<i>b</i>			
SKI 1	0.0034	3.22	0.0008	3.55	Langley et al (1993)		
SKI 3	0.0012	3.41	0.0095	3.47	Hurst & Bagley (1998)		
3. von Bertalanffy growth parameters							
	Male		Female				
	<i>L</i> _∞	<i>k</i>	<i>t</i> ₀	<i>L</i> _∞	<i>k</i>	<i>t</i> ₀	
East Northland	90.7	0.204	-0.49	122.7	0.114	-1.1	Langley et al (1993)
East Northland	88.4	0.235	-0.54	108.5	0.167	-0.71	Horn & Hurst (1999)
Wairarapa	90.8	0.287	0.00	103.4	0.231	-0.1	Horn & Hurst (1999)
West Northland	86.3	0.295	-0.11	103.4	0.209	-0.37	Horn & Hurst (1999)
North combined	87.4	0.266	-0.35	105	0.194	-0.55	Horn & Hurst (1999)
Southland	88.5	0.242	-0.66	104.2	0.178	-0.88	Horn & Hurst (1999)

3. STOCKS AND AREAS

In previous assessments, analysis of seasonal trends in gemfish fisheries indicated that there may be at least two stocks:

1. A southern/west coast stock (SKI 3 & 7), caught in the southern area in spring, summer and autumn, which presumably migrates to the west coast of the South Island to spawn and is caught there mainly in August–September. Spawning is thought to occur in late August/early September.
2. A northern/east coast stock (SKI 1E & SKI 2), caught mainly on the east coast in spring and summer, which migrates in May–June to spawn north of the North Island. Seasonal trends in commercial catch data from SKI 1E (QMA 1) are consistent with pre- and post-spawning migrations through the area; similar data from SKI 2 are inconclusive but indicate lower catches during the peak spawning months, although this could be partly due to target fishing on other species, particularly orange roughy, at this time.

The relationship of the pre-spawning fishery in SKI 1W (QMA 9) to the pre-spawning fishery in SKI 1E was investigated by Horn & Hurst (1999). They presented age frequency distributions from commercial catches for SKI 1E, SKI 1W, SKI 2 and from research sampling for SKI 3. Age distributions for the two SKI 1 spawning fisheries appear similar, with year classes in 1980, 1982, 1984, 1986 and 1991 appearing to be strong relative to other year classes. The SKI 2 distribution also exhibits the same pattern, although the relative dominance of the 1991 year class is greater, as might be expected from an area in which pre-recruit fish occur. The age distribution from SKI 3 gemfish showed that the 1982, 1984, 1985 and 1989 year classes were the stronger ones. There were no significant differences in the von Bertalanffy growth parameters calculated for northern and southern gemfish (Horn & Hurst 1999).

Recent biochemical analyses of Australasian gemfish suggested that there may be a very low level of mixing between eastern Australian and New Zealand gemfish, but not high enough to treat them as a single stock. There was also a suggestion of a difference between north-eastern and southern New Zealand gemfish.

Two alternative hypotheses have been proposed; that both SKI 1 and SKI 2 are one stock, or SKI 1W is separate from SKI 1E and SKI 2. The Middle Depths Working Group concluded that based on the close similarity in declines in CPUE indices and in age distributions from commercial catches that the northern gemfish should be assessed using SKI 1 and 2 combined.

4. STOCK ASSESSMENT

The assessment for the SKI 1 and SKI 2 stock was updated in 2007 with new standardised CPUE indices and addition of catch-at-age data up to 2005–06. Further analysis was carried out in 2008 incorporating SKI 2 catch-at-age for 2006–07. A number of changes were made to the 2003 model including the use of age-based selectivities and differential natural mortality.

The northern gemfish stock was assessed using the hypothesis of one stock (SKI 1 and SKI 2). The alternative hypothesis, that SKI 1W is separate from SKI 1E and SKI 2 was not modelled, as results from previous assessments were similar to those from SKI 1 and SKI 2 combined. Estimates of virgin biomass (B_0) and current mature biomass are presented below.

The stock assessment model includes two fishery types, based on spawning activity. The first is on the home ground, SKI 2, where all age classes occur and where fishing is mainly in the non-spawning season. The second is on the spawning migrations, SKI 1, where only mature age classes occur and where fishing is in the winter months. The non-spawning (SKI 2) and spawning (SKI 1) season landings used in the assessment are given in Table 4. This table also shows the split between east and west coast catches in SKI 1 from 1991 to 1997. The stock assessment was implemented as a Bayesian single stock model using the general-purpose stock assessment program CASAL v2.20 (Bull et al 2008). The assessment used catch-per-unit-effort time series, catch-at-age from the commercial fishery, and estimates of biological parameters.

New information from the previous assessment included a revised catch history, new CPUE abundance indices, four years of catch sampling proportions-at-age data for SKI 2, and one year of catch sampling proportions-at-age data for SKI 1.

The assessment of the southern stock (SKI 3 & 7) was not updated, as there were no new indices of biomass or proportion at age available. The results of the 1997 assessment are summarised below.

4.1 Auckland (SKI 1) and Central East (SKI 2)

4.1.1 Age composition of commercial catches

Commercial catch-at-age data included in the models were: SKI 1E for 1989 to 1994, 1997 to 1999, 2002, and 2006; SKI 1W for 1996 to 1999, and 2002; and SKI 2 for 1996 to 2005, and 2007. Age data for SKI 1E and SKI 1W were combined for the stock assessment model.

4.1.2 Estimates of abundance

Standardised CPUE indices for SKI 1 and SKI 2 were calculated for three fishery sub-groups in 2007: (1) target catch only; (2) all gemfish catch; and (3) all gemfish catch on TCEPR forms (Figures 2 and 3). The indices for TCEPR all gemfish catch (SKI 1 for 1990 to 2006, SKI 2 for 1994 to 2006) were used in the assessment (Table 7). The indices for SKI 1 are from SKI 1E and SKI 1W combined and for SKI 2 include both midwater and bottom trawl methods. Both time series show steep declines to the early 2000s, followed by marked increases in recent years.

GEMFISH (SKI)

In 2007, the Working Group considered year*area interactions in the CPUE model. This model was used to overcome the difference in timing of catch rate declines in different statistical areas of SKI 1. The catch rate in each statistical area had a different scale but a similar trend. Weighting of data would require relative population sizes (by area) to do correctly.

The Working Group thought at the time (2007) that the CPUE series should stop in 2001 when the quota was last reduced. Since then the indices are unlikely to be proportional to abundance in the stock given the changes observed in the fishery. The distribution of fishing in SKI 1 has shrunk to a small area in the Bay of Plenty and no fishing occurred on the WCNI in the last three years. In SKI 2 many vessels have left the area or have stopped targeting gemfish, therefore the CPUE series from 1994 to 2001 only should be used. The Working Group agreed in 2007 to use the CPUE indices from each fishery in the stock assessment based on TCEPR data including all SKI catch (Table 7).

Table 7: Standardised catch per unit effort indices and coefficient of variation (CV) for SKI 1 and SKI 2. The SKI 2 model is the combined mixed target species model (including SKI), based on daily effort data.

Year	Index	SKI 1		SKI 2	
		Index	CV	Index	CV
1990	1.94	0.10	0.061	6.28	0.061
1991	1.71	0.12	0.056	3.18	0.056
1992	1.36	0.10	0.053	1.52	0.053
1993	1.48	0.07	0.052	1.65	0.052
1994	1.73	0.06	0.051	1.24	0.051
1995	1.65	0.07	0.053	1.25	0.053
1996	1.05	0.06	0.063	0.76	0.063
1997	1.20	0.06	0.067	0.51	0.067
1998	0.86	0.06	0.068	0.38	0.068
1999	0.68	0.07	0.071	0.55	0.071
2000	0.66	0.07	0.074	0.53	0.074
2001	0.56	0.08	0.070	0.54	0.070
2002	–	–	0.070	0.66	0.070
2003	–	–	0.062	0.84	0.062
2004	–	–	0.060	1.18	0.060
2005	–	–	0.065	0.62	0.065
2006	–	–	0.061	0.52	0.061
2007	–	–	0.057	0.98	0.057
2008	–	–	0.063	1.05	0.063
2009	–	–	0.060	0.86	0.060
2010	–	–	0.056	0.83	0.056
2011	–	–	0.052	1.74	0.052
2012	–	–	0.053	1.74	0.053
2013	–	–	0.060	1.15	0.060

4.1.3 2014 SKI 2 CPUE update

The SKI 2 CPUE series was updated in 2014 with data up to the end of 2012–13. The SKI 1 series was not updated because of the cessation of fishing in East Northland and SKI 1W. The SKI 2 CPUE series differed from the previous series in a number of ways: a) only bottom trawl was used; b) data from all form types were amalgamated into a day of fishing by a vessel, selecting the modal target species and modal statistical area when there were multiple values within a day; c) target species (including SKI) was included in the analysis as an explanatory variable. Sensitivity analyses included excluding target SKI records and repeating both analyses using only the event-level forms in their original tow-by-tow stratification. These data were used to prepare lognormal models based on positive catch records and binomial models based on the presence/absence of gemfish, which were subsequently combined into a single model using the delta-lognormal method. Gemfish landings from the scampi target fishery were analysed separately as another sensitivity, recognising that this fishery is quite different from the finfish fisheries used in the other analyses, using slower towing speeds and a very different type of net. These data were also analysed using two different data preparation methods: daily amalgamated data or original event-level (tow-by-tow) stratification.

These analyses appear to be extremely robust, with only small differences in the models that excluded or included SKI as a target category (Figure 3). There was also good correspondence with the 2007 CPUE series (even with the SKI 1 series), except at the beginning and the end of the series (Figure 4). The scampi target models were much more variable, given the much smaller data sets being used, but there was broad general agreement in the CPUE indices calculated from all three data sets.

The two daily amalgamated series show a precipitous drop in the first two years of data, followed by a long slow decline up to the end of the 1990s, when the fishery was severely curtailed (Table 3). Since then, there appears to have been gradual increase in relative CPUE, with current levels 3 to 4 times greater than the lowest value observed in 1998 (Table 7). The two tow-by-tow series show the same pattern as the daily effort series over the period of overlap, without the initial steep decline because there are insufficient tow-by-tow data in the years before 1994 (Figure 3).

4.1.3 Assessment model

The assessment model partitions the stock into two areas (spawning (SKI 1E and 1W) and home ground (SKI 2)), two sexes and age groups 1–20, with no plus group. There are four time steps in the model (Table 8). In the first time step, the 1 year-olds are recruited to the population, which is then subjected to fishing mortality in SKI 2. In the second time step, fish migrate into SKI 1, and again are subjected to fishing mortality. In time step 3, fish ages are incremented, and spawning occurs. Fish migrate back to SKI 2 in the final time step.




Figure 2: Standardised CPUE indices for the three fishery subgroups in SKI 1: “target catch”, black solid; “all catch”, black dotted; “TCEPR all catch”, gray solid. Vertical bars represent 95% confidence interval.




Figure 3: Comparison of the four main combined 2014 SKI 2 CPUE series: a) mixed target species model (including SKI) (daily effort data); b) mixed target species model (without SKI) (daily effort data); b) mixed target species model (including SKI) (tow-by-tow data); b) mixed target species model (without SKI) (tow-by-tow data).




Figure 4: Comparison of the 2014 combined SKI 2 mixed target species model (including SKI) (daily effort data) with three of the 2006 SKI 1&2 CPUE models: SKI 1 mixed target species, SKI 2 mixed target species, SKI 2 target SKI.

Table 8: Annual cycle of the stock model for gemfish, showing the processes taking place at each time step, their sequence within each time step, and the available observations. Fishing and natural mortality that occur within a time step occur after all other processes, with half of the natural mortality for that time step occurring before and half after the fishing mortality.

Step	Period	Processes	M	Observations	
				Description	% M
1	Oct–Apr	Fishing (SKI 2)	0.58	CPUE (SKI 2)	50
		Recruitment		Proportions at age (SKI 2)	50
2	May–Jun	Migration to SKI 1	0.17	CPUE (SKI 1)	50
		Fishing (SKI 1)		Proportions at age (SKI 1)	50
3	Jul	Spawning	0.08		
		Increment age			
4	Aug–Sep	Migration to SKI 2	0.17		

1. M is the proportion of natural mortality that was assumed to have occurred in that time step.

2. % M is the percentage of the natural mortality within each time step that was assumed to have taken place at the time each observation was made.

The model used separate male and female age-based maturation ogives for SKI 1 and fishing ogives for SKI 2. The SKI 2 fishery was truncated into an early (before 2001) and a late period (after 2002), and separate fishing ogives were used. The SKI 1 fishing ogives were assumed known and were fixed at 1 for all ages.

The age-based fishing ogives for SKI 2 were assumed to be logistic, with male estimated relative to female. The model used logistic migration ogives, one for each sex to determine the rates that fish will mature.

The natural mortality was parameterised by the average of male and female, with the difference estimated within the model. A constant average natural mortality of 0.25 y^{-1} was used. The differential natural mortality, in conjunction with sex-specific fishing ogives were used to account for the between sex difference in proportions at age.

Maximum exploitation rates for gemfish were assumed to be 0.5 for SKI 2 and 0.7 for SKI 1. The choice of the maximum exploitation rate has the effect of determining the minimum possible virgin

biomass allowed by the model. This value was set relatively high as there was little external information from which to determine this value.

Lognormal errors, with known CVs, were assumed for all relative biomass and proportions-at-age observations. The CVs available for the relative abundance and catch-at-age observations allow for sampling error only. However additional variance, assumed to arise from differences between model simplifications and real world variation, was added to the sampling variance. The additional variance, termed process error, was estimated in early runs of the model using all available data from MPD fits. Hence, the overall CV assumed in the initial model runs for each observation was calculated by adding process error and observation error. The process error added was a CV of 0.14 and 0.20 for the SKI 1 and SKI 2 CPUE series respectively, and 0.48, 0.40, and 0.14 for the SKI 1, SKI 2 early period, and SKI 2 late period proportions-at-age data (run 2006_{YCS2000}, see Table 10).

Year class strengths were assumed known (and equal to one) for years prior to 1978 and after 2000 (run 2006_{YCS2000}, see Table 10) when inadequate or no age data were available. Otherwise year class strengths were estimated under the assumption that the estimates from the model should average one.

The assumed prior distributions used in the assessment are given in Table 9. All priors were intended to be relatively uninformed, and were estimated with wide bounds.

Table 9: The assumed priors assumed for key distributions (when estimated). The parameters are mean (in natural space) and CV for lognormal.

Parameter description	Distribution	Parameters		Bounds	
		Mean	CV	Lower	Upper
B_0	uniform-log	-	-	2 500	250 000
SKI 1 CPUE q	uniform-log	-	-	1×10^{-7}	0.01
SKI 2 CPUE q	uniform-log	-	-	1×10^{-7}	0.01
YCS	lognormal	1	0.9	0.01	10.0
Selectivity	uniform	-	-	0.1	80.0
Maturation	uniform	-	-	1.3	10.0
Difference in M	uniform	-	-	0	0.5
Process error CV.	uniform	-	-	1×10^{-3}	2.0

Penalty functions were used to constrain the model so that any combination of parameters that did not allow the historical catch to be taken was strongly penalised.

MCMC chains were estimated using a burn-in length of 10^6 iterations, with every 10 000th sample taken from the next 10^7 iterations (i.e., a final sample of length 1000 was taken from the Bayesian posterior). Autocorrelations, and single chain convergence tests of Geweke (1992) and Heidelberger & Welch (1983) were applied to resulting chains to determine evidence of non-convergence (Smith 2001).

4.1.4 Results

Estimates of biomass were obtained using the biological parameters and model input described earlier. Three model runs were considered, as there were concerns that the recent SKI 2 catch-at-age samples could be biased due to possible changes in the fishery. Model run “2006_{YCS2000}” used data up to 2006 and estimated year class strengths from 1978 to 2000; run “2006_{YCS2001}” used the same data but estimated the year class strengths from 1978 to 2001; run “2007_{YCS2003}” incorporated data up to 2007, with year class strengths estimated from 1978 to 2003. Table 10 describes the three model runs.

Table 10: Model run labels and descriptions for the base case and sensitivity model runs.

Model run	Description
2006 _{YCS2000}	Fitting to catch-at-age up to 2006, and CPUE indices based on TCEPR to 2001, and estimating YCSs 1978–00, using an average natural mortality of 0.25 yr^{-1} and separate age-based logistic fishing selectivities for SKI 2 fisheries before and after 2001.
2006 _{YCS2001}	2006 _{YCS2000} , but estimated YCS from 1978–2001.
2007 _{YCS2003}	2006 _{YCS2000} , but included 2007 SKI 1 and 2 catch and 2007 SKI 2 catch-at-age, and estimated YCSs 1978–2003.

For each model run, MPD fits were obtained and qualitatively evaluated. MPD estimates of biomass trajectories are shown in Figure 5. MCMC estimates of the posterior median and 95% percentile credible intervals for current and virgin biomass are reported in Table 11, and for year class strengths are shown in Figure 6.

No evidence of lack of convergence from the MCMC chains was found in the estimates of B_0 , although some estimates of selectivity parameters showed evidence of lack of convergence.

The between-sex difference in natural mortality was estimated to have a median of 0.02, with a 95% credible interval between 0.01 and 0.03. The median natural mortality was estimated to be about 0.26 for males and 0.24 for female.

The spawning maturation ogives appeared to be poorly estimated; both male and female ogives had broad posterior density estimates. It appears that males were 50% mature at age 6, and females at 7–8 years.

The selectivity ogives for males and females taken by the SKI 2 commercial trawl fishery for the early period were very steep and the 3–4 year-olds had broad posterior density estimates, suggesting considerable uncertainty. The selectivity ogives for the recent period was also steep but had narrow bounds. There were marked differences in the ogives: about 80% and 65% of males were estimated to be fully selected relative to females for the early and recent fishery respectively. There is no information outside the model that allows the shape of the estimated ogives to be verified.

Year class strengths were poorly estimated before 1990 when the only data available to determine year class strength were from older fish (see Figure 6). The estimates suggest a period of generally higher than average recruitment during the 1980s, followed by a period of generally lower than average recruitment (1992–2000). For run 2006_{YCS2001}, the 2001 year class strength was estimated to be weak. For run 2007_{YCS2003}, recruitment appeared to have improved in 2002 and 2003, but was still below average, and the estimate of 2003 year class strength was very uncertain.

The stock declined markedly during the early 1980s, followed by a small period of recovery due to recruitment of strong year classes in the late 1980s. Since 1992, the stock declined to its lowest level due to increasing exploitation rates combined with a long period of low recruitment since the early 1990s (see Figure 6). For model runs including data up to 2006, the estimated posterior median of B_{2006} was at about 32% of B_0 when the 2001 year class strength was fixed at 1, or 26% of B_0 when this year class was being estimated. More pessimistic estimates of biomass were obtained when 2007 catch-at-data were included, which suggest that the posterior median of B_{2007} was at about 22% of B_0 (see Table 11).

Table 11: Bayesian median and 95% credible intervals of B_0 , $B_{current}$, and $B_{current}$ as a percentage of B_0 for the three model runs. $B_{current}$ refers to B_{2006} for run 2006_{YCS2000} and 2006_{YCS2001}, and B_{2007} for run 2007_{YCS2003};

Model run	B_0	$B_{current}$	$B_{current} (\%B_0)$
2006 _{YCS2000}	12 672 (11 398–14 709)	4 007(2 759–5 766)	32(24–40)
2006 _{YCS2001}	11 691 (10 636–13 283)	3 008(2 024–4 593)	26(19–35)
2007 _{YCS2003}	10 900 (9 853–12 403)	2 443(1 448–3 924)	22(15–32)

The effect of using a lower and higher value of natural mortality was investigated for run 2007_{YCS2003}: with the average M set at 0.20, the current biomass is about 16% B_0 ; with an average M set at 0.30, the current biomass is about 28% B_0 . Estimates of other model parameters were relatively insensitive to the assumed value of natural mortality.




Figure 5: MPD biomass trajectories for the three model runs: 2006YCS2000, 2006YCS2001, and 2007YCS2003.




Figure 6: Bayesian median of year class strength for the three model runs 2006YCS2000, 2006YCS2001, and 2007YCS2003. Dotted lines are the 95% credible intervals for run 2007YCS2003.

4.1.5 Discussion of model results

This assessment updated the 2003 assessment using a similar model structure, revised catch history, revised CPUE indices, and addition of catch-at-age data. The model used sex-specific fishing selectivities and differential natural mortality to account for the sex ratio bias in the data, and the SKI 2 fishery was split into an early and a recent period to account for a possible change in selectivity. Several model runs were carried out, in consideration of the uncertainty of the most recent recruitment, arising from the possible bias in the catch-at-age data in the last few years. Model estimates of the state of the northern gemfish stock show that the current biomass is about 32% of virgin level if recruitments since 2001 were assumed to be average, or 22% of virgin level if more recent recruitments were estimated using the additional catch-at-age data in 2007.

The CPUE indices were only used up to 2001, as the recent indices were considered to be unlikely to track abundance. The fits to the CPUE indices were reasonable, though the SKI 2 indices declined slightly more than those predicted by the model. There appears to be some inconsistency between SKI 1 and SKI 2 CPUE indices. Both show declining trends, but the SKI 2 indices decline faster for the first few years, and are relatively flat for the remainder of the time series.

The fits to the catch at age data were reasonable and diagnostics showed no great departure from the assumption of normality for all model runs. The models explained most of the between-sex difference for the early and recent SKI 2 catch at age. The main outliers were the SKI 2 female observations in 2005, and it is possible that a larger proportion of female fish have been selected by the trawl. There appear to be some structures in the residuals of the older age classes for the SKI 1 catch at age as there are very few observed 14 and 15+ year old fish from 1989 to 1994.

The additional year class strengths estimated for run 2007_{YCS2003} show improvement of recruitment since 2001, which appears to be corroborated by the increase in the abundance indices of the last five years. However, the representativeness of the more recent SKI 2 catch-at-age data needs to be further examined (few age 3 males were observed in 2005, but the 2002 year class was one of the dominant year classes at age five in the 2007 catch at age data). More reliable abundance indices for SKI 1 and 2 fisheries need to be developed in order to obtain better estimates of the recent recruitment.

4.1.6 Yield estimates and projections

MCY and CAY were determined using stochastic sample-based simulations. One simulation run is done for each sample from the posterior, ultimately producing an estimate of yield that has been averaged over all samples (Bull et al 2008). Each run extended over 150 years with recruitment randomly sampled, but with the first 100 of those years discarded to allow the population to stabilise. Yield calculation was based on the procedures of Francis (1992), where yields were maximised subject to the constraint that spawning stock biomass should not fall below 20% of B_0 more than 10% of the time. For all model runs, the current stock status was at or below the estimated B_{MAY} (Table 12).

Table 12: Yield estimates (MCY and CAY) and associated parameters for the three model runs where simulations were based on recruits resampled from the entire period in which year class strengths were estimated.

Model run	B_{MCY} (t)	B_{MCY} (% B_0)	MCY (t)	B_{MAY} (t)	B_{MAY} (% B_0)	MAY (t)	CAY (t)
2006 _{YCS2000}	6 698	53	995	4 117	32	1 404	1 305
2006 _{YCS2001}	6 304	54	865	3 934	34	1 270	925
2007 _{YCS2003}	5 928	48	816	3 676	34	1 194	755

4.1.7 Projections

The projections were estimated for five years under four scenarios (two alternative recruitment assumptions and two alternative catch levels). Recruitment was randomly resampled from the entire period in which the year class strengths were estimated, or only the recent period (e.g., 1992 to 2000 for run 2006_{YCS2000}, 1992 to 2001 for run 2006_{YCS2001}, and 1992 to 2003 for run 2007_{YCS2003}). Future catches were set equal to the current TACC or the estimated CAY (see Table 12).

For all model runs, projections with recruitment resampled from the longer period suggest that the stock is likely to increase when future catches are assumed to be the current TAC, and is likely to decrease slightly when future catches are assumed to be the estimated CAY; projections with recruitment resampled from the recent period suggest that the future biomass is likely to decrease under the TAC, and is likely to decrease quickly under the estimated CAY (Table 13).

Table 13: Bayesian median and 95% credible intervals of projected biomass B_{PROJ} , B_{PROJ} as a percentage of B_0 , and $B_{PROJ}/B_{CURRENT}(\%)$ for the three model runs where future catches were fixed at either TAC or estimated CAY, and future recruitments were randomly sampled from the long period or from the recent period. B_{PROJ} and $B_{CURRENT}$ refer to B_{2011} and B_{2006} for run 2006_{YCS2000} and 2006_{YCS2001}, and B_{2012} and B_{2006} for run 2007_{YCS2003};

Model run	Catch (t)	Recruitment	B_{PROJ}	$B_{PROJ} (\%B_0)$	$B_{PROJ}/B_{CURRENT} (\%)$
2006 _{YCS2000}	450	1978–2000	6 060 (3 242–12 075)	47 (27–92)	151 (94–264)
	450	1992–2000	3 815 (2 128–6 071)	30 (18–44)	98 (74–122)
	1 305	1978–2000	3 472 (595–8 535)	27 (5–65)	85 (17–200)
	1 305	1992–2000	1 195 (135–3 414)	9 (1–24)	31 (5–66)
2006 _{YCS2001}	450	1978–2001	4 263 (2 010–8 844)	36 (18–74)	140 (76–286)
	450	1992–2001	2 436 (1 257–4 136)	21 (11–32)	81 (57–107)
	1 305	1978–2001	2 809 (630–7 744)	23 (6–64)	91 (24–235)
	1 305	1992–2001	999 (100–2 863)	9 (1–22)	34 (5–68)
2007 _{YCS2003}	450	1978–2003	3 580 (1 531–6 990)	33 (15–62)	139 (82–280)
	450	1992–2003	2 361 (1 019–4 509)	21 (10–38)	96 (62–137)
	755	1978–2003	2 497 (692–6 200)	23 (7–54)	99 (36–233)
	755	1992–2003	1 476 (199–3 481)	14 (2–29)	59 (13–105)

The projections suggest that unless recruitment improves and the catch remains at moderately low levels, the biomass is unlikely to increase in the short term.

4.2 South-East/Southland (SKI 3) and Challenger/Central (West) (SKI 7)

4.2.1 Estimation of fishery parameters and abundance

Estimates of relative abundance from two time series of trawl surveys used in the model for SKI 3 are presented in Table 14. Proportion-at-age data included in the model came from the *Tangaroa* trawl surveys. Model input parameters used in the assessment are given in Table 15.

Table 14: Biomass indices (t) and coefficients of variation (CV) from trawl surveys (assuming area availability, vertical availability and vulnerability = 1).

Fishstock	Area	Vessel	Trip code	Date	Biomass	% CV
SKI 3	Southland	<i>Shinkai Maru</i>	SHI8102	Feb 1981	3 900	17
			SHI8201	Mar–Apr 1982	3 100	31
			SHI8303	Apr 1983	5 500	33
SKI 3	Southland	<i>Tangaroa</i>	TAN9301	Feb–Mar 1993	1 066	17
			TAN9402	Feb–Mar 1994	406	18
			TAN9502	Feb–Mar 1995	539	25
			TAN9604	Feb–Mar 1996	529	23

Table 15: MIAEL model input parameters used in the SKI 3 & 7 assessment.

Parameter	Estimate
Steepness	0.75
Recruitment variability	1.0
Proportion spawning	0.95
M	0.23
Maximum exploitation (r_{MAX}) pre-spawning, spawning	0.6, 0.8
Minimum exploitation with maximum catch (r_{MMX})	0.1
Maturity ogive (ages 2–5)	0.1, 0.4, 0.8 1.0




Figure 7: Bayesian median of projected biomass (% B_0) for the three model runs, with future catch fixed at TAC or estimated CAY, and future recruitment randomly resampled from the long period or the recent period.

Year class strength was estimated in the model. As some year classes were exceptionally weak or strong, constraints were set to give more realistic estimates of year class strengths. The estimated year class strengths are given in Table 16. These year class strengths were poorly estimated and should be considered as indicative of poor and strong year classes only.

Table 16: Estimated or assumed (*) year class strengths for the base case SKI 3 & 7 assessment.

Year class	Estimate	Year class	Estimate	Year class	Estimate
1979	3.310	1986	0.300	1993	0.010*
1980	1.940	1987	0.001	1994	0.010*
1981	0.001	1988	0.010		
1982	5.690	1989	0.240		
1983	0.070	1990	0.010		
1984	4.250	1991	0.001*		
1985	2.250	1992	0.001*		

4.2.2 Biomass estimates

There was concern over the MIAEL point estimates due to the low value of the performance indices and therefore only the upper and lower bounds using r_{MMX} and r_{MAX} were reported. B_0 ranged from 26 000 to 73 000 t, B_{MID97} from 0 to 63%, and B_{BEG98} from 200 to 51 400 t (see also figure 1 in the 1997 Plenary Report).

4.2.3 Yield estimates and projections

Details of the modelling procedure which produced the B_0 estimates from which MCY was estimated for SKI 3 & 7 are given above. The MCY ranges from 990 to 2 770 t. MIAEL point estimates were not reported due to the low value of the performance indices.

Details of the modelling procedure which produced the B_{beg98} estimates from which CAY was estimated for SKI 3 & 7 are given above. The range of CAY for SKI 3 & 7 for 1998–99 was 20–5 900 t. MIAEL point estimates were not reported due to the low value of the performance indices.

5. STATUS OF THE STOCKS

Stock Structure Assumptions

Gemfish are assessed as two biological stocks, based on spawning migration and timing and the location of spawning grounds. These stocks are managed and assessed separately and are assumed to be non-mixing. The SKI 1&2 stock is based on the east coast North Island, migrating north to spawn north of the North Island during May–June. The SKI 3&7 stock occurs in the south of New Zealand and migrates to the west coast South Island to spawn in August–September.


A new stock assessment was completed for SKI 1 & 2 in 2008 and the CPUE from the SKI 2 fishery was updated in 2014.

SKI 1&2


Stock Status	
Year of Most Recent Assessment	2008: Stock Assessment 2014: CPUE update
Assessment Runs Presented	<u>Stock Assessment</u> Three cases are presented. There was no single preferred model. <u>CPUE Update</u> Combined (lognormal + binomial) model based on mixed target species (including SKI) using daily effort data for Statistical Areas 011–019
Reference Points	Management Target: 40% B_0 Soft Limit: 20% B_0 Hard Limit: 10% B_0 Overfishing threshold: -
Status in relation to Target	B_{2006} was estimated at 32% B_0 (2006 _{YCS2000}) and 26% B_0 (2006 _{YCS2001}), and B_{2007} at 22% B_0 (2007 _{YCS2003}) in the three models Unlikely (< 40%) to be at or above the target in 2006 The 2014 CPUE analysis indicates that relative abundance increased by 119% from the mean for 2005–2007 to the mean for

	2011–2013. Although biomass is increasing, it is not known whether the stock has reached the target
Status in relation to Limits	B_{2006} was estimated to be Unlikely (< 40%) to be below both the Soft Limit and the Hard Limit
Status in relation to Overfishing	-

Historical Stock Status Trajectory and Current Status



MPD biomass trajectories for the three model runs: 2006YCS2000, 2006YCS2001, and 2007YCS2003.



Historical CPUE Trajectory with combined SKI 1&2 landings and TACC (t)

Fishery and Stock Trends	
Recent Trend in Biomass or Proxy	Standardised CPUE has increased steadily since the late 1990s.
Recent Trend in Fishing Mortality or Proxy	Fishing pressure has declined with the decrease in TACC since 1999–2000.
Other Abundance Indices	-
Trends in Other Relevant Indicators or Variables	One strong year class was estimated to have occurred in 1991. Recruitment in recent years appears lower than seen previously.

Projections and Prognosis		
Stock Projections or Prognosis	With catches at the current TACC the stock is projected to increase if recruitment returns to the 1978–2000 average level, but decline slightly if recent (1992–2000) recruitment continues.	
Probability of Current Catch or TACC causing Biomass to remain below or to decline below Limits	Soft Limit: Very Unlikely (< 10%) Hard Limit: Very Unlikely (< 10%)	
Probability of Current Catch or TACC causing Overfishing to continue or to commence	-	
Assessment Methodology and Evaluation		
Assessment Type	Type 1 – Quantitative Stock Assessment (to 2006) Type 2 – Partial Quantitative Stock Assessment (2014)	
Assessment Method	Age-structured CASAL model with Bayesian estimation of posterior distributions	
Assessment Dates	Latest assessment: 2007 CPUE update: 2014	Next assessment: Unknown
Overall assessment quality rank	-	
Main data inputs (rank)	<u>Stock Assessment</u> Updated from previous assessment: <ul style="list-style-type: none">- Catch history- CPUE abundance indices- Proportions-at-age data (1 year SKI 1, 4 years SKI 2) <u>CPUE Analysis</u> MPI catch and effort data	
Data not used (rank)	-	
Changes to Model Structure and Assumptions	Incorporation of: <ul style="list-style-type: none">- Age based selectivities- Differential natural mortality- Additional year of age data	
Major Sources of Uncertainty	<u>Stock Assessment</u> Uncertainty in recent recruitment necessitated the development of multiple models, however, without more reliable abundance indices to estimate recent recruitment it is unwise to prefer a single model. <u>CPUE</u> Steep decline in first two years of series and sustained high catches suggest the first two data points may not reliably reflect abundance.	

Qualifying Comments
-

Fishery Interactions

Gemfish are common bycatch in the hoki, tarakihi, scampi and squid target fisheries, although some gemfish target fisheries do exist. Bycatch is variable but includes hoki, tarakihi, silver warehou and bluenose. Interactions with other species are currently being characterised.

SKI 3 & 7


The assessment of the southern gemfish stock has not been updated since 1997. Landings from SKI 7 increased from 2000 to be a level over twice the TACC in 2004–05, but have decreased since then.

Table 17: Summary of yields (t) from base case assessments, TACCs (t) and reported landings (t) for gemfish for the most recent fishing year.

Fishstock	QMA	FMA	MCY	CAY	2016–17 Actual TACC	2016–17 Reported landings
SKI 1	Auckland (East) (West)	1 & 9 }			210	244
SKI 2	Central (East)	2 }	816	-	240	236
SKI 3	South-East (Coast) (Chatham), Southland, Sub-Antarctic	3, 4, 5, & 6 }			300	248
SKI 7	Challenger, Central (West)	7 & 8 }	990–2 770	-	300	431
SKI 10	Kermadec	10	-	-	10	0
Total					1 060	1 159

6. FOR FURTHER INFORMATION

- Ayers, S (2003) Standardised CPUE analysis for the northern gemfish (*Rexea solandri*) fisheries in SKI 1 and SKI 2, 1989–2002. *New Zealand Fisheries Assessment Report 2003/34*. 17 p.
- Bull, B; Francis, R I C C; Dunn, A; McKenzie, A; Gilbert, D J; Smith, M H; Bian, R (2008) CASAL (C++ algorithmic stock assessment laboratory) CASAL user manual v2.20-2008/02/14. *NIWA Technical Report 130*. 275 p.
- Cordue, P L (1995) MIAEL estimation of biomass and fishery indicators for the 1995 assessment of hoki stocks. New Zealand Fisheries Assessment Research Document 1995/13. 48 p. (Unpublished document held by NIWA library, Wellington.)
- Cordue, P L (1998) Designing optimal estimators for fish stock assessment. *Canadian Journal of Fisheries and Aquatic Sciences* 55(2): 376–386.
- Dunn, A; Hurst, R J; Phillips, N L (2001) Stock assessment of northern gemfish (*Rexea solandri*) in SKI 1 and SKI 2 for the 2000–01 fishing year. *New Zealand Fisheries Assessment Report 2001/43*. 47p.
- Francis, R I C C (1992) Recommendations concerning the calculation of Maximum Constant Yield (MCY) and Current Annual Yield (CAY). New Zealand Fisheries Assessment Research Document 1992/8. 23 p. (Unpublished document held by NIWA library, Wellington.)
- Fu, D; Dunn, A; Hurst, R J (2008) Standardised CPUE analysis and stock assessment of northern gemfish (*Rexea solandri*) in SKI 1 and 2 for the 2005–06 fishing year. *New Zealand Fisheries Assessment Report 2008/1*. 68 p.
- Geweke, J (1992) Evaluating the accuracy of sampled-based approach to calculating posterior moments. In: Bayesian statistics. Bernardo, J M; Berger, J O; David, A P; Smith, A F M (Eds). pp 169–194. Clarendon Press, Oxford.
- Heidelberger, P; Welch, P (1983) Simulation run length control in the presence of an initial transient. *Operations Research* 31: 1109–1144.
- Hicks, A C; Cordue, P; Bull, B (2002) Estimating proportion at age and sex in the commercial catch of hoki (*Macruronus novaezelandiae*) using length frequency data. *New Zealand Fisheries Assessment Report 2002/43*. 51 p.
- Horn, P L; Hurst, R J (1999) Stock structure and age of gemfish (*Rexea solandri*) in New Zealand waters. *Marine and Freshwater Research* 50: 103–115.
- Hurst, R J; Bagley, N W (1998) A summary of the biology and commercial landings, and a stock assessment of southern (SKI 3 and SKI 7) gemfish *Rexea solandri* (Gempylidae) in New Zealand waters. New Zealand Fisheries Assessment Research Document 1998/3. 51p. (Unpublished document held by NIWA library, Wellington.)
- Hurst, R J; Coburn, R P; Bull, B (1999) Final Research Report to the Ministry of Fisheries Project SKI9801 (Objectives 2, 3, & 4). 35 p. (Unpublished document held by Fisheries New Zealand, Wellington.)
- Hurst, R J; Coburn, R P; Horn, P L (2000) Assessment of northern gemfish stocks (SKI 1 and SKI 2) for 2000. *New Zealand Fisheries Assessment Report 2000/18*. 41 p.
- Langley, A; Hartill, B; Walshe, C (1993) Summary of the northern gemfish (SKI 1) trawl fishery, 1989–92. Northern Fisheries Region Internal Report 14. 39 p. (Unpublished report held by NIWA library, Wellington.)
- Phillips, N L; Horn, P L (2003) Length and age composition for the Northern Gemfish fisheries (SKI 1 & 2). Final Research Report for Ministry of Fisheries Research Project SKI2002/01, Objective 4 & 5. 14 p. (Unpublished report held by Fisheries New Zealand, Wellington.)
- Renwick, J A; Hurst, R J; Kidson, J W (1998) Climatic influences on the recruitment of southern gemfish (*Rexea solandri*, Gempylidae) in New Zealand waters. *International Journal of Climatology* 18(15): 1655–1667.
- Smith, B J (2001) Bayesian output analysis program. Version 1.00 user's manual. 45 p. University of Iowa College of Public Health. <http://www.public-health.uiowa.edu/boa> (Unpublished report).
- Wynne-Jones, J; Gray, A; Hill, L; Heinemann, A (2014) National Panel Survey of Marine Recreational Fishers 2011–12: Harvest Estimates. *New Zealand Fisheries Assessment Report 2014/67*.

DARK GHOST SHARK (GSH)*(Hydrolagus novaezealandiae)***1. FISHERY SUMMARY****1.1 Commercial fisheries**

Two species (dark and pale ghost sharks) make up effectively all commercial ghost shark landings. Dark ghost shark (*Hydrolagus novaezealandiae*) was introduced into the QMS from the beginning of the 1998–99 fishing year for the 10 FMAs shown above.

Both ghost shark species are taken almost exclusively as a bycatch of other target trawl fisheries. In the 1990s, about 43% of ghost sharks were landed as a bycatch of the hoki fishery, with fisheries for silver warehou, arrow squid and barracouta combining to land a further 36%. The two ghost shark species were seldom differentiated on catch landing returns prior to the start of the 1998–99 fishing year. Estimated landings of both species by foreign licensed and joint venture vessels over the period 1 April 1978 to 30 September 1983 are presented in Table 1. Landings by domestic (inshore) vessels would have been negligible during this time period. The unknown quantities of ghost sharks that were discarded and not recorded will have resulted in an under-reported total, particularly before both species were included in the QMS.

In the early to mid-1980s about half of the reported ghost shark landings were from FMA 3. Virtually all the additional catch was spread over FMAs 4–7. In 1988–89, landings from west coast South Island (FMA 7) began to increase, almost certainly associated with the development of the hoki fishery. In 1990–91, significant landing increases were apparent on the Chatham Rise, off southeast South Island and on the Campbell Plateau. The development of fisheries for non-spawning hoki were probably responsible for these increases.

Estimated landings of dark ghost shark by QMA are shown in Table 2, while the historical landings and TACC for the main GSH stocks are depicted in Figure 1. Landings from 1983–84 to 1994–95 were derived by splitting all reported ghost shark landings into depth and area bins, and allocating to species based on distribution data derived from trawl surveys (*see* section 2). Landings from 1995–96 to 1998–99 were estimated assuming dark ghost shark made up 70% of the total ghost shark catch in FMAs 5 and 6, and 75% in all other FMAs. However this approach assumes that the proportion that each species contributes to the whole is consistent from year to year and does not change in response to various sources of mortality, fishing-induced or otherwise. As such, the data covered by this period of time should be treated with caution. Catches from the 1999–00 fishing year are more reliable, when pale ghost shark had also been included in the QMS, bringing both under the system.

DARK GHOST SHARK (GSH)

Table 1: Reported landings (t) of both ghost shark species by fishing year and EEZ area, taken by foreign licensed and joint venture vessels. An approximation of these areas with respect to current QMA boundaries is used to assign catches to QMAs. No data are available for the 1980–81 fishing year.

Year	QMA	EEZ Area												Total
		B	C(M)	C(1)	D	E(B)	E(P)	E(C)	E(A)	F(E)	F(W)	G	H	
78–79*		1	37	99	26	3	16	11	88	90	8	68	17	465
79–80*		1	55	54	426	10	4	28	138	183	7	1	5	912
80–81*														-
81–82*		0	84	28	117	0	2	6	29	71	9	4	0	350
82–83*		0	108	35	84	0	2	17	98	99	29	1	1	474
83–83#		0	84	41	73	0	0	17	5	16	17	0	0	253

* 1 April to 31 March

1 April to 30 Sept.

Table 2: Reported landings (t) for the main QMAs from 1931 to 1982.

Year	GSH 1	GSH 2	GSH 3	GSH 4	Year	GSH 1	GSH 2	GSH 3	GSH 4
1931–32	0	0	0	0	1957	0	0	0	0
1932–33	0	0	0	0	1958	0	0	0	0
1933–34	0	0	0	0	1959	0	0	0	0
1934–35	0	0	0	0	1960	0	0	0	0
1935–36	0	0	0	0	1961	0	0	0	0
1936–37	0	0	0	0	1962	0	0	0	0
1937–38	0	0	0	0	1963	0	0	0	0
1938–39	0	0	0	0	1964	0	0	0	0
1939–40	0	0	0	0	1965	0	0	0	0
1940–41	0	0	0	0	1966	0	0	0	0
1941–42	0	0	0	0	1967	0	0	0	0
1942–43	0	0	0	0	1968	0	0	0	0
1943–44	0	0	0	0	1969	0	0	0	0
1944	0	0	0	0	1970	0	0	0	0
1945	0	0	0	0	1971	0	0	0	0
1946	0	0	0	0	1972	0	0	103	0
1947	0	0	0	0	1973	0	0	0	0
1948	0	0	0	0	1974	0	0	7	0
1949	0	0	0	0	1975	0	0	8	0
1950	0	0	0	0	1976	0	0	19	0
1951	0	0	0	0	1977	0	0	2	0
1952	0	0	0	0	1978	0	0	54	0
1953	0	0	0	0	1979	0	2	486	383
1954	0	0	0	0	1980	0	0	150	230
1955	0	0	0	0	1981	0	0	233	243
1956	0	0	0	0	1982	0	0	320	97

Year	GSH 5	GSH 6	GSH 7	GSH 8	Year	GSH 5	GSH 6	GSH 7	GSH 8
1931–32	0	0	0	0	1957	0	0	0	0
1932–33	0	0	0	0	1958	0	0	0	0
1933–34	0	0	0	0	1959	0	0	0	0
1934–35	0	0	0	0	1960	0	0	0	0
1935–36	0	0	0	0	1961	0	0	0	0
1936–37	0	0	0	0	1962	0	0	0	0
1937–38	0	0	0	0	1963	0	0	0	0
1938–39	0	0	0	0	1964	0	0	0	0
1939–40	0	0	0	0	1965	0	0	0	0
1940–41	0	0	0	0	1966	0	0	0	0
1941–42	0	0	0	0	1967	0	0	0	0
1942–43	0	0	0	0	1968	0	0	0	0
1943–44	0	0	0	0	1969	0	0	0	0
1944	0	0	0	0	1970	0	0	0	0
1945	0	0	0	0	1971	0	0	0	0
1946	0	0	0	0	1972	11	0	0	0
1947	0	0	0	0	1973	0	0	0	0
1948	0	0	0	0	1974	1	0	0	0
1949	0	0	0	0	1975	1	0	0	0
1950	0	0	0	0	1976	2	0	0	1
1951	0	0	0	0	1977	0	0	0	0
1952	0	0	0	0	1978	100	30	15	2
1953	0	0	0	0	1979	178	131	268	2
1954	0	0	0	0	1980	92	144	144	28
1955	0	0	0	0	1981	111	35	17	17
1956	0	0	0	0	1982	223	29	11	7

Notes:

The 1931–1943 years are April–March but from 1944 onwards are calendar years. Data up to 1985 are from fishing returns: Data from 1986 to 1990 are from Quota Management Reports. Data for the period 1931 to 1982 are based on reported landings by harbour and are likely to be underestimated as a result of under-reporting and discarding practices. Data includes both foreign and domestic landings. Data were aggregated to FMA using methods and assumptions described by Francis & Paul (2013).

Table 3: Estimated landings (t) of dark ghost shark by Fishstock from 1982–83 to 2016–17, based on reported landings of both ghost shark species combined, and actual TACCs set from 1998–99. No landings have been recorded from FMA 10, and no TACC has been set for this area. QMS data from 1986 to present.

Fishstock FMA (s)	GSH 1		GSH 2		GSH 3		GSH 4		GSH 5	
	1		2		3		4		5	
	Landings	TAC	Landings	TAC	Landings	TAC	Landings	TAC	Landings	TAC
1982–83*	1	-	< 1	-	151	-	65	-	35	-
1983–84*	0	-	< 1	-	185	-	65	-	42	-
1984–85*	< 1	-	4	-	136	-	95	-	50	-
1985–86*	< 1	-	1	-	276	-	60	-	30	-
1986–87	3	-	13	-	472	-	97	-	34	-
1987–88	4	-	< 1	-	539	-	53	-	49	-
1988–89	9	-	27	-	460	-	21	-	67	-
1989–90	1	-	14	-	383	-	29	-	78	-
1990–91	1	-	40	-	665	-	271	-	70	-
1991–92	4	-	7	-	444	-	179	-	81	-
1992–93	8	-	5	-	399	-	151	-	76	-
1993–94	7	-	7	-	569	-	144	-	51	-
1994–95	3	-	2	-	737	-	187	-	63	-
1995–96	13	-	37	-	678	-	253	-	71	-
1996–97	17	-	66	-	817	-	402	-	94	-
1997–98	17	-	17	-	767	-	262	-	70	-
1998–99	18	15	60	37	950	1 187	318	373	64	109
1999–00	15	15	51	37	938	1 187	173	373	71	109
2000–01	15	10	50	33	1 111	1 185	179	370	85	109
2001–02	22	10	52	33	1 068	1 185	241	370	76	109
2002–03	17	10	58	33	1 371	1 185	265	370	93	109
2003–04	21	10	84	33	894	1 185	157	370	45	109
2004–05	14	10	74	33	880	1 185	282	370	80	109
2005–06	20	10	57	33	583	1 185	318	370	61	109
2006–07	20	22	60	66	654	1 185	396	370	115	109
2007–08	19	22	100	66	484	1 185	562	370	67	109
2008–09	14	22	71	66	490	1 185	251	370	61	109
2009–10	13	22	64	66	520	1 185	233	370	108	109
2010–11	17	22	95	66	640	1 185	311	370	73	109
2011–12	11	22	57	66	497	1 185	482	370	72	109
2012–13	12	22	51	66	420	1 185	210	370	111	109
2013–14	15	22	83	89	667	1 185	201	370	53	109
2014–15	16	22	44	89	406	1 185	217	370	42	109
2015–16	21	22	38	89	547	1 185	217	370	56	109
2016–17	21	22	47	89	493	1 185	223	370	83	109

Fishstock FMA (s)	GSH 6		GSH 7		GSH 8		GSH 9		Total	
	6		7		8		9			
	Landings	TAC	Landings	TAC	Landings	TAC	Landings	TAC	Landings	TAC
1982–83*	19	-	10	-	< 1	-	0	-	282	-
1983–84*	56	-	38	-	< 1	-	0	-	387	-
1984–85*	61	-	63	-	< 1	-	0	-	409	-
1985–86*	41	-	31	-	3	-	0	-	442	-
1986–87	36	-	71	-	4	-	0	-	729	-
1987–88	6	-	68	-	1	-	0	-	720	-
1988–89	6	-	133	-	2	-	0	-	725	-
1989–90	9	-	180	-	27	-	0	-	722	-
1990–91	94	-	217	-	3	-	0	-	1 361	-
1991–92	80	-	124	-	3	-	1	-	923	-
1992–93	68	-	221	-	11	-	0	-	938	-
1993–94	53	-	513	-	14	-	0	-	1 357	-
1994–95	61	-	703	-	3	-	0	-	1 778	-
1995–96	68	-	548	-	8	-	3	-	1 679	-
1996–97	135	-	926	-	9	-	11	-	2 477	-
1997–98	136	-	170	-	3	-	12	-	1 454	-
1998–99	110	95	409	1 121	7	12	22	14	1 958	2 963
1999–00	117	95	466	1 121	19	12	25	14	1 875	2 963
2000–01	76	95	475	1 121	22	12	31	8	2 043	2 943
2001–02	94	95	463	1 121	22	12	25	8	2 063	2 943
2002–03	99	95	593	1 121	15	12	20	8	2 531	2 943
2003–04	72	95	652	1 121	27	12	12	8	1 964	2 943
2004–05	53	95	694	1 121	31	12	10	8	2 118	2 943
2005–06	31	95	625	1 121	22	12	8	8	1 725	2 943
2006–07	43	95	696	1 121	16	22	6	22	2 006	3 012
2007–08	36	95	601	1 121	29	22	13	22	1 911	3 012
2008–09	49	95	991	1 121	24	22	16	22	1 967	3 012
2009–10	19	95	1 037	1 121	29	22	6	22	2 028	3 012
2010–11	38	95	1 129	1 121	33	22	6	22	2 341	3 012
2011–12	37	95	1 041	1 121	37	22	6	22	2 240	3 012
2012–13	70	95	767	1 121	32	22	10	22	1 683	3 012
2013–14	72	95	691	1 121	27	34	9	22	1 817	3 047
2014–15	72	95	458	1 121	20	34	7	22	1 283	3 047
2015–16	64	95	400	1 121	19	34	6	22	1 368	3 047
2016–17	59	95	423	1 121	19	34	14	22	1 382	3 047

* FSU data.

DARK GHOST SHARK (GSH)




Figure 1: Reported commercial landings and TACC for GSH stocks. From top GSH 2 (Central East), GSH 3 (South East Coast), GSH 4 (South East Chatham Rise).




Figure 1: Reported commercial landings and TACC for GSH stocks. From top GSH 5 (Southland), GSH 6 (Sub-Antarctic), and GSH 7 (West Coast South Island).

The TACs currently applied to dark ghost shark were initially intended to apply to a combined fishery for both species, and were based on the average catch of both species over various periods (see the “Review of Sustainability Measures and Other Management Controls for the 1998–99 Fishing Year - Final Advice Paper” dated 6 August 1998). No allowance for non-commercial interests was included in the final allocation because recreational and customary non-commercial catches are likely to be very small due to the depth distribution of this species.

DARK GHOST SHARK (GSH)

TACCs were increased from 1 October 2006 in GSH 1 to 22 t, in GSH 2 to 66 t, in GSH 8 to 22 t and in GSH 9 to 22 t. In these stocks landings were above the TACC for a number of years and the TACCs have been increased to the average of the previous 7 years plus an additional 10%. Landings exceeded the TACC slightly in GSH 3 in 2002–03, slightly in GSH 4 in 2006–07 and by 52% in 2007–08. Landings also exceeded the TACC slightly in GSH 5 in 2006–07, and GSH 6 in 1999–00 and 2002–03.

1.2 Recreational fisheries

Current catches of dark ghost sharks by recreational fishers are believed to be negligible in all areas.

1.3 Customary non-commercial fisheries

Quantitative information on the current level of customary non-commercial catch is not available but is likely to be negligible

1.4 Illegal catch

Quantitative information on the level of illegal catch is not available. In 1998–99 (when dark ghost shark were in the QMS, but pale ghost shark were not), a quantity of dark ghost shark were reported as pale ghost shark.

1.5 Other sources of mortality

Ghost sharks have been dumped and not reported in the past by commercial fishers in QMAs 1 and 2. Similar behaviour is believed to occur in all other QMAs. The extent of the unreported dumping is unknown in all areas.

2. BIOLOGY

Dark ghost shark (*Hydrolagus novaezelandiae*) occur through much of the New Zealand EEZ in depths from 30 to 850 m, but they are sparse north of 40° S and have not been recorded from the Bounty Platform. They are most abundant in waters 150–500 m deep on the west coast of the South Island and the Chatham Rise, and in depths of 150–700 m on the Stewart-Snares shelf and Southland/sub-Antarctic. Smaller sharks (under 40 cm chimaera length) are more abundant in waters shallower than 200 m, particularly in the Canterbury Bight.

Trawl surveys show that dark and pale ghost shark exhibit niche differentiation, with water depth being the most influential factor, although there is some overlap of habitat. On the Chatham Rise, the main overlap range appears quite compact (from about 340 to 540 m). In the Southland/sub-Antarctic region, the overlap range is wider (about 350 to 770 m). Stomach contents indicate that both species are predominantly benthic feeders.

No published information is available on the age or growth rate of any *Hydrolagus* species, or even any species in the family Chimaeridae. A research report by Francis & Ó Maolagáin (2001) found that eye lens diameter showed potential as an ageing technique but further work was needed. They calculated Von Bertalanffy parameters (Table 4) from trawl survey caught fish and found that growth rates were similar and moderately rapid for males and females with both sexes reaching 50 cm in 5–9 years. They caution the use of these parameters, however, as ageing of dark ghost sharks has not been validated. Length-frequency histograms indicate that females grow to a larger size than males. Without population age structures or confident estimates of longevity, it is not possible to estimate natural or total mortalities.

On the Chatham Rise, the estimated size at 50% sexual maturity for dark ghost sharks is 52–53 cm for males and 62–63 cm for females. As for most other elasmobranchs, ghost shark fecundity is likely to be low.

Length-weight parameters are shown in Table 5.

Table 4: Von Bertalanffy growth parameters for dark ghost shark. Source: Francis & Ó Maolagáin (2001).

Region	Sex	Von Bertalanffy growth parameters		
		L_{∞}	K	t_0
East coast South Island	Female	135.3	0.052	-0.94
	Male	89.0	0.091	-0.61
West coast South Island	Female	123.0	0.065	-1.15
	Male	123.4	0.044	-1.43
Stewart–Snares Shelf	Female	122.1	0.087	-1.01
	Male	108.0	0.073	-1.34
Chatham Rise	Female	97.0	0.090	-1.17
	Male	-	-	-

Table 5: Length-weight parameters for dark ghost shark.

1. $\text{Weight} = a (\text{length})^b$ (Weight in g, length in cm chimaera length)

FMA	Estimate		Source
	a	b	
Chatham Rise	0.002986	3.170546	O'Driscoll et al (2011)
Sub-Antarctic	0.001653	3.3256	Bagley et al (2013)

3. STOCKS AND AREAS

The only information which may indicate a stock boundary is an apparent difference in maximum size of dark ghost sharks, with both males and females from the Chatham Rise attaining a maximum size 3–4 cm greater than those in Southland/sub-Antarctic waters.

Horn (1997) proposed that ghost sharks be managed as three Fishstocks, i.e., east coast New Zealand (FMAs 1–4), Stewart-Snares shelf and Campbell Plateau (FMAs 5 and 6), and west coast New Zealand (FMAs 7, 8, and 9). Areas of narrow continental shelf separate these FMA groupings, so they could well provide barriers to stock mixing for pale ghost shark which have a preference for deeper water. This would be less influential for dark ghost shark, however, which are found much shallower. Pale ghost shark were given the QMAs recommended by Horn when introduced into the QMS, but dark ghost shark were already based on the generic FMAs.

4. STOCK ASSESSMENT

No assessment of any stocks of dark ghost shark has been completed. Therefore, no estimates of yield are available.

4.1 Estimates of fishery parameters and abundance

Estimates of fishery parameters are not available for dark ghost sharks. Several time series of relative biomass estimates are available from fishery independent trawl surveys (Table 6), but wide fluctuations between years suggest the need for caution in using these as indicators of relative abundance. The Chatham Rise time series may provide a reasonable index of abundance for GSH 4, but not GSH 3 as the survey does not fish shallower than 200 m where dark ghost shark are abundant. Much of GSH 3 is covered by the winter east coast South Island trawl survey however, which is optimised for dark ghost shark among other species.

DARK GHOST SHARK (GSH)

Table 6: Biomass indices (t) and coefficients of variation (CV). Estimates for the Chatham Rise and sub-Antarctic summer surveys on *Tangaroa* are for core strata only (200–800 and 300–800 m respectively).

FMA	Area	Vessel	Trip code	Date	Biomass	% CV
3 & 4	Chatham Rise	<i>Tangaroa</i>	TAN9106	Jan-Feb 1992	6 700	11.1
			TAN9212	Jan-Feb 1993	5 950	9.2
			TAN9401	Jan-94	10 360	15.3
			TAN9501	Jan-95	3 490	11.2
			TAN9601	Jan-96	6 170	12.4
			TAN9701	Jan-97	6 240	11.7
			TAN9801	Jan-98	6 720	14.1
			TAN9901	Jan-99	12 125	23.4
			TAN0001	Jan-00	9 154	25.2
			TAN0101	Jan-01	10 356	12
			TAN0201	Jan-02	9 997	11.1
			TAN0301	Jan-03	10 341	9.1
			TAN0401	Jan-04	10 471	15
			TAN0501	Jan-05	11 885	16.3
			TAN0601	Jan-06	11 502	12
			TAN0701	Jan-07	7 852	11
			TAN0801	Jan-08	9 391	10.9
			TAN0901	Jan-09	8 445	13.7
			TAN1001	Jan-10	11 596	16.8
			TAN1101	Jan-11	6 588	17
			TAN1201	Jan-12	13 162	20.6
			TAN1301	Jan-13	11 723	11.6
			TAN1401	Jan-14	9 050	18
			TAN1601	Jan-16	11 925	12
5 & 6	Southland Sub-Antarctic	<i>Tangaroa</i> (summer)	TAN9105	Nov-Dec 1991	1 030	25.4
			TAN9211	Nov-Dec 1992	710	43.2
			TAN9310	Nov-Dec 1993	1 060	33.6
			TAN0012	Nov-Dec 2000	1 459	89.6
			TAN0118	Nov-Dec 2001	1 391	35.7
			TAN0219	Nov-Dec 2002	175	37.7
			TAN0317	Nov-Dec 2003	382	48.9
			TAN0414	Nov-Dec 2004	843	41.7
			TAN0515	Nov-Dec 2005	517	40
			TAN0617	Nov-Dec 2006	354	32
			TAN0714	Nov-Dec 2007	659	37
			TAN0813	Nov-Dec 2008	1128	32
			TAN0911	Nov-Dec 2009	433	43
			TAN1117	Nov-Dec 2011	3 709	75
			TAN1215	Nov-Dec 2012	1 794	68.3
		<i>Tangaroa</i> (autumn)	TAN9204	Mar-Apr 1992	3 740	48.6
			TAN9304	Apr-May 1993	750	44.7
			TAN9605	Mar-Apr 1996	3 080	47.6
			TAN9805	Apr-May 1998	2 490	44
5	Stewart-Snares#	<i>Tangaroa</i>	TAN9301	Feb-Mar 1993	120	44
			TAN9402	Feb-Mar 1994	490	43
			TAN9502	Feb-Mar 1995	790	71
			TAN9604	Feb-Mar 1996	1 870	63
2	East coast North Island	<i>Kaharoa</i>	KAH9304	Mar-Apr 1993	450	61.5
			KAH9402	Feb-Mar 1994	40	41.3
			KAH9502	Feb-Mar 1995	10	48.6
			KAH9602	Feb-Mar 1996	80	33.5
3	ECSI winter surveys	<i>Kaharoa</i>	KAH9105	May-91	962	42
			KAH9205	May-92	934	44
			KAH9306	May-93	2 911	42
			KAH9406	May-94	2 702	25
			KAH9606	May-96	3 176	23
			KAH0705	May-07	4 483	25
			KAH0806	May-June-08	3 763	20
			KAH0905	May-Jun-09	4 330	24
			KAH1207	Apr-Jun-13	10 704	29
			KAH1402	Apr-Jun-14	13 137	26
			KAH1605	Apr-Jun-16	15 271	26

Table 6 [continued]

FMA	Area	Vessel	Trip code	Date	Biomass	% CV
3	ECSI summer surveys	<i>Kaharoa</i>	KAH9618	Dec '96 - Jan '97	3 066	18
			KAH9704	Dec '97 - Jan '98	5 870	33
			KAH9809	Dec '98 - Jan '99	7 416	27
			KAH9917	Dec '99 - Jan '00	2 512	19
			KAH0014	Dec '00 - Jan '01	2 950	18
7	West coast South Island	<i>Kaharoa</i>	KAH9204	Mar-Apr 1992	380	20
			KAH9404	Mar-Apr 1994	720	14.3
			KAH9504	Mar-Apr 1995	770	23.7
			KAH9701	Mar-Apr 1997	1 590	21.2
			KAH0004	Mar-Apr 2000	2 260	9
			KAH0304	Mar-Apr 2003	540	15
			KAH0503	Mar-Apr 2005	830	22
			KAH0704	Mar-Apr 2007	2 215	21
			KAH0904	Mar-Apr 2009	900	17
			KAH1104	Mar-Apr 2011	2 363	23
			KAH1305	Mar-Apr 2013	981	23

4.2 Biomass estimates

Biomass estimates from various trawl surveys are given in Table 6. Of those, ongoing estimates are available from random stratified bottom trawl surveys from the east coast South Island, Chatham Rise, sub-Antarctic, and west coast South Island trawl surveys.

Total biomass in the east coast South Island winter surveys core strata (30–400 m) increased 16-fold between 1992 and 2016 (Table 6, Figure 2). All surveys had a large component of pre-recruit biomass ranging from 30–61%. In 2016 the pre-recruit biomass was relatively high at 54% of total biomass. The juvenile and adult biomass (based on length-at-50% maturity) of both sexes have generally increased proportionately over the time series and juvenile biomass comprised about half of the total biomass. In 2016 the juvenile biomass was 49% of total biomass. (Beentjes et al 2016).

Distribution over the ECSI winter trawl survey time series was similar and was confined to the continental slope and edge mainly in the Canterbury Bight, although the larger biomass from 2007 to 2016 is commensurate with a slightly expanded distribution throughout the survey area in this depth range and into Pegasus Bay. The size distributions in each of the last nine surveys (1993–2016) were similar and generally bimodal (Beentjes et al 2016). The 2012, 2014 and 2016 length frequency distributions were distinct from previous years with relatively large numbers of adults or mature fish. The distributions differ from those of the Chatham Rise and Southland/Sub-Antarctic surveys in that ECSI has a large component of juvenile fish, suggesting that this area may be an important nursery ground for dark ghost shark.




Figure 2: Biomass for dark ghost shark from the east coast South Island winter trawl surveys in core strata (30–400 m). Error bars are ± 2 standard errors.

The Chatham Rise trawl survey time series is not optimised for dark ghost shark and there has been some year-to-year variation between surveys, particularly for the first ten years (Figure 3). This time

DARK GHOST SHARK (GSH)

series may provide a reasonable index of abundance for that part of the eastern fishery (see Section 5) covered by GSH 4. However the survey extends into GSH 3 where commercial catches of dark ghost shark are significant but shallower than the survey's starting depth of 200 m.




Figure 3: Biomass for dark ghost shark from the Chatham Rise trawl survey. Error bars are ± 2 standard errors.

Biomass indices from the sub-Antarctic trawl survey time series are significantly lower than those for the east coast South Island and Chatham Rise surveys. Indices have fluctuated somewhat (Figure 4). The large spike seen in 2011 is due to randomly allocated stations within stratum 6 (300–600 m) being located at the shallower, northern end of the stratum where dark ghost shark are more likely to be encountered. The starting depth of 300 m may mean that this survey is unlikely to be a reliable index of abundance.




Figure 4: Biomass trends $\pm 95\%$ CI (estimated from survey CVs assuming a lognormal distribution) from the Sub-Antarctic trawl survey.

Biomass estimates from the west coast South Island inshore trawl survey are lower than those from the east coast South Island and Chatham Rise surveys. Estimates fluctuate considerably and are unlikely to reflect real changes in abundance (Figure 5).




Figure 5: Biomass trends $\pm 95\%$ CI (estimated from survey CVs assuming a lognormal distribution) from the West Coast South Island trawl survey.

4.3 Estimation of Maximum Constant Yield (*MCY*)

As there are no available estimates of biomass or harvest rates, the only possible method of calculating maximum constant yield is $MCY = cY_{AV}$ (Method 4). However, it was decided that no estimates of *MCY* would be presented because:

- i. *M* (and hence, the natural variability factor *c*) is unknown;
- ii. the level of discarding is unknown and may have been considerable; and
- iii. no sufficiently long period of catches was available where there were no systematic changes in catch or effort (noting that the period of catches from which Y_{AV} is derived should be at least half the exploited life span of the fish).

4.4 Estimation of Current Annual Yield (*CAY*)

In the absence of estimates of current biomass, *CAY* has not been estimated.

4.5 Other yield estimates and stock assessment results

No other yield estimates are available.

4.6 Other factors

Elasmobranchs are believed to have a strong stock-recruit relationship; the number of young born is related directly to the number of adult females. Ghost shark fecundity is unknown, but is probably low. Assuming a strong stock-recruit relationship, Francis & Francis (1992) showed that the estimates of *MCY* obtained using the equations in current use in New Zealand stock assessments were overly optimistic for rig, and it is likely that they are also unsuitable for ghost sharks.

A data informed qualitative risk assessment was completed on all chondrichthyans (sharks, skates, rays and chimaeras) at the New Zealand scale in 2014 (Ford et al 2015). Dark ghost shark was ranked seventh highest in terms of risk of the eleven QMS chondrichthyan species. Data were described as existing but poor for the purposes of the assessment and consensus over this risk score was achieved by the expert panel. This risk assessment does not replace a stock assessment for this species but may influence research priorities across species.

5. STATUS OF THE STOCKS

Stock Structure Assumptions

Based on differences in length frequency distributions between the sub-Antarctic and Chatham Rise trawl surveys, and the location of commercial catches, there are most likely two main stocks of dark ghost shark.

1. The eastern fishery; extending from the upper east coast of the South Island and out east across the Chatham Rise.
2. The southern fishery; extending from the lower east coast of the South Island, south around the Stewart/Snares Shelf, Campbell Plateau, and Puysegur trench.


Further work needs to be done to investigate what if any relationship there is between dark ghost shark caught on the west coast of the South Island, around both coasts of the North Island, and the eastern and southern stocks.

- **Chatham Rise**


Stock Status	
Year of Most Recent Assessment	2016
Assessment Runs Presented	-
Reference Points	Management Target: 40% B_0 Soft Limit: 20% B_0 Hard Limit: 10% B_0 Overfishing threshold: Not defined
Status in relation to Target	Unknown
Status in relation to Limits	Unknown
Status in relation to Overfishing	Unknown

Historical Stock Status Trajectory and Current Status

ECSI core strata



Biomass trajectory for dark ghost shark from the Chatham Rise trawl survey. Error bars are ± 2 standard errors.



Fishery and Stock Trends

Recent Trend in Biomass or Proxy	Biomass indices from the east coast South Island inshore trawl survey time series have been steadily increasing since 2009. Biomass indices from the Chatham Rise have fluctuated somewhat over the time series. Estimates from the last ten years have been more stable.
Recent Trend in Fishing Intensity or Proxy	Landings have been stable for the last five years from GSH 3, and relatively stable from GSH 4, apart from a small spike in the 2007–08 fishing year.
Other Abundance Indices	-
Trends in Other Relevant Indicators or Variables	-

Projections and Prognosis

Stock Projections or Prognosis	Unknown
Probability of Current Catch or TACC causing Biomass to remain below or to decline below Limits	Soft Limit: Unknown Hard Limit: Unknown
Probability of Current Catch or TACC causing Overfishing to continue or to commence	Unknown, but there is no evidence of a systematic decline in biomass indices from either the east coast of the South Island or the Chatham Rise.

Qualifying Comments


-

Fishery Interactions

Dark ghost shark in the eastern fishery is caught exclusively as bycatch in other target fisheries with the two most important ones being hoki followed by arrow squid. For both target fisheries, incidental interactions and associated mortalities are noted for New Zealand fur seals and seabirds, and low productivity species taken in the fisheries include basking sharks and deepsea skates.

DARK GHOST SHARK (GSH)

- Southern stock**

Stock Status	
Year of Most Recent Assessment	2011
Assessment Runs Presented	-
Reference Points	Management Target: 40% B_0 Soft Limit: 20% B_0 Hard Limit: 10% B_0 Overfishing threshold: Not defined
Status in relation to Target	Unknown
Status in relation to Limits	Unknown
Status in relation to Overfishing	Unknown
Historical Stock Status Trajectory and Current Status	
 <p>Biomass trends $\pm 95\%$ CI (estimated from survey CVs assuming a lognormal distribution) from the Sub-Antarctic trawl survey.</p>	

Fishery and Stock Trends	
Recent Trend in Biomass or Proxy	Biomass indices from the summer sub-Antarctic trawl survey time series have been relatively flat for the last few years apart from a large spike in 2011 due to a number of randomly allocated stations occurring at the shallower end of the depth range for dark ghost shark.
Recent Trend in Fishing Intensity or Proxy	Unknown. Landings have fluctuated somewhat from GSH 5 in recent years, and have been relatively stable from GSH 6.
Other Abundance Indices	-
Trends in Other Relevant Indicators or Variables	-

Projections and Prognosis	
Stock Projections or Prognosis	Unknown
Probability of Current Catch or TACC causing Biomass to remain below or to decline below Limits	Soft Limit: Unknown Hard Limit: Unknown
Probability of Current Catch or TACC causing Overfishing to continue or to commence	Unknown, but there is no evidence of a systematic decline in biomass indices from the sub-Antarctic survey.

Qualifying Comments
-


Fishery Interactions
Dark ghost shark in the southern fishery is caught exclusively as bycatch in other target fisheries with the two most important ones being arrow squid followed by hoki. For both target fisheries, incidental interactions and associated mortalities have been recorded for New Zealand fur seals and seabirds, and low productivity species taken in the fisheries include basking sharks and deepsea skates. Interactions with other species are currently being characterised.

Table 7: Summary of TACCs (t) and reported landings (t) for dark ghost shark for the most recent fishing year.

			2016–17 Actual TACC	2016–17 Estimated Landings
Fishstock		QMA		
GSH 1	Auckland (East)	1	22	21
GSH 2	Central (East)	2	89	44
GSH 3	South-east (Coast)	3	1 185	493
GSH 4	South-east (Chatham)	4	370	223
GSH 5	Southland	5	109	83
GSH 6	Sub-Antarctic	6	95	59
GSH 7	Challenger	7	1 121	423
GSH 8	Central (West)	8	34	19
GSH 9	Auckland (West)	9	22	14
GSH 10	Kermadec	10	0	0
Total			3 047	1 379

6. FOR FURTHER INFORMATION

- Bagley, N W; O'Driscoll, R L; Oeffner, J (2013) Trawl survey of hoki and middle-depth species in the Southland and Sub-Antarctic areas, November–December 2011 (TAN1117). *New Zealand Fisheries Assessment Report 2013/23*. 70 p.
- Beentjes, M P; MacGibbon, D; Parkinson, D (2016) Inshore trawl survey of Canterbury Bight and Pegasus Bay, April–June 2016 (KAH1605). *New Zealand Fisheries Assessment Report 2016/61*. 135 p.
- Ford, R B; Galland, A; Clark, M R; Crozier, P; Duffy, C A J; Dunn, M R; Francis, M P; Wells, R (2015) Qualitative (Level 1) Risk Assessment of the impact of commercial fishing on New Zealand Chondrichthyans. *New Zealand Aquatic Environment and Biodiversity Report No. 157*. 111 p.
- Francis, M P; Francis, R I C C (1992) Growth, mortality, and yield estimates for rig (*Mustelus lenticulatus*). New Zealand Fisheries Assessment Research Document 1992/5. 32 p. (Unpublished document held by NIWA library, Wellington.)
- Francis, M P; McMillan, P; Lasenby, R; Didier, D (1998) How to tell dark and pale ghost sharks apart. *Seafood New Zealand 6 (11)*: 29–30.
- Francis, M P; Ó Maolagáin, C O (2001) Development of ageing techniques for dark ghost shark (*Hydrolagus novaezelandiae*). Final Research Report for Ministry of Fisheries Research Project MOF2000/03C. 10 p. (Unpublished report held by Fisheries New Zealand Wellington.)
- Francis, M P.; Paul, L J (2013) New Zealand inshore finfish and shellfish commercial landings, 1931–82. *New Zealand Fisheries Assessment Report 2013/55*. 136 p.
- Horn, P L (1997) A summary of biology and commercial landings, and a stock assessment of ghost sharks (*Hydrolagus* spp.) in New Zealand waters. New Zealand Fisheries Assessment Research Document 97/3. 36 p. (Unpublished document held by NIWA library, Wellington.)
- MacGibbon, D J; Stevenson, M L (2013) Inshore trawl survey of the west coast South Island and Tasman and Golden Bays, March–April 2013 (KAH1305) *New Zealand Fisheries Assessment Report 2013/66*. 115 p.
- O'Driscoll, R L; MacGibbon, D; Fu, D; Lyon, W; Stevens, D W (2011) A review of hoki and middle depth trawl surveys of the Chatham Rise, January 1992–2010. *New Zealand Fisheries Assessment Report 2011/47*.
- Stevens, D; Livingston, M; Bagley, N (2001) Trawl survey of hoki and middle depth species on the Chatham Rise, January 2001 (TAN0101). Trawl survey of hoki and middle depth species on the Chatham Rise, January 2001 (TAN0101). *NIWA Technical Report 116*. 61 p.
- Stevens, D W; O'Driscoll, R L; Ballara, S L; Bagley, N; Horn, P L (2011) Chatham Rise Trawl Survey, 2 Jan - 28 Jan 2011 (TAN1011). WG-HOK-2011/X. (Unpublished report held by Fisheries New Zealand, Wellington.)
- Stevens, D.W.; O'Driscoll, R.L.; Oeffner, J.; Ballara, S.L.; Horn, P.L. (2014). Trawl survey of hoki and middle depth species on the Chatham Rise, January 2013 (TAN1301). *New Zealand Fisheries Assessment Report 2014/02*. 110 p.
- Stevens, D.W.; O'Driscoll, R.L.; Ldroit, Y.; Ballara, S.L.; MacGibbon, D.J.; Horn, P.L. (2015). Trawl survey of hoki and middle depth species on the Chatham Rise, January 2014 (TAN1401). *New Zealand Fisheries Assessment Report 2015/19*. 119 p.
- Stevens, D W; O'Driscoll, R L; Ballara, S L; Ldroit, Y (2017) Trawl survey of hoki and middle-depth species on the Chatham Rise, January 2016 (TAN1601). *New Zealand Fisheries Assessment Report 2017/08*. 131 p.

PALE GHOST SHARK (GSP)*(Hydrolagus bemisi)***1. FISHERY SUMMARY****1.1 Commercial fisheries**

Two species (dark and pale ghost sharks) make up virtually all the commercial ghost shark landings. Pale ghost shark (*Hydrolagus bemisi*) was introduced into the QMS from the beginning of the 1999–00 fishing year as three Fishstocks: GSP 1 (FMAs 1 to 4, and 10), GSP 5 (FMAs 5 and 6) and GSP 7 (FMAs 7, 8 and 9).

Both ghost shark species are taken almost exclusively as a bycatch of other target trawl fisheries. In the 1990s, about 43% of ghost sharks were landed as a bycatch of the hoki fishery, with fisheries for silver warehou, arrow squid and barracouta combining to land a further 36%. The two ghost shark species were seldom differentiated on catch landing returns prior to the start of the 1998–99 fishing year. Estimated landings of both species by foreign licensed and joint venture vessels over the period 1 April 1978 to 30 September 1983 are presented in Table 1. Landings by domestic (inshore) vessels would have been negligible during this time period. The unknown quantities of ghost sharks that were discarded and not recorded are likely to have resulted in under-reported total catches over the full period for which data are available.

Table 1: Reported landings (t) of both ghost shark species by fishing year and EEZ area, taken by foreign licensed and joint venture vessels. An approximation of these areas with respect to current FMA boundaries is used to assign catches to QMAs. No data are available for the 1980–81 fishing year.

Year	EEZ Area												Total
	B	C(M)	C(I)	D	E(B)	E(P)	E(C)	E(A)	F(E)	F(W)	G	H	
FMA	1&2	3		4	5			6	7		8	9	
1978–79*	1	37	99	26	3	16	11	88	90	8	68	17	465
1979–80*	1	55	54	426	10	4	28	138	183	7	1	5	912
1980–81*													-
1981–82*	0	84	28	117	0	2	6	29	71	9	4	0	350
1982–83*	0	108	35	84	0	2	17	98	99	29	1	1	474
1983–83#	0	84	41	73	0	0	17	5	16	17	0	0	253

* 1 April to 31 March. # 1 April to 30 Sept

In the early to mid 1980s, about half of the reported ghost shark landings were from FMA 3. Virtually all the additional catch was spread over FMAs 4–7. In 1988–89, landings from west coast South Island (FMA 7) began to increase, almost certainly associated with the development of the hoki fishery. In 1990–91, significant increases in landings were apparent on the Chatham Rise, off

PALE GHOST SHARK (GSP)

southeast South Island, and on the Campbell Plateau. The development of fisheries for non-spawning hoki was probably responsible for these increases.

Estimated landings of pale ghost shark by QMA are shown in Table 2. Landings from 1983–84 to 1994–95 were derived by splitting all reported ghost shark landings into depth and area bins, and allocating to species based on distribution data derived from trawl surveys (Section 2). Landings from 1995–96 to 1998–99 were estimated assuming that pale ghost shark made up 30% of the total ghost shark catch in FMAs 5 and 6, and 25% in all other FMAs.

From 1 Oct 1999 TACCs were set for pale ghost shark fishstocks as follows: GSP 1 509 t, GSP 5 118 t and GSP 7 176 t. The TAC in each case was set equal to the TACC. Estimated and reported landings for this period are shown in Table 3, while Figure 1 shows the historical landings and TACC values for the main GSP stocks. The fisheries in GSP 1 and GSP 5 exceeded the TACC by large amounts, possibly as a result of better reporting of catches. From 1 October 2004 the TACCs for GSP 1 and GSP 5 were increased to 1150 t and 454 t respectively, the level of catch being reported from the fisheries. Catches have since declined to well below the TACC levels.

In GSP 1, catches are mainly taken on the Chatham Rise while in GSP 5 catches are mainly taken in the Sub-Antarctic area; both as bycatch of the hoki trawl fisheries. Estimated catches appear to have been under-reported both before and after the introduction to the QMS. The original TACCs were based on estimated catches, but these are likely to have been much lower than the actual catches. Estimated catches on TCEPR forms since 1999–2000 have been only 25–30% of the QMR totals.

Table 2: Estimated landings (t) of pale ghost shark by Fisheries Management Area for fishing years 1982–83 to 1998–99 based on the reported landings of both species combined. The estimated landings up to 1994–95 are based on data in the 1997 Plenary Report. Landings from 1995–96 to 1998–99 were estimated assuming pale ghost shark made up 30% of the total ghost shark catch in FMAs 5 and 6, and 25% in all other FMAs.

	FMA										
	1	2	3	4	5	6	7	8	9	10	Total
1982–83	1	1	74	35	21	13	2	1	0	0	148
1983–84	0	1	63	24	11	15	7	1	0	0	122
1984–85	1	1	60	49	16	19	12	0	0	0	158
1985–86	1	1	96	23	10	14	7	1	0	0	153
1986–87	1	2	110	27	11	12	13	1	0	0	177
1987–88	1	1	138	21	13	2	15	1	0	0	192
1988–89	2	7	124	9	19	2	34	1	0	0	198
1989–90	1	3	86	8	41	5	33	5	0	0	182
1990–91	1	7	148	63	61	82	39	1	0	0	402
1991–92	1	2	218	95	64	54	35	2	1	0	472
1992–93	2	1	227	99	77	55	53	7	0	0	521
1993–94	1	2	173	42	36	32	99	4	0	0	389
1994–95	1	1	246	62	27	26	234	1	0	0	598
1995–96	4	12	226	84	30	29	183	3	1	0	572
1996–97	6	22	272	134	40	58	309	3	3	0	847
1997–98	6	6	256	87	30	58	57	1	4	0	505
1998–99	6	20	315	107	27	47	136	2	7	0	667

1.2 Recreational fisheries


Current catches of ghost sharks by recreational fishers are believed to be negligible in all areas.

1.3 Customary non-commercial fisheries

Quantitative information on the current level of customary non-commercial take is not available.

Table 3: Estimated landings (t) of pale ghost shark by Fishstock for 1999–2000 to 2016–17 and actual TACCs set from 1999–2000 (QMR data).

Fishstock FMA (s)	GSP 1 1,2,3,4,10		GSP 5 5,6		GSP 7 7,8,9		Total	
	Landings	TACC	Landings	TACC	Landings	TACC	Landings	TACC
1999–00	577	509	216	118	35	176	828	803
2000–01	1 142	509	454	118	16	176	1 613	803
2001–02	1 033	509	545	118	71	176	1 649	803
2002–03	1 277	509	602	118	16	176	1 895	803
2003–04	1 009	509	529	118	15	176	1 553	803
2004–05	635	1 150	247	454	5	176	887	1 780
2005–06	565	1 150	134	454	9	176	708	1 780
2006–07	553	1 150	226	454	15	176	794	1 780
2007–08	473	1 150	329	454	16	176	818	1 780
2008–09	486	1 150	294	454	15	176	795	1 780
2009–10	534	1 150	206	454	11	176	751	1 780
2010–11	395	1 150	203	454	13	176	611	1 780
2011–12	447	1 150	201	454	10	176	659	1 780
2012–13	510	1 150	163	454	25	176	697	1 780
2013–14	409	1 150	286	454	33	176	727	1 780
2014–15	476	1 150	243	454	38	176	759	1 780
2015–16	493	1 150	171	454	26	176	690	1 780
2016–17	577	1 150	324	454	25	176	926	1 780

**Figure 1: Reported commercial landings and TACC for the three main GSP stocks. From top: GSP 1 (Auckland East), GSP 5 (Southland). Note that these figures do not show data prior to entry into the QMS.**

PALE GHOST SHARK (GSP)




Figure 1 Continued: Reported commercial landings and TACC for the three main GSP stocks. GSP 7 (Challenger). Note that these figures do not show data prior to entry into the QMS.

1.4 Illegal catch

Quantitative information on the level of illegal catch is not available. In 1998–99 (when dark ghost shark were in the QMS, but pale ghost shark were not), a quantity of dark ghost shark were reported as pale ghost shark.

1.5 Other sources of mortality

Ghost sharks have been dumped and not reported in the past by commercial fishers in FMAs 1 and 2. Similar behaviour is believed to occur in all other FMAs. The extent of the unreported dumping is unknown in all areas.

2. BIOLOGY

Pale ghost shark occur throughout the EEZ and have been recorded in depths ranging from 270 to 1200 m. They are most abundant in depths of 400–1000 m on the Chatham Rise and Southland/Sub-Antarctic, but are uncommon north of 40° S and appear to inhabit a narrower depth range in that region (600–950 m).

Trawl surveys show that dark and pale ghost shark exhibit niche differentiation, with water depth being the most influential factor, although there is some overlap of habitat. On the Chatham Rise, the main overlap range appears quite compact (from about 340 to 540 m). In the Southland/Sub-Antarctic region, the overlap range is wider (about 350 to 770 m). Stomach contents indicate that both species are predominantly benthic feeders.

No published information is available on the age or growth rate of any *Hydrolagus* species, or even any species in the family Chimaeridae. Length-frequency histograms indicate that females grow to a larger size (and presumably have a faster growth rate) than males. Hard parts of pale ghost shark have not yet been examined to check the existence of any banding pattern that may represent annual growth zones. Without population age structures or confident estimates of longevity it is not possible to estimate natural or total mortalities. A recent study has shown that eye lens measurements and spine band counts are potentially useful ageing techniques for dark ghost sharks (Francis & Ó Maolagáin 2001). However, these techniques have yet to be validated.

On the Chatham Rise, the estimated size at 50% sexual maturity for pale ghost sharks is 59–60 cm for males and 69–70 cm for females. As for most other elasmobranchs, ghost shark fecundity is likely to be low.

Biological parameters relevant to the stock assessment are shown in Table 4.

Table 4: Estimates of biological parameters for pale ghost shark, from Horn (1997).

FMA	Estimate	
1. Weight = a (length) ^b (Weight in g, length in cm chimaera length)		
Pale ghost shark	a	b
3 & 4	0.00512	3.037
5 & 6	0.00946	2.883

3. STOCKS AND AREAS

Horn (1997) proposed that ghost sharks be managed as three Fishstocks, i.e., east coast New Zealand (FMAs 1–4), Stewart-Snares shelf and Campbell Plateau (FMAs 5 and 6), and west coast New Zealand (FMAs 7, 8, and 9). Areas of narrow continental shelf separate these FMA groupings, so they could well provide barriers to stock mixing, particularly for the pale ghost shark. The deep water separating the Bounty Platform from the Campbell Plateau may also provide a barrier to mixing, and these areas may hold separate stocks.

4. STOCK ASSESSMENT

No assessment of any stocks of ghost shark has been completed. Therefore, no estimates of yield are available.

4.1 Estimates of fishery parameters and abundance

Table 5: Biomass indices (t) and coefficients of variation (CV)

GSP	Area	Vessel	Trip code	Date	Pale ghost shark	
					Biomass	% CV
1	Chatham Rise	<i>Tangaroa</i>	TAN9106	Jan–Feb 1992	6 060	5.7
			TAN9212	Jan–Feb 1993	3 570	7
			TAN9401	Jan-94	5 900	8.6
			TAN9501	Jan-95	2 750	8.4
			TAN9601	Jan-96	7 900	10
			TAN9701	Jan-97	2 870	12.2
			TAN9801	Jan-98	4 052	9.3
			TAN9901	Jan-99	5 272	9.7
			TAN0001	Jan-00	4 892	7.6
			TAN0101	Jan-01	7 094	9
			TAN0201	Jan-02	4 896	10
			TAN0301	Jan-03	4 653	12.1
			TAN0401	Jan-04	3 627	8.6
			TAN0501	Jan-05	4 061	9.2
			TAN0601	Jan-06	3 237	11
			TAN0701	Jan-07	4 766	9.0
			TAN0801	Jan-08	3 235	6.1
			TAN0901	Jan-09	3 995	7.6
			TAN1001	Jan-10	3 216	11.7
			TAN1101	Jan-11	2 550	14.2
5	Southland Sub-Antarctic	<i>Tangaroa</i>	TAN1201	Jan-12	4 327	8.5
			TAN1301	Jan-13	4 270	18.0
			TAN9105	Nov–Dec 1991	11 210	6.1
			TAN9211	Nov–Dec 1992	4 750	7.2
			TAN9310	Nov–Dec 1993	11 670	9.4
			TAN0012	Nov–Dec 2000	17 823	12.4
			TAN0118	Nov–Dec 2001	11 219	8.8
			TAN0219	Nov–Dec 2002	9 297	9.3
			TAN0317	Nov–Dec 2003	10 360	8.7

PALE GHOST SHARK (GSP)

Table 5 Continued:

			TAN0414	Nov–Dec 2004	8 549	10.3
			TAN0515	Nov–Dec 2005	9 416	10
			TAN0617	Nov–Dec 2006	12 619	10
			TAN0714	Nov–Dec 2007	13 107	11
			TAN0813	Nov–Dec 2008	10 098	13
			TAN0911	Nov–Dec 2009	13 553	9
			TAN1117	Nov–Dec 2011	11 677	9.6
			TAN1215	Nov–Dec 2012	16 181	12.6
5	Southland	Tangaroa	TAN9204	Mar–Apr 1992	10 530	6.1
	Sub-Antarctic		TAN9304	Apr–May 1993	14 640	9.5
			TAN9605	Mar–Apr 1996	16 380	9.9
			TAN9805	Apr–May 1998	15 758	10

Estimates of fishery parameters are not available for ghost sharks. Several time series of relative biomass estimates are available from trawl surveys (Table 5). In 2004, the Plenary agreed that the trawl survey series for both GSP 1 and GSP 5 indicated that previous catch levels had made little impact on the biomass of pale ghost shark, however, the actual level of catch is not known. The recorded catch history for this species is likely to underestimate actual catches. The trawl series fluctuates over time and decreases in 2010 and 2011 on the Chatham Rise. In the Sub-Antarctic the trawl biomass indices have increased since 2005.

4.2 Biomass estimates

No biomass estimates are available for ghost shark.

4.3 Yield estimates and projections

As no estimate of biomass or harvest rate are available, the only possible method of calculating maximum constant yield is $MCY = cY_{AV}$ (Method 4).

However, it was decided that no estimates of MCY would be presented because:

- i. M (and hence, the natural variability factor c) is unknown;
- ii. the level of discarding is unknown and may have been considerable; and
- iii. no sufficiently long period of catches was available where there were no systematic changes in catch or effort (noting that the period of catches from which Y_{AV} is derived should be at least half the exploited life span of the fish).

In the absence of estimates of current biomass, CAY has not been estimated.

4.4 Other factors

Elasmobranchs are believed to have a strong stock-recruit relationship; the number of young born is related directly to the number of adult females. Ghost shark fecundity is unknown, but is probably low. Assuming a strong stock-recruit relationship, Francis & Francis (1992) showed that the estimates of MCY obtained using the equations in current use in New Zealand stock assessments were overly optimistic for rig, and it is likely that they are also unsuitable for ghost sharks.

A data informed qualitative risk assessment was completed on all chondrichthyans (sharks, skates, rays and chimaeras) at the New Zealand scale in 2014 (Ford et al 2015). Pale ghost shark was ranked ninth highest in terms of risk of the eleven QMS chondrichthyan species. Data were described as existing but poor for the purposes of the assessment and no consensus over this risk score was achieved by the expert panel. This risk assessment does not replace a stock assessment for this species but may influence research priorities across species.


5. STATUS OF THE STOCKS

No estimates of current and reference biomass are available for pale ghost shark.

GSP 1

Stock Status	
Year of Most Recent Assessment	2011
Assessment Runs Presented	
Reference Points	Target: 40% B_0 Soft Limit: 20% B_0 Hard Limit: 10% B_0 Overfishing threshold:-
Status in relation to Target	Unknown
Status in relation to Limits	Unlikely (< 40%) to be below soft limit Very Unlikely (< 10%) to be below hard limit
Status in relation to Overfishing	-

Historical Stock Status Trajectory and Current Status



Doorspread biomass estimates of pale ghost shark (error bars are \pm two standard deviations) from the Chatham Rise, from *Tangaroa* surveys from 1992 to 2011.

Fishery and Stock Trends	
Recent Trend in Biomass or Proxy	Biomass estimates from trawl surveys on the Chatham Rise have fluctuated over the time series showing a decreasing trend since 2001. Precision is generally good in this time series (< 10%). The Working Group considered this index to be suitable to monitor major trends in this stock.
Recent Trend in Fishing Mortality or Proxy	Unknown
Other Abundance Indices	-
Trends in Other Relevant Indicators or Variables	Catches have been well below the TACC since 2004–05.

Projections and Prognosis	
Stock Projections or Prognosis	-
Probability of Current Catch or	Soft Limit: Unlikely (< 40%) at recent catch levels; unknown at

PALE GHOST SHARK (GSP)

TACC causing Biomass to remain below or to decline below Limits	the TACC Hard Limit: Very Unlikely (< 10%) at recent catch levels; unknown at the TACC
Probability of Current Catch or TACC causing Overfishing to continue or to commence	-


Assessment Methodology and Evaluation		
Assessment Type	Level 2 – Partial Quantitative Stock Assessment	
Assessment Method	Evaluation of trawl survey indices on the Chatham Rise	
Assessment Dates	Latest assessment: 2011	Next assessment: Unknown
Overall assessment quality rank		
Main data inputs (rank)	- Research time series of abundance indices (trawl surveys)	
Data not used (rank)	-	
Changes to Model Structure and Assumptions	-	
Major Sources of Uncertainty	The core strata in the trawl survey do not cover the full depth distribution of pale ghost shark.	

Qualifying Comments
The catch history for this species is likely to underestimate actual catches.

Fishery Interactions
The pale ghost shark in GSP 1 is mainly taken as bycatch of the hoki fishery. Interactions with other species are currently being characterised.

GSP 5

Stock Status	
Year of Most Recent Assessment	2011
Assessment Runs Presented	-
Reference Points	Target: 40% B_0 Soft Limit: 20% B_0 Hard Limit: 10% B_0 Overfishing threshold:-
Status in relation to Target	Unknown
Status in relation to Limits	Unlikely (< 40%) to be below soft limit Very Unlikely (< 10%) to be below hard limit
Status in relation to Overfishing	-

Historical Stock Status Trajectory and Current Status

Doorspread biomass estimates of pale ghost shark (error bars are \pm two standard deviations) from the Sub-Antarctic, from *Tangaroa* summer surveys from 1991 to 1993, and 2000 to 2009 (solid line) and autumn surveys from 1992 to 1998 (dashed line).

Fishery and Stock Trends

Recent Trend in Biomass or Proxy	Biomass estimates from trawl surveys on the Sub-Antarctic have increased in recent years. Precision is generally good in this time series (about 10%). The Working Group considered this index to be suitable to monitor major trends in this stock.
Recent Trend in Fishing Mortality or Proxy	Unknown
Other Abundance Indices	-
Trends in Other Relevant Indicators or Variables	Catches have been well below the TACC since 2004–05.

Projections and Prognosis

Stock Projections or Prognosis	Stock size is Unlikely (< 40%) to change much at current catch levels in FMA 5&6.
Probability of Current Catch or TACC causing Biomass to remain below or to decline below Limits	Soft Limit: Unlikely (< 40%) at recent catch levels; unknown at the TACC Hard Limit: Very Unlikely (< 10%) at recent catch levels; unknown at the TACC
Probability of Current Catch or TACC causing overfishing to continue or to commence	-

Assessment Methodology

Assessment Type	Level 2 - Quantitative stock assessment	
Assessment Method	Evaluation of trawl survey indices on the Chatham Rise	
Assessment Dates	Latest assessment: 2011	Next assessment: Unknown
Overall assessment quality rank	-	
Main data inputs	- Research time series of abundance indices (trawl surveys)	
Data not used (rank)		
Changes to Model Structure and Assumptions	-	
Major Sources of Uncertainty	-	

PALE GHOST SHARK (GSP)

Qualifying Comments

The early catch history for this species is likely to underestimate actual catches.

Fishery Interactions

The pale ghost shark in GSP 5 is mainly taken as bycatch of the hoki fishery. Interactions with other species are currently being characterised.

GSP 7

There are no accepted stock monitoring indices available for GSP 7.

TACCs and reported landings for the 2016–17 fishing year are summarised in Table 6.


Table 6: Summary of TACCs (t) and reported landings (t) of pale ghost shark for the most recent fishing year.

Fishstock		FMA	2016–17 Actual TACC	2016–17 Estimated landings
GSP 1	Auckland (East), Central (East) South-East (Coast) (Chatham), Kermadec	1, 2, 3, 4, 10	1 150	577
GSP 5	Southland, Sub-Antarctic	5, 6	454	324
GSP 7	Challenger, Central (West), Auckland (West)	7, 8, 9	176	25
Total			1 780	926

6. FOR FURTHER INFORMATION

- Ford, R B; Galland, A; Clark, M R; Crozier, P; Duffy, C A J; Dunn, M R; Francis, M P; Wells, R (2015) Qualitative (Level 1) Risk Assessment of the impact of commercial fishing on New Zealand Chondrichthyans. *New Zealand Aquatic Environment and Biodiversity Report No. 157*. 111p.
- Francis, M P; Francis, R I C C (1992) Growth, mortality, and yield estimates for rig (*Mustelus lenticulatus*). New Zealand Fisheries Assessment Research Document 1992/5. 32 p. (Unpublished document held by NIWA library, Wellington.)
- Francis, M P; McMillan, P; Lasenby, R; Didier, D (1998) How to tell dark and pale ghost sharks apart. *Seafood New Zealand* 6 (11): 29–30. (December 1998.)
- Francis, M P; Ó Maolagáin, C (2001) Development of ageing techniques for dark ghost shark (*Hydrolagus novaezelandiae*). Final Research Report for Ministry of Fisheries Research Project MOF2000/03C. 10 p. (Unpublished report held by Fisheries New Zealand, Wellington.)
- Horn, P L (1997) A summary of biology and commercial landings, and a stock assessment of ghost sharks (*Hydrolagus* spp.) in New Zealand waters. New Zealand Fisheries Assessment Research Document 1997/3. 36 p. (Unpublished document held by NIWA library, Wellington.)
- Stevens, D; Livingston, M; Bagley, N (2001) Trawl survey of hoki and middle depth species on the Chatham Rise, January 2001 (TAN0101). Final Research Report for Ministry of Fisheries Research Project HOK2000/02, Objectives 1 and 2. 13 p. (Unpublished report held by Fisheries New Zealand, Wellington.)

GIANT SPIDER CRAB (GSC)

(Jacquinotia edwardsii)

1. FISHERY SUMMARY

1.1 Commercial fisheries

The giant spider crab (*Jacquinotia edwardsii*) was introduced into the Quota Management System on 1 April 2004 with a combined TAC of 451 t and TACC of 419 t. There are no allowances for customary or recreational take, and there is an allowance for other sources of mortality of 32 t. The fishing year is from 1 April to 31 March and commercial catches are measured in greenweight. Up until 2001–02, reported commercial catches of this crab were generally low (Table 1). Since then total reported landings have risen from about 8 t to more than 180 t (Table 1). There was exploratory fishing for this crab in the late 1960s and early 1970s in the Auckland Islands and Pukaki Rise areas and then little interest until, according to Ministry data, the 1999–2000 fishing year. Figure 1 shows the historical landings and TACC for the main GSC stocks.

Table 1: TACCs and reported landings (t) of giant spider crab by Fishstock from 2001–02 to 2016–17 from CELR and CLR data. (N/A = no TACC set). [Continued on next page].

Fishstock	GSC 1		GSC 3		GSC 4		GSC 5		GSC 6	
	Landings	TACC	Landings	TACC	Landings	TACC	Landings	TACC	Landings	TACC
1990–91	< 1	-	0	-	0	-	0	-	0	-
1991–92	0	-	0	-	0	-	0	-	0	-
1992–93	0	-	0	-	0	-	0	-	< 1	-
1993–94	< 1	-	0	-	0	-	0	-	< 1	-
1994–95	0	-	0	-	0	-	0	-	0	-
1995–96	0	-	0	-	0	-	0	-	0	-
1996–97	< 1	-	0	-	0	-	< 1	-	0	-
1997–98	0	-	0	-	0	-	< 1	-	0	-
1998–99	< 1	-	0	-	0	-	0	-	0	-
1999–00	0	-	< 1	-	0	-	0	-	< 1	-
2000–01	0	-	< 1	-	0	-	0	-	< 1	-
2001–02	0	-	< 1	-	0	-	1	-	7	-
2002–03	0	-	< 1	-	0	-	< 1	-	3	-
2003–04	0	1	< 1	14	< 1	N/A	2	19	7	N/A
2004–05	0	1	< 1	14	N/A	N/A	5	19	N/A	N/A
2005–06	0	1	< 1	14	N/A	N/A	8	19	N/A	N/A
2006–07	0	1	< 1	14	N/A	N/A	5	19	N/A	N/A
2007–08	0	1	< 1	14	N/A	N/A	11	19	N/A	N/A
2008–09	< 1	1	13	14	N/A	N/A	10	19	N/A	N/A
2009–10	< 1	1	12	14	N/A	N/A	25	19	N/A	N/A
2010–11	0	1	1	14	N/A	N/A	19	19	N/A	N/A
2011–12	0	1	2	14	N/A	N/A	14	19	N/A	N/A
2012–13	< 1	1	< 1	14	N/A	N/A	54	19	N/A	N/A
2013–14	0	1	2	14	N/A	N/A	72	19	N/A	N/A
2014–15	0	1	14	14	N/A	N/A	80	19	N/A	N/A
2015–16	0	1	2.1	14	N/A	N/A	39	19	N/A	N/A
2016–17	0	1	6.4	14	N/A	N/A	48	19	N/A	N/A

GIANT SPIDER CRAB (GSC)

Table 1 [Continued].

Fishstock	GSC 6A		GSC 6B		GSC 8		GSC 10		TOTAL	
	Landings	TACC	Landings	TACC	Landings	TACC	Landings	TACC	Landings	TACC
1990-91	0	-	0	-	0	-	0	-	< 1	-
1991-92	0	-	0	-	0	-	0	-	0	-
1992-93	0	-	0	-	0	-	0	-	0	-
1993-94	0	-	0	-	0	-	0	-	1	-
1994-95	0	-	0	-	0	-	0	-	0	-
1995-96	0	-	0	-	< 1	-	0	-	< 1	-
1996-97	0	-	0	-	0	-	0	-	< 1	-
1997-98	0	-	0	-	0	-	0	-	< 1	-
1998-99	0	-	0	-	0	-	0	-	0	-
1999-00	0	-	0	-	0	-	0	-	2	-
2000-01	0	-	0	-	0	-	0	-	< 1	-
2001-02	0	-	0	-	0	-	0	-	8	-
2002-03	0	-	0	-	0	-	0	-	4	-
2003-04	0	148	0	237	0	N/A	0	0	27	419
2004-05	24	148	2	237	N/A	N/A	0	0	35	419
2005-06	63	148	1	237	N/A	N/A	0	0	72	419
2006-07	23	148	< 1	237	N/A	N/A	0	0	30	419
2007-08	16	148	2	237	N/A	N/A	0	0	29	419
2008-09	13	148	< 1	237	N/A	N/A	0	0	36	419
2009-10	44	148	3	237	N/A	N/A	0	0	84	419
2010-11	23	148	< 1	237	N/A	N/A	0	0	43	419
2011-12	83	148	< 1	237	N/A	N/A	0	0	99	419
2012-13	80	148	5	237	N/A	N/A	0	0	140	419
2013-14	52	148	<1	237	N/A	N/A	0	0	127	419
2014-15	128	148	2	237	N/A	N/A	0	0	224	419
2015-16	37	148	2	237	N/A	N/A	0	0	80	419
2016-17	132	148	<1	237	N/A	N/A	0	0	186	419





Figure 1: Reported commercial landings and TACC for GSC 5 (Southland), and GSC 6A (Southern Islands). Note that these figures do not show data prior to entry into the QMS.

1.2 Recreational fisheries

There are no known records of recreational use of this crab.

1.3 Customary non-commercial fisheries

There are no known records of customary use of this crab.

1.4 Illegal catch

There is no known illegal catch of this crab.

1.5 Other sources of mortality

There is no quantitative information on other sources of mortality, although this crab is often taken as a bycatch in orange roughy fishing.

2. BIOLOGY

Jacquiniotia is found from the intertidal to over 500 m in the southeast and south of New Zealand from near Mernoo Gap to Campbell Island. It appears to attain highest densities southeast of the Snares, on the Pukaki Rise, and around the Auckland Islands. Ryff & Voller (1976) recorded *Jacquiniotia* in highest quantities on the Pukaki Rise and at the Auckland Islands, then decreasing quantities at the Campbell Islands, Bounty Islands, Stewart Island, Stewart Island Shelf, Puysegur Bank, and off Otago Heads, an observation consistent with earlier resource surveys (Ritchie 1970, 1973; Webb 1972). At the Auckland Islands they appear to be most abundant between 20 m and 40 m, but on the Pukaki Rise between 140 m and 160 m.

This spider crab, also sometimes known as the southern spider crab or the Auckland Islands crab, is a large, conspicuous brachyuran with a brick red carapace and bright red to yellowish-white chelae. The male grows much larger than the female, to at least 20 cm across the back and, together with its up to 40 cm long clawed legs, can give a total spread approaching 1 m. The males at least seem to be migratory. There have been reports of 'mounding' behaviour associated with moulting and mating (Bennett 1964, Ritchie 1970) in which large numbers of crabs form clumps, particularly in spring and autumn.

Large males have been observed feeding on ribbed mussels (*Aulacomya maoriana*) and they probably also feed on other shellfish, both bivalves (*Mytilus*, *Macra*) and gastropods (*Haliotis*, *Maurea*, *Struthiolaria*). In contrast, females are detritus feeders on sandy substrates, and juveniles seem to feed on drift algae. These differences mean that although both males and females may enter pots, only males have been observed feeding on fish bait.

Sexes are separate and in both there appears to be a terminal moult. Males reach maturity at 110 mm carapace length (CL) and females at 100 mm CL. It appears that, at least near land masses, large males migrate between shallow and deep water seasonally. Pairs form in shallow water (less than 10 m) or just out of the water in September–November, when females are in late berry. Egg extrusion probably takes place in September to February and larval release in September to November. A female of 101 mm CL carries about 37 500 eggs; a female of 126 mm CL about 71 200 eggs. Only one batch of eggs is produced each year and the interval between hatching of one lot of eggs and extrusion of the next batch is very short. In summer, females and pre-puberty males occur mainly in shallow water while large males are found deeper.

Larval duration, survival, behaviour, and settlement are poorly known. There are two zoeal stages but the megalopa is unknown. Zoea probably occur in the plankton during September to November. Juveniles have been found in large numbers close inshore at the Auckland Islands, where shoreline rock meets the deeper mud and sand flats. Seaweed present here was apparently both food and shelter for the young crabs.

There is little or no information available on age, growth and natural mortality. Moulting appears to take place between November and March. Males reach 220 mm CL; females 144 mm. According to Ritchie (1970), *M* for mature females is 13–25%, and may be slightly higher for mature males.

GIANT SPIDER CRAB (GSC)

3. STOCKS AND AREAS

For management purposes stock boundaries are based on FMAs, however, there is currently no biological or fishery information which could be used to identify stock boundaries.

4. STOCK ASSESSMENT

4.1 Estimates of fishery parameters and abundance

There are no estimates of fishery parameters or abundance for any giant spider crab fishstock.

4.2 Biomass estimates

There are no biomass estimates for any giant spider crab fishstock.

4.3 Yield estimates and projections

There are no estimates of *MCY* for any giant spider crab fishstock.

There are no estimates of *CAY* for any giant spider crab fishstock.

5. STATUS OF THE STOCKS


There are no estimates of reference or current biomass for any giant spider crab fishstock.

6. FOR FURTHER INFORMATION

- Bennett, E W (1964) The marine fauna of New Zealand: Crustacea, Brachyura. *New Zealand Oceanographic Memoir* 22. 120 p.
- Chilton, C (1911) Scientific results of the New Zealand government trawling expedition, 1907. *Crustacea. Records of the Canterbury (New Zealand) Museum* 1: 285–312.
- McClay, C L (1988) Brachyura and crab-like Anomura of New Zealand. *Leigh Laboratory Bulletin* No. 22.
- Melville-Smith, R (1982) A brief exploitation of the stone crab *Lithodes murrayi* (Henderson) off South West Africa, 1979/80. *Fisheries Bulletin of South Africa* 16: 45–55.
- O'Driscoll, R L; Booth, J D; Bagley, N W; Anderson, O F; Griggs, L H; Stevenson, M L; Francis, M P (2001) Areas of importance for spawning, pupping or egg-laying, and juveniles of New Zealand deepwater fish, pelagic fish, and invertebrates. Final Research Report for Ministry of Fisheries Research Project ENV2000/04. Objectives 1, 2, & 3. (Unpublished report held by Fisheries New Zealand, Wellington.) 377 p.
- Paul, L J (2000) *New Zealand fishes. Identification, natural history & fisheries*. Reed. 253 p.
- Ritchie, L D (1970) Southern spider crab (*Jacquintia edwardsii* (Jacquinot, 1853)) survey - Auckland Islands and Campbell Island. *Fisheries Technical Report* No. 52.
- Ritchie, L D (1973) Commercial fishing for southern spider crab (*Jacquintia edwardsii*), at the Auckland Islands, October 1971. *Fisheries Technical Report* No. 101.
- Roberts, P E (1972) The plankton of Perseverance Harbour, Campbell Island, New Zealand. *Pacific Science* 26: 296–309.
- Ryff, M R; Voller, R W (1976) Aspects of the southern spider crab (*Jacquintia edwardsii*) fishery of southern New Zealand islands and Pukaki Rise. *Fisheries Technical Report* No. 143.
- Thomson, G M (1913) The natural history of Otago Harbour and the adjacent sea, together with a record of the researches carried out at the Portobello Marine Fish Hatchery: Part 1. *Transactions of the New Zealand Institute* 45: 225–251.
- Webb, B F (1972) Report on the investigations of the 'Lloret Lopez II' - 8 January to 2 April 1970. Section 3 Crab survey - 18 February to 27 February 1970. *Fisheries Technical Report* No. 97.
- Webber, W R; Wear, R G (1981) Life history studies on New Zealand Brachyura. 5. Larvae of the family Majidae. *New Zealand Journal of Marine and Freshwater Research* 15: 331–383.

GREEN-LIPPED MUSSEL (GLM)

(*Perna canaliculus*)
Kuku, Kutai



1. FISHERY SUMMARY

1.1 Commercial fisheries

Commercial harvesting of green-lipped mussels began with handpicking of inter-tidal beds in the late nineteenth century, and expanded in 1927 with the development of a dredge fishery for sub-tidal mussels in the Hauraki Gulf. Following a brief decline in catch rates from 1935–45, landings increased steadily to peak in 1961 at more than 2 000 t. Overexploitation of the Hauraki Gulf beds caused the fishery to close in 1966. A second dredge fishery developed in Tasman Bay and Kenepuru Sound in 1962; however, under an open access regime this fishery also declined within five years. Since 2004 reported landings have been dominated by GLM 7A and GLM 9. Total landings have been low and declining compared to the total TACC. Recent estimated landings of green-lipped mussels are shown in Table 1, while Figure 1 shows the historical landings and TACC for the three main GLM stocks.

Table 1: Reported landings (t) of Green-lipped mussel and actual TACCs (t) from 2004–05 to the present.

Fishstock (QMA)	GLM1		GLM2		GLM3		GLM7A		GLM9		Total	
	Landings	TACC	Landings	TACC	Landings	TACC	Landings	TACC	Landings	TACC	Landings	TACC
2004–05	6.2	10	0	10	0.2	10	410.9	1 500	121.3	180	539	1 720
2005–06	12.5	10	0.2	10	0.2	10	229.0	1 500	93.0	180	335	1 720
2006–07	7.8	10	0	10	0	10	84.3	1 500	136.9	180	229	1 720
2007–08	3.5	10	0	10	<0.1	10	7.4	1 500	141.7	180	153	1 720
2008–09	6.7	10	0	10	<0.1	10	0.1	1 500	67.9	180	75	1 720
2009–10	4.4	10	0	10	<0.1	10	<1	1 500	183.3	180	187	1 720
2010–11	1.0	10	0	10	0	10	1.4	1 500	78.1	180	80	1 720
2011–12	0.5	10	0	10	0	10	0.1	1 500	162.0	180	163	1 720
2012–13	0.6	10	0	10	0	10	0	1 500	129.0	180	130	1 720
2013–14	0.1	10	0	10	0	10	8.3	1 500	159.9	180	167	1 720
2014–15	<0.1	10	0	10	0	10	8.3	1 500	207.0	180	215	1 720
2015–16	0.1	10	0	10	0	10	0	1 500	203.4	180	203	1 720
2016–17	0.2	10	0	10	0	10	0	1 500	208.9	180	209	1 720

Spat collecting is the other commercial venture with green-lipped mussels. Until green-lipped mussels were introduced into the QMS a permit was required to harvest spat attached to beach cast seaweed. Green-lipped mussels were introduced into the Quota Management System on 1 October 2004 with TAC and TACC listed in Table 2.

GREEN-LIPPED MUSSEL (GLM)

Table 2: Recreational and Customary non-commercial allowances, TACC and TAC for green-lipped mussel.

Fishstock	Recreational allowance	Customary non-commercial allowance	TACC	TAC
GLM 1	162	243	10	415
GLM 2	10	15	10	35
GLM 3	58	87	10	155
GLM 7A	19	29	1 500	1 548
GLM 7B	5	8	100	23
GLM 8	17	26	0	43
GLM 9	39	59	180	278
GLM 10	0	0	0	0
Total	310	467	1 720	2 497






Figure 1: Reported commercial landings and TACC for the three main GLM stocks. From top left: GLM 1 (Auckland East), GLM 7A (Nelson Marlborough), and GLM 9 (Auckland West). Note that these figures do not show data prior to entry into the QMS.

1.2 Recreational fisheries

Recreational harvest estimates for green-lipped mussels have been obtained from the 1996, 2000 and 2001 national telephone diary surveys of recreational fishers (Table 3). Estimates of green-lipped mussels from the 1996 survey are only available for FMA 1. No weights were available from the surveys to estimate recreational harvest by tonnage. The Recreational Technical Working Group has reviewed the harvest estimates from the national telephone diary surveys and considered that the estimates from the 1996 survey are unreliable because the survey contained a methodological error. The estimated number of green-lipped mussels from the 2000 and 2001 surveys is also considered to be unreliable.

Table 3: Harvest estimates of mussels (000s of individuals of *P. canaliculus* combined) from the 1996, 2000 and 2001 national recreational surveys, by FMA (Bradford 1998, Boyd et al 2004).

FMA	1996 Harvest	2000 Harvest	2001 Harvest
1	818	1 308	949
2		8	22
3		402	187
5		1	36
7		3	363
8		242	-
9		25	148

1.3 Customary non-commercial fisheries

Green-lipped mussels are very important to customary fishing. This species was used extensively by Māori, appearing in middens throughout the country. The species continues to be important to Māori and, anecdotally, a number of customary fishers have noted its importance as a resource in a number of areas. While no information is available, the green-lipped mussel remains an important element of customary fishing throughout many parts of New Zealand.

2. BIOLOGY

The green-lipped mussel is a filter-feeding mollusc. While distributed throughout New Zealand, it is most common in central and northern parts where it frequently forms dense beds of up to 100 m². This species is absent from the Chatham Islands and other offshore islands. It is typically a bivalve of the lower shore and open coast and is found from the mid-littoral to depths of over 50 m. The species can grow to over 240 mm in shell length (anterior-posterior axis).

The green-lipped mussel is a dioecious (uni-sexual) broadcast spawner. Gonadal development takes place at temperatures above 11°C and is also related to food availability. Most spawning occurs in late spring to early autumn, but larvae can be present all year. Sexual maturity has been observed in some populations to begin from 27 mm shell length, with most individuals sexually mature by 40 mm shell length. Sexual maturity is reached in the first year, and females can produce up to 100 million eggs per season. Fertilisation is largely dependent on the proximity of adults.

Settlement processes associated with marine farms have been well studied, but less is known about natural settlement. The planktonic stage (pediveligers) of the green-lipped mussel is ready to settle at 220–350 µm in length, after a three to five week larval phase. The larvae swim only vertically but they can be transported large distances by currents and tides. Settlement is most intense from late winter to early summer, but is highly variable spatially and temporally. In the wild, larvae settle over a wide range of depths, preferring fine filamentous substrata including hydroids, bryozoans, and filamentous and turfing algae. Settlement is completed with the attachment of byssus threads and subsequent metamorphosis.

Primary settlement onto beds of adult mussels is uncommon, but can take place on surrounding algae and on the byssi of adults. Secondary settlement, after a form of byssopelagic migration or mucous drifting, is thought to be the means by which most juveniles recruit into mussel beds. The spat detaches from the substrate by severing the byssus threads and the secreted mucous strand, this enables it to swim or drift to new areas for attachment. Juvenile mussels may move numerous times like this before settling on adult mussel beds. This drifting ability is lost once spat reach about 6 mm in shell length.

GREEN-LIPPED MUSSEL (GLM)

There is little information on age, growth and natural mortality, particularly for wild populations. Green-lipped mussels in suspended culture typically grow from 10 to 75 mm shell length in six months, to 111–115 mm in one year, and to 195 mm in three and a half years. Growth is typically faster in cultured situations compared with natural beds, which are often overcrowded, are on exposed coasts, and are not constantly submerged so feeding is discontinuous. At Piha and West Tamaki Head, green-lipped mussel growth is variable, with individuals reaching 20–70 mm shell length in their first year.

3. STOCKS AND AREAS

Green-lipped mussels are distributed in seven of the ten FMAs (1–3, 5 and 7–9) but are most common in the central and northern parts of New Zealand.

There is little information on stock structure, recruitment patterns, or other biological characteristics. There appears to be strong genetic structuring of the New Zealand green-lipped mussel population, with a northern and southern group being differentiated by frequency shifts in common haplotypes, and the occurrence of a unique haplotype in the south island west coast population. The southern-northern population split occurs south of Cook Strait.

4. STOCK ASSESSMENT

There are no stock assessments or biomass estimates for green-lipped mussels.

5. STATUS OF THE STOCKS


No estimates of reference or current biomass are available for any green-lipped mussel fishstock. It is not known whether green-lipped mussel stocks are at, above, or below a level that can produce *MSY*.

6. FOR FURTHER INFORMATION

- Alfaro, A C; Jeffs, A G; Hooker, S H (2001) Reproductive behavior of the green-lipped mussel, *Perna canaliculus*, in northern New Zealand. *Bulletin of Marine Science* 69(3):1095–1108.
- Apte, S; Star, B; Gardner, J (2003) A comparison of genetic diversity between cultured and wild populations, and a test for genetic introgression in the New Zealand greenshell mussel *Perna canaliculus* (Gmelin 1791). *Aquaculture* 219: 193–220.
- Bradford, E (1998) Harvest estimates from the 1996 national recreational fishing surveys. New Zealand Fisheries Assessment Research Document. 1998/16. 27 p. (Unpublished document held by NIWA library, Wellington.)
- Boyd, R O; Gowing, L; Reilly, J L (2004) 2000–2001 National marine recreational fishing survey: diary results and harvest estimates. New Zealand Fisheries Research Report. (Unpublished report held by Fisheries New Zealand, Wellington.)
- Boyd, R O; Reilly, J L (2004) 1999/2000 National marine recreational fishing survey: harvest estimates. New Zealand Fisheries Research Report. (Unpublished report held by the Fisheries New Zealand, Wellington.)
- Greenway, J P C (1969) Surveys of mussels (Mollusca: Lamellibranches) of Thames, 1961–67. *New Zealand Journal of Marine and Freshwater Research* 3: 304–17.

GREY MULLET (GMU)

(*Mugil cephalus*)
Kanae, Hopuhopu



1. FISHERY SUMMARY

1.1 Commercial fisheries

Commercial fishing for grey mullet occurs predominantly in GMU 1, where annual landings increased from approximately 128 t in 1931 to a maximum of 1 142 t in 1983–84 (Table 1; 2). Marked changes in fishing effort occurred during this period through the development of more efficient fishing techniques and an increase in the market demand for this species. Before the introduction of the QMS, total domestic catches declined from the maximum (1 160 t) in 1983–84 to 901 t in 1985–86. The TACC was consistently under caught after GMU 1 was introduced into the QMS (Figure 1). The Minister of Fisheries therefore reduced the TACC for GMU 1 to 925 t, beginning in 1998–99. The reduction in TACC had little effect on the annual catches, and it has only ever been reached in GMU 1 in 2004–05 and 2013–14 (Table 2).




Figure 1: Reported commercial landings and TACC for the main GMU stock; GMU 1 (Auckland).

GREY MULLET (GMU)

Table 1: Reported landings (t) for the main QMAs from 1931 to 1990.

Year	GMU 1	GMU 2	GMU 3	GMU 7	Year	GMU 1	GMU 2	GMU 3	GMU7
1931–32	128	0	0	0	1957	204	1	0	0
1932–33	138	0	0	0	1958	262	0	0	0
1933–34	78	0	0	0	1959	244	0	0	0
1934–35	111	0	0	0	1960	213	0	0	0
1935–36	147	0	0	0	1961	230	0	0	0
1936–37	80	0	0	0	1962	191	0	0	0
1937–38	82	0	0	0	1963	199	0	0	0
1938–39	117	1	0	1	1964	214	0	0	0
1939–40	91	0	0	0	1965	222	2	3	0
1940–41	77	0	0	0	1966	240	0	0	0
1941–42	48	2	0	0	1967	243	0	0	0
1942–43	44	2	0	0	1968	256	0	0	0
1943–44	35	0	0	0	1969	283	1	1	0
1944	104	0	0	0	1970	248	1	0	0
1945	138	0	0	0	1971	253	1	0	0
1946	141	0	0	0	1972	305	0	1	0
1947	151	0	0	0	1973	393	1	4	2
1948	114	0	0	0	1974	386	0	0	0
1949	100	0	0	0	1975	360	0	0	0
1950	129	0	0	0	1976	394	0	0	0
1951	108	0	0	0	1977	557	0	0	0
1952	136	0	0	0	1978	604	0	0	0
1953	166	0	0	0	1979	735	0	0	0
1954	190	0	0	0	1980	494	0	0	0
1955	188	0	0	0	1981	612	0	0	0
1956	193	0	0	0	1982	990	0	8	2

Notes:

1. The 1931–1943 years are April–March but from 1944 onwards are calendar years.
2. Data up to 1985 are from fishing returns; Data from 1986 to 1990 are from Quota Management Reports.
3. Data for the period 1931 to 1982 are based on reported landings by harbour and are likely to be underestimated as a result of under-reporting and discarding practices. Data includes both foreign and domestic landings.

Table 2: Reported landings (t) of grey mullet by Fishstock from 1983–84 to 2016–17 and actual TACCs (t) for 1986–87 to 2016–17. QMS data from 1986-present. There have been no report landings for GMU 10.

Fishstock QMA (s)	GMU 1 1 & 9		GMU 2 2 & 8		GMU 3 3, 4, 5 & 6		GMU 7 7		GMU 10 10	Total	
	Landings	TACC	Landings	TACC	Landings	TACC	Landings	TACC	TACC	Landings	TACC
1983–84*	1 142	-	6	-	5	-	7	-	-	1 160	-
1984–85*	1 069	-	5	-	0	-	15	-	-	1 089	-
1985–86*	881	-	10	-	0	-	10	-	-	901	-
1986–87	595	910	3	20	<1	30	0	20	10	598	990
1987–88	751	941	3	20	0	30	0	20	10	754	1 021
1988–89	792	963	3	20	0	30	0	20	10	795	1 043
1989–90	907	990	2	20	0	30	4	20	10	913	1 070
1990–91	875	994	2	20	1	30	<1	20	10	879	1 073
1991–92	848	1 006	1	20	2	30	1	20	10	852	1 086
1992–93	711	1 006	<1	20	<1	30	0	20	10	712	1 086
1993–94	743	1 006	<1	20	<1	30	0	20	10	706	1 086
1994–95	776	1 006	0	20	<1	30	10	20	10	787	1 086
1995–96	866	1 006	0	20	<1	30	<1	20	10	866	1 086
1996–97	870	1 006	<1	20	1	30	<1	20	10	872	1 086
1997–98	730	1 006	<1	20	<1	30	<1	20	10	730	1 086
1998–99	750	925	<1	20	<1	30	<1	20	10	750	1 005
1999–00	749	925	<1	20	0	30	<1	20	10	750	1 005
2000–01	797	925	1	20	0	30	<1	20	10	798	1 005
2001–02	782	926	2	20	<1	30	<1	20	10	784	1 005
2002–03	797	926	1	20	<1	30	0	20	10	798	1 005
2003–04	886	926	<1	20	0	30	<1	20	10	796	1 005
2004–05	941	926	<1	20	0	30	0	20	10	941	1 005
2005–06	878	926	<1	20	<1	30	0	20	10	878	1 005
2006–07	847	926	1	20	0	30	<1	20	10	845	1 005
2007–08	848	926	1	20	<1	30	<1	20	10	849	1 005
2008–09	814	926	1	20	0	30	0	20	10	815	1 005
2009–10	746	926	<1	20	0	30	0	20	10	746	1 005
2010–11	825	926	<1	20	<1	30	<1	20	10	826	1 006
2011–12	848	926	<1	20	<1	30	<1	20	10	848	1 006
2012–13	871	926	<1	20	<1	30	<1	20	10	871	1 006
2013–14	981	926	<1	20	0	30	0	20	10	981	1 006
2014–15	900	926	<1	20	0	30	<1	20	10	901	1 006
2015–16	827	926	<1	20	0	30	0	20	10	827	1 006
2016–17	835	926	<1	20	0	30	0	20	10	836	1 006

*FSU data.

1.2 Recreational fisheries

Grey mullet are a popular recreational species particularly in the Auckland FMA. Information is available on the relative levels of commercial and amateur catch of this species in the Manukau Harbour and the lower Waikato River based on limited tagging work undertaken in 1987. Of the number of tags returned 38% were from amateur fishers, suggesting that recreational use of the resource was relatively high.

Telephone-diary surveys in 1993–94 (Teirney et al 1997), 1996 (Bradford 1998), and 2000 (Boyd et al 2004) were used to estimate the annual recreational catch from GMU 1 as 150, 106, and 100 t, respectively (Table 3). The Minister of Fisheries provided an allowance for customary harvest of 100 t beginning in 1998–99.

The harvest estimates provided by telephone-diary surveys between 1993 and 2001 are no longer considered reliable for various reasons. A Recreational Technical Working Group concluded that these harvest estimates should be used only with the following qualifications: a) they may be very inaccurate; b) the 1996 and earlier surveys contain a methodological error; and c) the 2000 and 2001 estimates are implausibly high for many important fisheries. In response to these problems and the cost and scale challenges associated with onsite methods, a National Panel Survey was conducted for the first time throughout the 2011–12 fishing year. The panel survey used face-to-face interviews of a random sample of 30 390 New Zealand households to recruit a panel of fishers and non-fishers for a full year. The panel members were contacted regularly about their fishing activities and harvest information collected in standardised phone interviews. Harvest estimates for grey mullet are given in Table 3 using a mean weight of 0.922 kg per fish (from Wynne-Jones et al. 2014 and Hartill & Davey 2015).

Table 3: Estimated number of grey mullet harvested by recreational fishers by Fishstock and survey year, the corresponding estimated survey harvest, and the estimated Fishstock harvest.

Survey	Fishstock	Number	CV	Harvest range (t)	Harvest estimate (t)
1994 telephone-diary	GMU 1	170 000	0.19	90–210	150
1996 telephone-diary	GMU 1	110 000	0.25	80–130	106
2000 telephone-diary	GMU 1	110 000	0.33	68–136	102
2011-12 panel survey	GMU 1	29 622	0.41	-	27.3
2011-12 panel survey	GMU 2	1 531	0.53	-	2.8
2011-12 panel survey	GMU 3	5 252	0.93	-	4.8
2011-12 panel survey	GMU 7	191	0.73	-	0.2
2011-12 panel survey	GMU total	38 127	0.34	-	35.2

A repeat of the National Panel Survey is being conducted over the 2017–18 October fishing year. Results are expected in early 2019.

1.3 Customary non-commercial fisheries

No quantitative information is available on the current level of customary non-commercial take. The Minister of Fisheries provided an allowance for customary harvest of 100 t per annum beginning in 1998–99.

1.4 Illegal catch

Estimates of illegal catch are unknown but anecdotal evidence suggests 10–20% under-reporting is plausible. In the latest stock assessment, an annual under-reporting of 20% was assumed for the period before 1986 and 10% thereafter.

1.5 Other sources of mortality

No quantitative estimates are available regarding the impact of other sources of mortality on grey mullet stocks. Grey mullet principally occur in sheltered harbours and estuarine ecosystems. Some of these habitats are known to have suffered environmental degradation.

2. BIOLOGY

Grey mullet has a worldwide distribution, occurring commonly along coasts, in estuaries, and in lower river systems between latitudes of 42° N and 42° S. Overseas and New Zealand tagging studies indicate that movement patterns of adult grey mullet are complex. Some schools remain in one locality, while others appear to be on the move almost continuously. Recorded movements of tagged grey mullet of 160 km within a few weeks of release are not uncommon.

Females grow faster than males and attain a larger size. Both sexes mature at 3 years of age at an average size of 33 cm fork length (FL) for males and 35 cm FL for females. Maximum ages appear to be 12 to 14 years, with ages 4–8 making up the bulk of the commercial fishery.

Natural mortality was estimated from the equation $M = \log_e 100/\text{maximum age}$, where maximum age is the age to which 1% of the population survives in an unexploited stock. Using 15 years for the maximum age results in an estimate of $M = 0.33$. (Note: the maximum age of 15 years was obtained from an exploited population, so M is likely to be less than 0.33).

Grey mullet commonly occur in schools, which generally become larger and more prevalent in the spawning season. Spawning in northern New Zealand occurs during November to February. Females are highly fecund and may release up to 1 million eggs in a spawning event. It is likely that grey mullet spawn at sea, because running-ripe females have only been caught off coastal beaches or in offshore waters, and eggs and larvae are a component of the offshore coastal plankton at certain times of the year. Small post-larval grey mullet occur seasonally in estuaries, which serve as nursery grounds for juveniles.

Adult grey mullet typically feed on diatom algae and small invertebrates which are gulped along with surface scum or with detrital ooze and sifted by fine teeth and gill-rakers.

Biological parameters relevant to stock assessment are shown in Table 4.

Table 4: Estimates of biological parameters of grey mullet.

Fishstock	Estimate						Source
1. Natural mortality (<i>M</i>)							
GMU 1	0.33						NIWA (unpubl. data)
2. Weight = a(length) ^b (Weight in g, length in cm fork length).							
	Both Sexes						
	a	b					
GMU 1	0.04236	2.826					Breen & McKenzie (unpublished)
3. Von Bertalanffy growth parameters							
	Females			Males			
	<i>L</i> _∞	<i>k</i>	<i>t</i> ₀	<i>L</i> _∞	<i>k</i>	<i>t</i> ₀	
GMU 1	40.1	0.587	1.3469	37.0	0.619	1.3257	Breen & McKenzie (unpublished)

3. STOCKS AND AREAS

There is little biological data to determine the level of sub stock separation within GMU 1. Results from a small scale tagging program in the Manukau Harbour and the Lower Waikato River indicated that there is fish movement between these two localities and also north along the west coast but the net level of movement cannot be ascertained. There is evidence in the CPUE data that GMU 1 may be comprised of six populations with low to moderate mixing between them (McKenzie 1997).

GMU 1 has been divided into two sub-stocks (east coast and west coast) for the purposes of fisheries stock assessment. The boundary between the two sub-stocks is assumed to be due north from North Cape.

4. STOCK ASSESSMENT

4.1 Estimates of fishery parameters and abundance

Standardised CPUE analyses were undertaken for the six largest catching areas in GMU 1. The analysis was based on setnet catch and effort data for the years 1990–91 to 2005–06 (McKenzie & Vaughan 2008), and updated to 2010–11 (Kendrick & Bentley 2012). However, internal and anecdotal evidence suggest that method is being misreported in these fisheries and that standardized CPUE is unlikely to reflect relative abundance for GMU. CPUE was therefore rejected as an index of relative abundance for all sub-areas within GMU 1.

4.2 Biomass estimates

West coast GMU 1

A stock assessment was undertaken for the west GMU 1 substock using a stochastic dynamic age-structured observation-error time series model (Breen & McKenzie 1998), but this did not prove to be robust and the results were rejected by the Working Group.

4.3 Yield estimates and projections

There is insufficient information with which to revise the yield estimates of either the West or East coast GMU 1 substocks. The *MCY* estimate derived in 1986 using the equation $MCY = cY_{AV}$ (Method 4) remains the accepted yield estimate for GMU 1.

Annual landings of grey mullet in the Auckland QMA for the period 1974–84 showed an increasing trend to a maximum in 1984. There were some fluctuations throughout this period. A general increase in fishing effort occurred during this time. Fishing effort between 1983–84 and 1985–86 appeared relatively constant, and catches during these years were averaged to estimate Y_{AV} . The constant ' c ' was set at 0.8. This is not consistent with the maximum observed age of 14 years, which equates with an estimate of $M = 0.33$ and $c = 0.7$. However, it is believed that they live to older ages in unexploited populations. Therefore, the accuracy of *MCY* derived for grey mullet is uncertain. The estimate of *MCY* for GMU 1 is shown in Table 5. *MCY* cannot be estimated for the other fish stocks.

Table 5: Estimate of *MCY* (t) rounded to the nearest 5 t.

Fishstock	QMA	Y_{AV}	<i>MCY</i>
GMU 1	Auckland 1 & 9	1 030	825

The level of risk to the stock by harvesting the population at the estimated *MCY* level cannot be determined. No estimates of current biomass, fishing mortality, or other information are available which would permit the estimation of *CAY*.

4.5 Other Factors

The minimum legal mesh size for use in the grey mullet fishery is 89 mm. However, fishers typically use mesh larger than 89 mm when fishing for grey mullet (Fisheries New Zealand data). There are no data available to compare the selectivity characteristics of different mesh sizes. It is possible that a significant fraction of the grey mullet stock comprising larger older fish is poorly selected by the fishery. If this is true then the von Bertalanffy parameter estimates, which are based on random samples from the 1997–98 setnet landings, are likely to be biased: L_{∞} will be biased low, K biased high.

Grey mullet have been exploited by customary, commercial, and recreational fishers for over a hundred years. They are found predominantly in harbours and these environments have undergone considerable change over this period due to a range of anthropogenic sources. The impact of these changes on potential carrying capacity and productivity are not understood and this potentially has impacts on the yields of GMU.

Characterisation shows an overall trend away from set netting towards ring netting, and, within the nominal setnet method, a trend towards shorter nets; a trend that is not seen in flatfish setnet fisheries in the same areas. This suggests there have been systematic changes in fishing strategy that are not

GREY MULLET (GMU)

captured by the CELR form. Anecdotal information from interviews of net fishers suggests that fishers use the various net method codes interchangeably, and that the methods describe differences in strategy rather than in gear, from passive fishing to spotting and encircling schools of fish. While the passive form of set netting is an appropriate sampling tool, any contamination by ring net or similarly 'directed' fishing could mask trends in the abundance of the underlying population.

The Working Group agreed that given the misreporting issues and its consequences, that standardized CPUE is unlikely to reflect relative abundance for GMU.

5. STATUS OF THE STOCKS

Given the misreporting of method and its consequences, standardized CPUE is unlikely to reflect relative abundance for GMU. CPUE was therefore rejected as an index of relative abundance for all sub-areas within GMU 1.

Yields, TACCs and reported landings are summarised in Table 6.

Table 6: Summary of yields (t), TACCs (t), and reported landings (t) of grey mullet for the most recent fishing year.


Fishstock	QMA	MCY	2016–17 Actual TACC	2016–17 Reported landings
GMU 1	Auckland (East) (West) 1 & 9	825	926	835
GMU 2	Central (East) (West) 2 & 8	-	20	< 1
GMU 3	South-East (Coast) (Chatham) 3, 4,			
	Southland and Sub-Antarctic 5 & 6	-	30	0
GMU 7	Challenger 7	-	20	0
GMU 10	Kermadec 10	-	10	0
Total		-	1 006	836

6. FOR FURTHER INFORMATION

- Annala, J H; Sullivan, K J; O'Brien, C J (1999) Report from the Fishery Assessment Plenary, April 1999: Stock assessments and yield estimates. (Unpublished report held by Fisheries New Zealand, Wellington.) 430 p.
- Anon (1989) Effects of commercial fishing on the fisheries of the Manukau Harbour and lower Waikato River. (Unpublished internal report held by NIWA library.)
- Boyd, R O; Gowing, L; Reilly, J L (2004) 2000–2001 national marine recreational fishing survey: diary results and harvest estimates. (Draft New Zealand Fisheries Assessment Report held by Fisheries New Zealand Wellington.) 93 p.
- Boyd, R O; Reilly, J L (2005) 1999/2000 National marine recreational fishing survey: harvest estimates. Final Research Report for Ministry of Fisheries Research Project REC9803. (Unpublished report held by Fisheries New Zealand, Wellington.)
- Bradford, E (1998) Harvest estimates from the 1996 national recreational fishing surveys. New Zealand Fisheries Assessment Research Document 1998/16. 27 p. (Unpublished report held by Fisheries New Zealand, Wellington.)
- Breen, P A; McKenzie, J R (1998) A simple age-structured model for grey mullet (*Mugil cephalus*) stock assessment.
- Hartill, B (2004) Characterisation of the commercial flatfish, grey mullet, and rig fisheries in the Kaipara Harbour. *New Zealand Fisheries Assessment Report 2004/1*. 23 p.
- Hartill, B; Davey, N (2015) Mean weight estimates for recreational fisheries in 2011–12. *New Zealand Fisheries Assessment Report 2015/25*.
- Hore, A (1985) Grey mullet. In: Colman, J A; McKoy, J L; Baird, G G (1985) Background papers for the 1985 Total Allowable Catch recommendations, pp. 82–85. (Unpublished report held at NIWA library, Wellington.) 259 p.
- Kendrick, T H; Bentley, N (2012) Fishery characterisation and setnet catch-per-unit-effort indices for grey mullet in GMU 1, 1989–90 to 2010–11. Final Research Report for project INS2011-01. (Unpublished report held by Fisheries New Zealand, Wellington.)
- King, M R (1985) Fish and shellfish landings by domestic fishermen, 1974–1982. *Fisheries Research Division Occasional Publication*: 20.
- McKenzie, J (1997) Catch per Unit Effort analysis of the Northern (GMU 1) target grey mullet (*Mugil cephalus*) setnet fishery 1983–96. Draft Fisheries Assessment Research Document
- McKenzie, J; Paul, L; Ó Maolagáin, C (1999) Length and age composition of commercial grey mullet landings from the west coast setnet fishery (GMU 1), 1997–98. Final Research Report for Ministry of Fisheries project GMU9701. (Unpublished report held by Fisheries New Zealand, Wellington.)
- McKenzie, J; Vaughan, M (2008) CPUE and characterisation of grey mullet (*Mugil cephalus*) setnet fisheries in GMU 1 between 1989 and 2006. *New Zealand Fisheries Assessment Report 2008/57*. 37 p.
- Teirney, L D; Kilner, A R; Millar, R E; Bradford, E; Bell, J D (1997) Estimation of recreational catch from 1991–92 to 1993–94. New Zealand Fisheries Assessment Research Document 1997/15. 43 p. (Unpublished document held by NIWA library, Wellington.)
- Watson, T; McKenzie, J; Hartill, B (2005) Catch Per Unit Effort Analysis of the Northern (GMU 1) Grey Mullet (*Mugil cephalus*) Setnet Fishery 1989–2002. *New Zealand Fisheries Assessment Report 2005/22*. 36 p.
- Wells, R D S (1976) The utilisation of the lower Waikato Basin by the grey mullet *Mugil cephalus*. (Unpublished MSc thesis, University of Waikato).
- Wynne-Jones, J; Gray, A; Hill, L; Heinemann, A (2014) National Panel Survey of Marine Recreational Fishers 2011–12: Harvest Estimates. *New Zealand Fisheries Assessment Report 2014/67*.

GROPER (HPB)*(Polyprion oxygeneios, Polyprion americanus)*

Hapuku, Moeone

**1. FISHERY SUMMARY****1.1 Commercial fisheries**

Both groper species, *Polyprion oxygeneios* (hapuku) and *P. americanus* (bass), occur in shelf and slope waters of the New Zealand mainland and offshore islands, from the Kermadecs to the Auckland Islands. The groper fishery takes both species, but in different proportions by region, depth, fishing method and season, and these have changed over time. Reported catches generally do not distinguish between species, and published data combine them. In earlier years, bluenose (*Hyperoglyphe antarctica*) landings were sometimes also combined with groper. In this document, groper is used as collective term for hapuku and bass. Historical estimated and recent reported groper landings and TACCs are shown in Tables 2 and 3, while Figure 1 shows the historical and recent landings and TACC values for the main groper stocks.

Table 1: Reported total New Zealand landings (t) of groper from 1948 to 1983.

Year	Landings	Year	Landings	Year	Landings	Year	Landings
1948	1 665	1957	1 368	1966	1 222	1975	1 422
1949	1 969	1958	1 532	1967	1 314	1976	1 512
1950	1 709	1959	1 310	1968	1 073	1977	1 942
1951	1 396	1960	1 223	1969	1 122	1978	1 488
1952	1 430	1961	1 203	1970	1 499	1979	2 078
1953	1 403	1962	1 173	1971	1 346	1980	2 435
1954	1 364	1963	1 194	1972	1 120	1981	2 379
1955	1 305	1964	1 370	1973	1 312	1982	2 218
1956	1 399	1965	1 249	1974	1 393	1983	2 511

Reported foreign catches are included from 1974.

Source: Fisheries data.

The main fishery comprises a number of domestic fishers working small to medium sized vessels - longliners, setnetters and trawlers, at a variety of depths (according to method) out to 500 m (Paul 2002a). Over 90% of early (to 1950) total groper catches were taken by longline. Trawl catches rose from 5–10% during this period to 20–30% by the late 1970s. A setnet fishery developed in the late 1970s and early 1980s, mainly at Kaikoura, taking 14% in 1983 and then subsequently declining. From 1950 to the mid-1980s, line-fishing took 70–80% of the catch. After the introduction of the QMS in 1986, the proportion of the catch taken by lines appeared to drop.

GROPER (HPB)

The Cook Strait region has always supported the main groper fishery, followed by the Canterbury Bight; both show the same slow decline from 1949 to 1986 (equivalent regional data from subsequent years are not available). Northland, Bay of Plenty and Hawke Bay fisheries developed at different rates during the 1960s and 1970s. In most other areas, the groper fishery has been small and/or variable.

The first recorded landings of about 1 500 t in 1936 were typical of the range of catches (1 000–2 000 t) from then until 1978. After a decrease during the war when effort was restricted, landings in the total fishery slowly declined from almost 2 000 t in 1949 to about 1 300 t in the mid-1970s. They then increased sharply to 2 700 t in 1983–84 (Tables 1 and 2). Figure 1 shows the historical landings and TACC values for the main HPB stocks.

Landings and TACCs for all Fishstocks are given in Table 3. Total landings of groper were relatively stable throughout the mid-1990s, remaining below 1 500 t until 1998–99. From 1999–2000 onwards, catches have generally ranged between 1 200 t and 1 700 t. Although the TACC in HPB 3 has been exceeded in some years, catches have generally remained within the quotas for individual Fishstocks and have never exceeded the TACC.

For the 1991–92 fishing year the conversion factor for headed and gutted groper was increased from 1.40 to 1.45, for fish landed in this state (about 75% of the total), this will result in a reduction in removals from the stock of 3.5% for the same nominal quota.

Table 2: Reported landings (t) for the main QMAs from 1931 to 1982.

Year	HPB 1	HPB 2	HPB 3	HPB 4	Year	HPB 1	HPB 2	HPB 3	HPB 4
1931–32	231	0	207	2	1957	133	380	419	23
1932–33	201	276	242	0	1958	115	473	458	30
1933–34	198	330	173	25	1959	147	406	350	54
1934–35	204	304	212	57	1960	122	394	331	48
1935–36	179	201	146	70	1961	135	369	348	50
1936–37	129	445	115	12	1962	163	355	298	40
1937–38	119	523	315	15	1963	197	315	321	56
1938–39	90	621	479	8	1964	224	397	365	41
1939–40	118	502	409	12	1965	212	368	325	68
1940–41	120	444	286	9	1966	213	415	315	4
1941–42	80	450	302	10	1967	229	448	275	0
1942–43	69	287	315	9	1968	139	357	264	0
1943–44	59	316	271	8	1969	197	454	220	0
1944	55	332	286	9	1970	259	670	239	2
1945	106	311	271	3	1971	191	562	289	4
1946	154	326	409	7	1972	401	370	188	0
1947	98	401	563	5	1973	419	481	215	0
1948	111	450	526	11	1974	356	457	208	2
1949	174	498	547	7	1975	227	315	213	18
1950	141	423	555	9	1976	183	220	350	107
1951	104	353	381	19	1977	277	301	265	87
1952	112	368	373	35	1978	348	470	194	10
1953	105	349	431	33	1979	620	487	355	147
1954	156	355	397	32	1980	956	376	414	40
1955	142	351	419	26	1981	693	373	457	59
1956	106	404	439	32	1982	957	336	402	26

Year	HPB 5	HPB 7	HPB 8	Year	HPB 5	HPB 7	HPB 8
1931–32	130	13	13	1957	92	246	76
1932–33	91	98	53	1958	96	250	109
1933–34	99	127	53	1959	68	198	87
1934–35	115	106	56	1960	100	150	77
1935–36	33	109	33	1961	82	139	80
1936–37	29	156	50	1962	101	142	75
1937–38	29	148	52	1963	75	159	71
1938–39	75	156	50	1964	76	193	74
1939–40	59	155	43	1965	48	176	52
1940–41	54	142	41	1966	49	163	62
1941–42	46	150	44	1967	49	228	85

Table 2 [Continued]

Year	HPB 5	HPB 7	HPB 8	Year	HPB 5	HPB 7	HPB 8
1942–43	44	115	35	1968	67	176	70
1943–44	42	112	42	1969	30	138	84
1944	60	188	117	1970	54	175	97
1945	65	173	128	1971	41	181	78
1946	83	229	190	1972	29	99	33
1947	142	250	175	1973	30	136	32
1948	140	275	151	1974	43	140	72
1949	142	364	236	1975	55	379	62
1950	116	281	184	1976	101	445	37
1951	102	267	171	1977	47	575	113
1952	100	281	162	1978	59	280	67
1953	96	252	137	1979	113	276	71
1954	77	235	112	1980	199	315	105
1955	82	197	88	1981	218	381	166
1956	114	227	77	1982	133	256	46

Notes:

1. The 1931–1943 years are April–March but from 1944 onwards are calendar years.
2. Data up to 1985 are from fishing returns: Data from 1986 to 1990 are from Quota Management Reports.
3. Data for the period 1931 to 1982 are based on reported landings by harbour and are likely to be underestimated as a result of under-reporting and discarding practices. Data includes both foreign and domestic landings.

Table 3: Reported landings (t) of groper by Fishstock from 1983–84 to present and actual TACCs (t) from 1986–87 to present. QMS data from 1986–present. [Continued on next page].

Fishstock FMA (s)	HPB 1 1 & 9		HPB 2 2		HPB 3 3		HPB 4 4		HPB 5 5 & 6	
	Landings	TACC	Landings	TACC	Landings	TACC	Landings	TACC	Landings	TACC
1983–84*	974	-	493	-	505	-	55	-	395	-
1984–85*	642	-	388	-	418	-	52	-	228	-
1985–86*	569	-	270	-	391	-	53	-	126	-
1986–87	238	360	179	210	260	270	42	300	131	410
1987–88	248	388	202	219	268	286	43	315	91	414
1988–89	231	405	187	248	259	294	49	315	70	425
1989–90	310	465	179	263	283	318	40	322	127	430
1990–91	350	480	225	263	311	326	77	323	120	436
1991–92	277	480	252	263	298	326	58	323	112	446
1992–93	375	480	273	264	299	327	68	323	128	446
1993–94	363	480	287	264	306	330	90	323	147	446
1994–95	334	481	259	264	274	335	149	323	161	451
1995–96	335	481	214	264	321	335	173	323	144	451
1996–97	331	481	234	264	301	335	131	323	149	451
1997–98	375	481	260	266	329	335	88	323	91	451
1998–99	433	481	256	266	348	335	121	323	97	451
1999–00	471	481	229	266	385	335	66	323	169	451
2000–01	450	481	220	266	381	335	45	323	188	451
2001–02	427	481	226	266	343	335	82	323	169	451
2002–03	442	481	273	266	350	335	79	323	212	451
2003–04	433	481	281	266	335	335	87	323	166	451
2004–05	433	481	263	266	371	335	147	323	208	451
2005–06	425	481	280	266	406	335	185	323	167	451
2006–07	483	481	245	266	394	335	222	323	157	451
2007–08	439	481	253	266	341	335	241	323	138	451
2008–09	415	481	253	266	391	335	138	323	153	451
2009–10	374	481	249	266	358	335	213	323	152	451
2010–11	371	481	222	266	322	335	231	323	128	451
2011–12	312	481	193	266	336	335	265	323	158	451
2012–13	314	481	206	266	337	335	156	323	140	451
2013–14	319	481	224	266	301	335	169	323	143	451
2014–15	314	481	180	266	280	335	156	323	126	451
2015–16	270	481	143	266	315	335	144	323	143	451
2016–17	287	481	162	266	342	335	152	323	156	451

	HPB 7 7		HPB 8 8		HPB 10 10		Total	
	Landings	TACC	Landings	TACC	Landings	TACC	Landings	TACC
1983–84*	174	-	46	-	0	-	2 698	-
1984–85*	207	-	33	-	0	-	2 039	-
1985–86*	199	-	25	-	0	-	1 697	-
1986–87	149	210	35	60	0	10	1 036	1 830
1987–88	158	215	66	76	0	10	1 076	1 923
1988–89	132	226	39	78	1	10	968	2 001
1989–90	119	229	43	80	0	10	1 098	2 117
1990–91	128	235	48	80	23#	10	1 282	2 153
1991–92	175	235	50	80	83#	10	1 319	2 163
1992–93	186	236	62	80	22#	10	1 405	2 165
1993–94	193	236	69	80	0	10	1 455	2 167

GROPER (HPB)

Table 3 [Continued]

	HPB 7		HPB 8		HPB 10		Total	
	Landings	TACC	Landings	TACC	Landings	TACC	Landings	TACC
1994-95	192	236	68	80	0	10	1 437	2 179
1995-96	214	236	78	80	0	10	1 479	2 179
1996-97	186	236	71	80	15	10	1 418	2 179
1997-98	147	236	60	80	33#	10	1 406	2 181
1998-99	218	236	78	80	3#	10	1 562	2 181
1999-00	165	236	65	80	0#	10	1 561	2 181
2000-01	171	236	64	80	0#	10	1 519	2 181
2001-02	204	236	62	80	< 1	10	1 514	2 181
2002-03	233	236	72	80	0	10	1 661	2 181
2003-04	239	236	66	80	0	10	1 607	2 181
2004-05	240	236	80	80	0	10	1 742	2 181
2005-06	207	236	56	80	0	10	1 728	2 181
2006-07	206	236	66	80	0	10	1 773	2 181
2007-08	195	236	44	80	0	10	1 651	2 181
2008-09	207	236	71	80	0	10	1 628	2 181
2009-10	221	236	66	80	0	10	1 633	2 181
2010-11	191	236	80	80	0	10	1 543	2 181
2011-12	173	236	61	80	0	10	1 187	2 181
2012-13	209	236	75	80	0	10	1 436	2 181
2013-14	182	236	63	80	0	10	1 401	2 181
2014-15	132	236	67	80	0	10	1 254	2 181
2015-16	148	236	73	80	0	10	1 236	2 181
2016-17	141	236	69	80	0	10	1 309	2 181

* FSU data.

Values in HPB 10 included catches taken under exploratory permit.




Figure 1: Total reported landings and TACC for the seven main HPB stocks. From top to bottom: HPB 1 (Auckland) and HPB 2 (Central East) [Continued on the next page].




Figure 1 [Continued]: Total reported landings and TACC for the seven main HPB stocks. From top to bottom: HPB 3 (South East Coast), HPB 4 (Chatham Rise), and HPB 5 (Southland, Sub-Antarctic). [Continued on next page].

GROPER (HPB)




Figure 1 [Continued]: Total reported landings and TACC for the seven main HPB stocks. From top to bottom: HPB 7 (Challenger) and HPB 8 (Central).

1.2 Recreational fisheries

Groper are taken by handline and setline, and to a lesser extent by setnets. Recreational catch estimates from surveys undertaken in the 1990s are given in Tables 4–6.

Table 4: Estimated number of groper harvested by recreational fishers by Fishstock and survey, the corresponding estimated survey harvest and the estimated Fishstock harvest. Surveys were carried out in different years in the MAF Fisheries regions: South in 1991–92, Central in 1992–93 and North in 1993–94 (Teirney et al 1997).

Fishstock	Survey	Total		Survey harvest (t)
		Number	CV (%)	
HPB 1	North	22 000	17	190–220
HPB 2	North	1 000	-	5–10
HPB 2	Central	10 000	37	45–85
HPB 3	Central	3 000	-	10–30
HPB 3	South	4 000	40	10–30
HPB 5	Central	7 000	36	20–40
HPB 5	South	2 000	-	5–15
HPB 7	Central	12 000	40	45–115
HPB 8	Central	1 000	-	5–10

Table 5: Results of a national diary survey of recreational fishers in 1996, indicating estimated number of groper harvested by recreational fishers by Fishstock and the corresponding harvest tonnage. The mean weights used to convert numbers to catch weight are considered the best available estimates. Estimated harvest is also presented as a range to reflect the uncertainty in the estimates (from Bradford 1998).

Fishstock	Number caught	CV (%)	Harvest range (t)	Point Estimate (t)
HPB 1	11 000	17	40–60	49
HPB 2	23 000	22	75–125	100
HPB 3	4 000	-	-	-
HPB 5	2 000	-	-	-
HPB 7	9 000	-	-	-
HPB 8	< 500	-	-	-

Table 6: Results of the 1999–2000 national diary survey of recreational fishers (Dec 1999–Nov 2000). Estimated number of groper harvested by recreational fishers by Fishstock, and the corresponding harvest tonnage. Estimated harvest is presented as a range to reflect the uncertainty in the estimates (Boyd & Reilly 2002).

Fishstock	Number caught	CV (%)	Harvest range (t)	Point estimate (t)
HPB 1	60 000	39	209–476	342
HPB 2	56 000	33	307–608	457
HPB 3	52 000	50	97–293	195
HPB 5	6 000	70	14–80	47
HPB 7	17 000	37	79–172	125
HPB 8	2 000	67	6–32	19

The harvest estimates provided by telephone-diary surveys between 1993 and 2001 are no longer considered reliable for various reasons. A Recreational Technical Working Group concluded that these harvest estimates should be used only with the following qualifications: a) they may be very inaccurate; b) the 1996 and earlier surveys contain a methodological error; and c) the 2000 and 2001 estimates are implausibly high for many important fisheries. In response to these problems and the cost and scale challenges associated with onsite methods, a National Panel Survey was conducted for the first time throughout the 2011–12 fishing year. The panel survey used face-to-face interviews of a random sample of 30 390 New Zealand households to recruit a panel of fishers and non-fishers for a full year. The panel members were contacted regularly about their fishing activities and harvest information collected in standardised phone interviews. Harvest estimates for groper are given in Table 7 (from Wynne-Jones et al 2014 and Hartill & Davey 2015).

Table 7: Recreational harvest estimates for groper stocks (Wynne-Jones et al 2014). Mean fish weights were obtained from boat ramp surveys; for groper the value used was 5.854 kg (Hartill & Davey 2015).

Stock	Year	Method	Number of fish	Total weight (t)	CV
HPB 1	2011/12	Panel survey	14 264	83.5	0.37
HPB 2	2011/12	Panel survey	10 179	59.6	0.28
HPB 3	2011/12	Panel survey	6 383	37.4	0.31
HPB 5	2011/12	Panel survey	138	0.8	1.00
HPB 7	2011/12	Panel survey	2 163	12.7	0.41
HPB 8	2011/12	Panel survey	4 376	25.6	0.54
HPB total	2011/12	Panel survey	37 502	219.5	0.18

A repeat of the National Panel Survey is being conducted over the 2017–18 October fishing year. Results are expected in early 2019.

1.3 Customary non-commercial fisheries

Groper (hapuku and bass) were certainly taken by early Maori, and would have been available in greater numbers at shallower depths than is the case at present. Traditional groper grounds are known in several regions. Quantitative information on the current level of customary non-commercial catch is not available.

1.4 Illegal catch

Quantitative information on the level of illegal catch is not available.

1.5 Other sources of mortality

None are apparent.

2. BIOLOGY

Both hapuku and bass are widely distributed around New Zealand, generally over rough ground from the central shelf (about 100 m) to the shelf edge and down the upper slope. Their lower limits are ill-defined, but hapuku extends to at least 300 m and bass to 500 m.

Hapuku mature sexually between 10 and 13 years old and may live in excess of 60 years (Francis et al 1999). Cook Strait hapuku mature over a wide size range, with the size at 50% maturity at 80–85 cm total length (TL) and 85–90 cm TL for males and females respectively (Paul 2002d). Spawning occurs during winter, anecdotally earlier in the north of New Zealand than in the south, but running ripe fish are seldom caught and spawning grounds are unknown. The smallest juveniles are virtually unknown, but are mottled, pelagic and epi-pelagic, perhaps schooling in association with drifting weed.

The size range of commercially caught hapuku is 50–140 cm TL, with a broad mode between 70 and 100 cm TL. Bass are slightly larger at 60–150 cm TL, with a mode at 80–110 cm TL, but much bulkier and heavier at equivalent lengths.

There appear to be some regional differences in the size structure of populations. Trawl-caught hapuku on the Stewart-Snares Shelf are mainly 50–80 cm, modal length 60 cm, and therefore juveniles. Trawl-caught hapuku on the Chatham Rise are slightly larger, 50–100 cm, modal length 70 cm, with those on the shelf around the islands having their main mode at 60–75 cm; most of these fish are also juveniles. These offshore regions may be important nurseries.

Both groper species are assumed to be long-lived. Natural mortality in the past was assumed to be 0.2, however, a study of a South American (Juan Fernandez) population suggested that it may be lower (0.13–0.16) (Pavez & Oyarzun 1985). Furthermore, preliminary unvalidated ageing in New Zealand has indicated that maximum age may be greater than 40 years, and that M may be 0.1 or less (Francis et al 1999). This value of M will be retained until clearer information becomes available from ageing. Parker et al (2011) compared regional differences in the catch composition from observer collected data. This report noted that the proportion of age 10+ fish in the catch in the Kermadec and Northeastern regions (FMA 2) was greater than that of Southland.

Migration patterns are also little known, but are probably related to spawning. Tagging of mostly immature fish in Cook Strait has shown a high level of site fidelity, but about 5% of these fish have moved up to 160 km north and south. Other information is largely anecdotal and speculative. It is known that good fishing grounds, particularly pinnacles and reefs or ledges, can be quickly fished out and take some time to recover, suggesting a high level of residency (except, perhaps, for during the spawning season). On the other hand, trawlers sometimes catch groper on the flat and clear seafloor, and it is not known whether this represents their normal habitat, whether they are simply dispersing by travelling from one rough ground to another, or whether they are on a purposeful spawning migration.

Hapuku and bass prey on a wide variety of fish and invertebrates, including red cod, tarakihi, blue cod, hoki and squid. In Cook Strait, they are preyed upon by sperm whales, although probably neither heavily nor selectively. Biological parameters relevant to stock assessment are shown in Table 8.

Table 8: Estimates of biological parameters of groper.

Fishstock	Estimate	Source
1. Natural mortality (M)		
All	$M = 0.1$	Francis et al (1999)
2. Weight = a (length) ^b (Weight in g, length in cm fork length)		
	Both sexes combined	
BAS 1	$a = 0.2734$ $b = 2.382$	Johnston (1993)
HAP 1	$a = 0.0142$ $b = 3.003$	Johnston (1993)
HAP 2	$a = 0.0242$ $b = 2.867$	Johnston (1993)
HAP 7, 8	$a = 0.0142$ $b = 2.998$	Johnston (1983)

(HAP = hapuku, BAS = bass groper)

3. STOCKS AND AREAS

Tagging studies reveal considerable mixing of hapuku between Otago, South Canterbury and Cook Strait. Fishstock boundaries in Cook Strait separate Cook Strait hapuku into three separate "stocks" (HPB 2, HPB 7, and HPB 8), none of which include Otago-Canterbury fish (HPB 3). Current Fishstock boundaries appear inappropriate for the management of Cook Strait and South Island hapuku. Current stock boundaries are based on QMAs and do not reflect biological stocks. Existing data cannot describe the stock structure of New Zealand groper (Paul 2002b). Electrophoretic studies suggest that separate stocks of hapuku could occur. However, the genetic heterogeneity of Cook Strait hapuku, seasonal movements of hapuku through this area, moderately long-distance movements of some tagged hapuku, the presence of both species on open ground and the eventual recovery of heavily exploited reefs, suggest that either each stock is moderately mobile or that there is essentially only one stock (of each species) with some small geographic or temporal genetic differences.

4. STOCK ASSESSMENT

Yield estimates for HPB 4 and HPB 5 have been removed because the previous method used is now considered obsolete. The yield estimates for the other Fishstocks have been revised based on a revision of the estimate of M .

4.1 Estimates of fishery parameters and abundance

Estimates of fishery parameters and abundance are not available. Paul (2002c) found that CPUE indices could not be developed for hapuku and bass either separately or in combination.

4.2 Biomass estimates

Estimates of current and reference biomass are not available. Data for hapuku from the East Coast South Island trawl surveys have moderate CVs (average over all years = 28.17; range 19–35) and although the survey does not extend to the entire habitat range, the survey may be monitoring settled juveniles (Figure 2).

4.4 Yield estimates and projections

Current biomass cannot be estimated, so CAY cannot be determined. Yield estimates are summarised in Table 9.

Table 9: Yield estimates (t).

Parameter	Fishstock	Estimate
	HPB 4	Cannot be determined
	HPB 5	Cannot be determined
	Total	Cannot be determined
CAY	All	Cannot be determined

GROPER (HPB)




Figure 2: Biomass estimates $\pm 95\%$ CI (estimated from survey CV's assuming a lognormal distribution) and the time series mean (dotted line) from the East Coast South Island trawl survey.

4.5 Other factors

Although no distinct stocks of either groper species have been identified, results from trawl surveys suggest that there are reasonably large but dispersed populations over the Stewart-Snares Shelf and the Chatham Rise. The relationship between these "offshore" and the more traditionally fished "inshore" populations is not known due to the lack of information on groper movements. Little is known of the species composition and population structure of groper on the rough bottom shelf and ridges extending northwards from New Zealand.

The relative quantity of groper taken as target and non-target catch has not been investigated, but is likely to have varied both spatially and temporally. Groper have been taken by the foreign licensed, chartered and New Zealand-owned trawlers working offshore grounds; although being regarded as a small bycatch they were not accurately reported before 1986. The *MCY* may therefore be underestimated.

There are three regions where the groper catch has been substantially lower than the TACC.

HPB 1 - Three features of the fishery appear to explain the under-catch of the TACC. (i) A considerable part of the fishing effort which had generated the high catches in the early 1980s left the fishery. (ii) The allocated quota is widely distributed in small units among fishers who appear to use only a modest proportion of it to cover bycatch. (iii) The fishers who hold larger amounts of quota generally also use only a proportion of it to land high-quality fish (in contrast to the earlier bulk landings of lower-quality fish).

HPB 4 and 5 - The original yield estimates made before the introduction of the QMS and the original TAC were based on trawl surveys, not catch histories. The TACCs for these Fishstocks can only be economically targeted around the Chatham Islands in HPB 4, and a few localities in HPB 5. Elsewhere, it is used to cover a small bycatch from trawlers. A moderate quantity of quota is held, unused, by companies which would require it should they resume target fishing for ling and associated species.

5. STATUS OF THE STOCKS

No estimates of current biomass are available. An estimate of B_{AV} is available for HPB 5.

It is not known if current catches or the TACCs are sustainable or at levels that will allow the stocks to move towards a size that will support the maximum sustainable yield.

Yield estimates, TACCs and reported landings are summarised in Table 10.

Table 10: Summary of TACCs (t) and reported landings (t) of groper for the most recent fishing year.

Fishstock	QMA	FMAs	2016–17	2016–17
			Actual TACC	Reported Landings
HPB 1	Auckland (East, West)	1 & 9	481	287
HPB 2	Central (East)	2	266	162
HPB 3	South-east (Coast)	3	335	342
HPB 4	South-east (Chatham)	4	323	152
HPB 5	Southland, Sub-Antarctic	5 & 6	451	156
HPB 7	Challenger	7	236	141
HPB 8	Central (West)	8	80	69
HPB 10	Kermadec	10	10	0
Total			2 181	1 309

6. FOR FURTHER INFORMATION

- Beentjes, M P; Francis, M P (1999) Movements of hapuku, *Polyprion oxygeneios* determined from tagging studies. *New Zealand Journal of Marine and Freshwater Research* 33(1): 1–12.
- Boyd, R O; Reilly, J L (2002) 1999/2000 national marine recreational fishing survey: harvest estimates. Draft New Zealand Fisheries Assessment Report. (Unpublished report held by Fisheries New Zealand.)
- Bradford, E (1998) Harvest estimates from the 1996 national recreational fishing surveys. New Zealand Fisheries Assessment Research Document. 1998/16. 27 p. (Unpublished report held by Fisheries New Zealand.)
- Francis, M P; Mulligan, K P; Davies, N M; Beentjes, M P (1999) Age and growth estimates for New Zealand hapuku, *Polyprion oxygeneios*. *Fishery Bulletin*. 97(2): 227–242.
- Hartill, B; Davey, N (2015) Mean weight estimates for recreational fisheries in 2011–12. *New Zealand Fisheries Assessment Report* 2015/25.
- Hurst, R J; Bagley, N W; Uozumi, Y (1990) New Zealand-Japan trawl survey of shelf and upper slope species off southern New Zealand, June 1986. *New Zealand Fisheries Technical Report* No. 18. 50 p.
- Johnston, A D (1983) The southern Cook Strait groper fishery. *Fisheries Technical Report* No. 159. 33 p.
- Johnston, R G (Ed.) (1993) Report from the Conversion Factors Working Group and Steering Committee 1992. MAF Fisheries, Greta Point Internal Report No. 201. 171 p. (Unpublished report held in NIWA library, Wellington.)
- McDougall, C R (1975) Age and growth of *Polyprion oxygeneios* (Pisces: Serranidae) in Cook Strait. (Unpublished B.Sc. (Hons) thesis (Zoology), Victoria University of Wellington.)
- Parker, S J; Paul, L J; Francis, M P (2011) Age structure characteristics of hapuku *Polyprion oxygeneios* stocks from existing samples of otoliths. *New Zealand Fisheries Assessment Report* 2011/31. 42 p.
- Paul, L (2002a) A description of the New Zealand fisheries for the two groper species, hapuku (*Polyprion oxygeneios*) and bass (*P. americanus*). *New Zealand Fisheries Assessment Report* 2002/13. 47 p.
- Paul, L (2002b) Can existing data describe the stock structure of the two New Zealand groper species, hapuku (*Polyprion oxygeneios*) and bass (*P. americanus*)? *New Zealand Fisheries Assessment Report* 2002/14. 24 p.
- Paul, L (2002c) Can separate CPUE indices be developed for the two groper species, hapuku (*Polyprion oxygeneios*) and bass (*P. americanus*)? *New Zealand Fisheries Assessment Report* 2002/15. 24 p.
- Paul, L (2002d) Size structure of hapuku (*Polyprion oxygeneios*) and bass (*P. americanus*) populations in New Zealand. *New Zealand Fisheries Assessment Report* 2002/16. 17 p.
- Paul, L J (1985) The estimation of hapuku and bass yields for New Zealand fishing regions. Fisheries Research Division Internal Report No. 26. 31 p. (Unpublished report held in NIWA library, Wellington.)
- Paul, L J; Davies, N M (1988) Groper. New Zealand Fisheries Assessment Research Document 88/15. 27 p. (Unpublished report held by NIWA library, Wellington.)
- Pavez, P; Oyarzun, M E (1985) Determination of the relative efficiency of hooks, and growth parameters of the Juan Fernandez "cod" *Polyprion oxygeneios* Bloch and Schneider, 1801, in the Robinson Crusoe and Santa Clara Islands. In Arana, P (Ed.), "Investigaciones en el Archipelago de Juan Fernandez", pp. 341–345. Escuela de Ciencias del Mar, UCV, Valparaiso. [In Spanish, English summary.]
- Teirney, L D; Kilner, A R; Millar, R E; Bradford, E; Bell, J D (1997) Estimation of recreational catch from 1991/92 to 1993/94 New Zealand Fisheries Assessment Research Document 1997/15. 43 p. (Unpublished report held by NIWA library, Wellington.)
- Teirney, L; McKinnon, S; Kilner, A; Sylvester, T (1991) Marine Recreational Fisheries Working Group Report — November 1991. New Zealand Fisheries Working Group Report 91/1. 46 p.
- Teirney, L D; Olsen, D L (1992) Marine Recreational Fisheries Group Report — November 1992. New Zealand Fisheries Working Group Report 92/1. 13 p.
- Wynne-Jones, J; Gray, A; Hill, L; Heinemann, A (2014) National Panel Survey of Marine Recreational Fishers 2011–12: Harvest Estimates. *New Zealand Fisheries Assessment Report* 2014/67.

Fisheries Assessment Plenary

May 2018

Stock Assessment and Stock Status
Volume 1: Introductory Section to Groper

ISBN: 978-1-77665-886-2 (online)
ISBN: 978-1-77665-887-9 (print)

Fisheries New Zealand
Tini a Tangaroa
PO Box 2526, Wellington, 6140
New Zealand
www.mpi.govt.nz



Fisheries New Zealand
Tini a Tangaroa

New Zealand Government